

CONCISE REPORT

# Hemin-Induced Conversion of Pyruvate Kinase Isozymes in K562 Cells

By S. Takegawa, T. Shinohara, and S. Miwa

The effects of hemin on the conversion of pyruvate kinase (PK) isozymes from the M<sub>2</sub>-type to the L-type in K562 cells were investigated. Immunofluorescence, ion exchange chromatography, and electrophoretic studies showed that the untreated K562 cells contained only the M<sub>2</sub>-type PK, while eight to 20 days after induction with hemin, concomi-

tant with hemoglobin F synthesis, L-type PK levels increased while M<sub>2</sub>-type PK levels decreased. Electrophoretic study revealed three hybrid isozymes of the L-type and M<sub>2</sub>-type PK. We conclude that the conversion of PK isozymes from the M<sub>2</sub>-type to the L-type in erythroid precursor cells occurs in the early stage of maturation.

**WE** HAVE RECENTLY REPORTED the presence of M<sub>2</sub>-type pyruvate kinase (PK) in the proerythroblast stage and conversion of the M<sub>2</sub>-type

PK to L-type PK with erythroid maturation<sup>1</sup> and demonstrated a compensatory production of M<sub>2</sub>-type PK in the orthochromatic erythroblasts of PK-deficient patients.<sup>2</sup> Lozzio and Lozzio<sup>3</sup> found that the K562 cell line obtained from the pleural effusion of a patient with Philadelphia chromosome-positive chronic myelogenous leukemia (CML) in blastic crisis can be differentiated into erythroid precursors after the addition of hemin.<sup>4</sup> Hence, this cell line seems to be suitable for examining changes in PK isozymes in erythroid precursor cells. In this report, we describe conversion of PK isozymes from the M<sub>2</sub>-type to the L-type in K562 cells after exposure to an inducing agent, hemin.

*From the Department of Internal Medicine, The Institute of Medical Science, The University of Tokyo, and the Department of Human Genetics, Japan Red Cross Medical Center, Tokyo.*

*Supported in part by scientific research grants from the Ministry of Education and the Ministry of Health and Welfare, Japan.*

*Submitted April 27, 1984; accepted May 4, 1984.*

*Address reprint requests to Dr Susumu Takegawa, Department of Internal Medicine, The Institute of Medical Science, The University of Tokyo, 4-6-1, Shirokanedai Minato-ku, Tokyo 108, Japan.*

© 1984 by Grune & Stratton, Inc.

0006-4971/84/6403-0026\$03.00/0





Fig 1. Karyotype of a pseudotriploid cell with 46 chromosomes stained with the Giemsa-banding technique. The Philadelphia chromosome (Ph<sup>1</sup>) is identified by the arrow. \*, abnormal chromosomes; M<sub>1-5</sub>, markers.



**Fig 2.** Immunofluorescent stain of K-562 cells. K-562 cells reacted with FITC-labeled anti-M<sub>2</sub>-type PK (M<sub>2</sub>), anti-L-type PK (L), and anti-hemoglobin F (Hb F). HEMIN(-), untreated K-562 cells; HEMIN(+) 8d, cells on day 8 of induction; HEMIN(+) 20d, cells on 20th day of induction.


**MATERIALS AND METHODS**

The K562 cells were a kind gift of Dr Yoshikura at the Department of Bacteriology, The University of Tokyo. Chromosome analysis was performed with the trypsin-Giemsa banding technique. The cultures contained 10<sup>5</sup> cells per milliliter in RPMI 1640 medium (GIBCO, Grand Island, NY) supplemented with 10% fetal calf serum with or without 50 μmol/L hemin (Sigma Chemical Co, St Louis).<sup>4</sup> After 20 days the cells were harvested by centrifugation at 4 °C. Lysis was performed by addition of four volumes of 10 mmol/L sodium phosphate buffer, pH 6.0, containing 2 mmol/L MgSO<sub>4</sub>, 0.2 mmol/L fructose-1,6-diphosphate (FDP) and 0.1% 2-mercaptoethanol (buffer A) and by sonication for two minutes. The cellular debris was removed by centrifugation at 4 °C for 30 minutes at 20,000 g.

Pyruvate kinase activities were determined at 37 °C according to the International Committee for Standardization in Haematology<sup>5</sup> (ICSH) recommended methods<sup>5</sup> using a Gilford Model 250 spectrophotometer. The column was loaded with 0.08 to 0.1 U of the enzyme preparation. For linear gradient elution, a CM-Sephadex C-50 (Pharmacia Fine Chemicals, Piscataway, NJ) column (1 × 18 cm) was equilibrated with buffer A. A 70-mL linear gradient of sodium phosphate buffer from 10 to 100 mmol/L was used for elution; 60 fractions were collected. All the operations were carried

out at 4 °C. Pyruvate kinase electrophoresis was performed using polyacrylamide thin layer gel.<sup>7</sup>

Detection of L-type and M<sub>2</sub>-type PK by the immunofluorescent method was described previously.<sup>1</sup> Hemoglobin F was determined by the immunofluorescent technique described by Wood et al.<sup>6</sup>



**Fig 3.** Change of PK in K-562 cell. CM-Sephadex column chromatography. Abbreviations used are the same as those in Fig 2.

## RESULTS

*Cytogenetic Studies*

The K562 cells were aneuploid with a prominent, near-triploid mode (66 to 72). The karyotypic analysis revealed that all the cells contained at least seven abnormal chromosomes and three to five markers (Fig 1). The Philadelphia chromosome was identified, but the 9q+ chromosome was not observed. The banding pattern showed that the 17q+ chromosome was probably a translocation between one chromosome 15 and one chromosome 17.

*Immunofluorescent Studies*

Figure 2 shows the results of the immunofluorescent studies. The fluorescence of  $M_2$ -type PK was decreased after eight days of hemin induction, while that of L-type PK and hemoglobin F was increased with induction duration.

*Ion Exchange Chromatography*

The chromatographic patterns are shown in Fig 3. Purified L-type PK was eluted at an earlier position than the  $M_2$ -type PK (Fig 3A). In the sample from uninduced K562 cells, the elution profiles showed a single peak at the same position as  $M_2$ -type PK. On the other hand, in hemin-induced cells, a broad fraction was seen with a peak in a position between L- and  $M_2$ -type PK (Fig 3D).

*Pyruvate Kinase Electrophoresis*

Electrophoretic studies revealed three bands in addition to the L-type PK band, the mobilities of which were between L-type and  $M_2$ -type PK (Fig 4).

## DISCUSSION

We have reported the conversion of PK isozymes from the  $M_2$ -type to the L-type during early maturation of normal and PK-deficient erythroblasts.<sup>1,2</sup> K562 cells established by Lozzio and Lozzio<sup>3</sup> are known to differentiate into erythroid cells after the addition of hemin.<sup>4</sup> Many studies concerning embryonic hemoglobin synthesis and membrane proteins have been reported, but little is known about the glycolytic enzymes of this cell line.




Fig 4. Pyruvate kinase electrophoresis. H, hemolysate;  $M_2$ ,  $M_2$ -type PK; L, L-type PK.

Our immunofluorescent studies and CM-Sephadex chromatographic study confirm that conversion of PK isozymes occurs by induction with hemin, ie,  $M_2$ -type PK decreased and L-type PK increased. Furthermore, electrophoretic studies showed hybrid enzymes between  $M_2$  and L. Cardenas et al<sup>8</sup> detected five isozymes of bovine PK by using the hybridization technique. We applied a similar technique to 20-day samples, but the zymogram showed only four bands (data not shown). There appears to be two reasons for this result. First, the ratio of  $M_2$ -type PK to L-type PK was not equal. Second, the condition of dissociation of PK subunits was not appropriate, and homotetramer isozyme, ie,  $(M_2)_4$  could not form.

Recently, Jansen et al<sup>9</sup> showed the presence of L-type PK in K562 cells, but they failed to demonstrate the conversion of PK isozymes,<sup>9</sup> possibly because of conditions such as the batches of serum used and concentration of the inducers, which are of great importance in cell cultures.<sup>10</sup> Furthermore, the K562 cells used by Jansen et al<sup>9</sup> exhibited different karyotypic characteristics from our K562 cells.

## REFERENCES

1. Takegawa S, Fujii H, Miwa S: Change of pyruvate kinase isozymes from  $M_2$ - to L-type during development of the red cell. *Br J Haematol* 54:467, 1983
2. Takegawa S, Miwa S: Change of pyruvate kinase (PK) isozymes in classical type PK deficiency and other PK deficiency cases during red cell maturation. *Am J Hematol* 16:53, 1984
3. Lozzio CB, Lozzio BB: Human chronic myelogenous leukemia cell-line with positive Philadelphia chromosome. *Blood* 45:321, 1975
4. Rutherford TR, Clegg JB, Weatherall DJ: K562 human leukaemia cells synthesize embryonic haemoglobin in response to haemin. *Nature* 280:164, 1979
5. Beutler E, Blume KG, Kaplan JC, Löhr GW, Ramot B, Valentine WN: International committee for standardization in

haematology: Recommended methods for red-cell enzyme analysis. *Br J Haematol* 35:331, 1977

6. Wood WG, Stamatoyannopoulos G, Lim G, Nute PE: F-cells in the adult: Normal values and levels in individuals with hereditary and acquired elevation of Hb F. *Blood* 46:671, 1975

7. Miwa S, Boivin P, Blume KG, Arnold H, Black JA, Kahn A, Staal GEJ, Nakashima K, Tanaka KR, Paglia DE, Valentine WN, Yoshida A, Beutler E: International committee for standardization in haematology: Recommended methods for the characterization of

red cell pyruvate kinase variants. *Br J Haematol* 43:275, 1979

8. Cardenas JM, Dyson RD: Bovine pyruvate kinase. II. Purification of the liver isozyme and its hybridization with skeletal muscle pyruvate kinase. *J Biol Chem* 248:6938, 1973

9. Jansen G, Rijkse G, deGast GC, Staal GEJ: Glycolytic enzymes of an erythroleukemic cell line, K562, before and after hemoglobin induction. *Exp Hematol* 11:626, 1983

10. Harrison PR: New human myeloid leukaemia cell line undergoes red shift. *Nature* 281:632, 1979