

The Role of Tumor Cell–Derived Connective Tissue Growth Factor (CTGF/CCN2) in Pancreatic Tumor Growth

Kevin L. Bennewith,¹ Xin Huang,¹ Christine M. Ham,^{1,2} Edward E. Graves,¹ Janine T. Erler,¹ Neeraja Kambham,³ Jonathan Feazell,¹ George P. Yang,² Albert Koong,¹ and Amato J. Giaccia¹

Departments of ¹Radiation Oncology, ²Surgery, and ³Pathology, Stanford University School of Medicine, Stanford, California

Abstract

Pancreatic cancer is highly aggressive and refractory to existing therapies. Connective tissue growth factor (CTGF/CCN2) is a fibrosis-related gene that is thought to play a role in pancreatic tumor progression. However, CCN2 can be expressed in a variety of cell types, and the contribution of CCN2 derived from either tumor cells or stromal cells as it affects the growth of pancreatic tumors is unknown. Using genetic inhibition of CCN2, we have discovered that CCN2 derived from tumor cells is a critical regulator of pancreatic tumor growth. Pancreatic tumor cells derived from CCN2 shRNA-expressing clones showed dramatically reduced growth in soft agar and when implanted s.c. We also observed a role for CCN2 in the growth of pancreatic tumors implanted orthotopically, with tumor volume measurements obtained by positron emission tomography imaging. Mechanistically, CCN2 protects cells from hypoxia-mediated apoptosis, providing an *in vivo* selection for tumor cells that express high levels of CCN2. We found that CCN2 expression and secretion was increased in hypoxic pancreatic tumor cells *in vitro*, and we observed colocalization of CCN2 and hypoxia in pancreatic tumor xenografts and clinical pancreatic adenocarcinomas. Furthermore, we found increased CCN2 staining in clinical pancreatic tumor tissue relative to stromal cells surrounding the tumor, supporting our assertion that tumor cell–derived CCN2 is important for pancreatic tumor growth. Taken together, these data improve our understanding of the mechanisms responsible for pancreatic tumor growth and progression, and also indicate that CCN2 produced by tumor cells represents a viable therapeutic target for the treatment of pancreatic cancer. [Cancer Res 2009;69(3):775–84]

Introduction

Pancreatic cancer is the tenth most common cancer type diagnosed in the United States each year, and is the fourth most common cause of cancer deaths. The 5-year survival rate for patients with pancreatic ductal adenocarcinoma is ~5%, and has not substantially improved over the past 25 years (1). There is a clear need for increased understanding of the mechanisms driving pancreatic cancer growth and progression so that novel therapeutic strategies can be devised to improve treatment. As with most

solid tumors, pancreatic ductal adenocarcinomas contain tumor cells that are at low oxygen tensions (hypoxic), and tumor hypoxia is known to induce the expression of a variety of genes associated with tumor progression and aggressiveness. Pancreatic tumors also contain extensive desmoplasia, and this fibrotic tissue can store a variety of secreted factors that facilitate tumor progression (2). Genes such as connective tissue growth factor are thought to play a role in the formation of desmoplastic tissue and also in tumor progression and, therefore, represent an attractive therapeutic target for the treatment of pancreatic cancer.

Connective tissue growth factor (CTGF/CCN2) is a member of the CCN family of proteins and is thought to be involved in extracellular matrix production, desmoplasia, tumor cell proliferation, adhesion, migration, angiogenesis, and metastasis (3, 4). The myriad of functions assigned to CCN2 may be partly explained by the modular domain structure of the protein. CCN2 has four structural domains, each of which is thought to have a distinct biological function (5), and cleaved versions of CCN2 have been detected in body fluids (6). CCN2 is known to interact with transforming growth factor- β (TGF- β) and bone morphogenetic protein-4 to modulate their activity (7), and also binds a variety of integrins involved in cell adhesion and migration (8–10). Thus, CCN2 function in different cell types is likely dictated by a number of factors including CCN2 expression levels, structure of CCN2 molecules, and interaction with other proteins.

Although the role of CCN2 in normal tissue fibrosis has been well-studied (11), the function of CCN2 in cancer is not as well-understood. Interestingly, CCN2 has been identified as an oncogene in a variety of cancer types but is considered a tumor-suppressor gene in other forms of cancer. Overexpression of CCN2 correlates with decreased survival in patients with esophageal adenocarcinoma (12), glioblastoma (13), breast cancer (14), gastric cancer (15), and adult acute lymphoblastic leukemia (16). Increased CCN2 expression has been associated with progression of cervical tumors (17), esophageal squamous cell carcinoma (18), and Wilms' tumor (19). Conversely, CCN2 expression levels correlated with increased survival in chondrosarcoma patients (20) and in patients with lung cancer (21). Whether these disparities in the literature are due to diversity in CCN2 structure, expression levels, and binding partners in different tumor types is an open question. Notably, CCN2 expression has been observed in tumor cells, tumor-associated fibroblasts, and endothelial cells, raising the question that the effects of CCN2 on tumor progression may be influenced by the source of CCN2 production.

Previous studies have shown that immunologic inhibition of CCN2 delays the growth and metastasis of xenografted human pancreatic tumors (22, 23). However, the CCN2 antibody used in these studies cross-reacts with both human CCN2 (tumor cell–derived) and mouse CCN2 (stromal cell–derived), not permitting insight into the effect of tumor cell or stromal cell sources of CCN2

Note: Supplementary data for this article are available at Cancer Research Online (<http://cancerres.aacrjournals.org/>).

X. Huang and C.M. Ham contributed equally to this work.

Requests for reprints: Amato J. Giaccia, CCSR-South, Room 1255, 269 Campus Drive, Stanford, CA 94305. Phone: 650-723-7366; Fax: 650-723-7382; E-mail: giaccia@stanford.edu.

©2009 American Association for Cancer Research.
doi:10.1158/0008-5472.CAN-08-0987

on pancreatic tumor growth. Determining the relative importance of tumor cell-derived CCN2 or stromal cell-derived CCN2 in pancreatic tumor growth would improve our understanding of pancreatic cancer progression, and would also provide a means to select patients that are more likely to benefit from immunologic CCN2 inhibition.

We show herein that pancreatic tumor growth in both s.c. and orthotopic sites is critically dependent on CCN2 expressed by tumor cells, and that there is a robust *in vivo* selection for tumor cells that express high levels of CCN2. We observed elevated levels of CCN2 in clinical pancreatic adenocarcinomas when compared with either normal pancreatic tissue or to stromal cells surrounding the tumor. We also found that CCN2 colocalized with tumor hypoxia in clinical pancreatic adenocarcinomas and in human tumor xenografts, and that CCN2 secretion was increased in hypoxic pancreatic tumor cells *in vitro*. CCN2 increased the growth of pancreatic tumor cells in soft agar and decreased apoptosis of pancreatic tumor cells in response to hypoxic stress *in vitro*, potentially explaining the *in vivo* selection for tumor cells that express high levels of CCN2. Taken together, these data indicate the importance of tumor cell-derived CCN2 in pancreatic tumor growth and support the inhibition of CCN2 in clinical pancreatic cancer therapy.

Materials and Methods

Patient samples. Human tissue was obtained from pancreatic cancer patients undergoing pancreaticoduodenectomies at Stanford Hospital; informed consent was obtained before the procedure under the approval of the Stanford Institutional Review Board. Paraffin-embedded samples were stained as previously described for CCN2 (22) or carbonic anhydrase-IX (CAIX; ref. 24). CCN2 staining intensity was scored on a scale from 0+ to 3+ (where 0+ = no staining and 3+ = strong staining), and the fraction of tissue with a given level of CCN2 staining intensity was estimated. The overall CCN2 staining scores were based on the most intense staining found in a given section, such that a staining intensity of 3+ in >50% of tumor tissue were 3, a staining intensity of 2+ in >50% of tumor tissue were 2, and a staining intensity of 1+ in >50% of tumor tissue were 1. In sections that contained lesser amounts of stained tissue, the overall CCN2 staining scores were based on a staining intensity of 3+ in <50% of tumor tissue were 2, a staining intensity of 2+ in <50% of tumor tissue were 1, and a staining intensity of 0 to 1+ in <50% of tumor tissue were 0.

CAIX staining intensities were typically classified as either 3+ or 0+, and the overall CAIX staining scores were derived from the relative tissue area stained with 3+ CAIX intensity: 3+ staining in >50% of tumor tissue were 3, 3+ staining in <50% and >15% of tumor tissue were 2, 3+ staining in <15% of tumor tissue were 1, and no observable 3+ staining were 0.

Tumor cells. Human pancreatic Panc-1 and Su86.86 tumor cells were obtained from the American Type Culture Collection and used within 10 passages. Panc-1 cells were maintained in DMEM supplemented with 10% fetal bovine serum (FBS), whereas Su86.86 cells were maintained in RPMI 1640 supplemented with 10 mmol/L HEPES, 1 mmol/L sodium pyruvate, and 10% FBS. CCN2 shRNAs were designed using the Dharmacon shRNA design algorithm: A, 5'-CTATGATTAGACCAACTG-3'; B, 5'-GCTGACCTG-GAAGAGAACA-3'; C, 5'-GAGAACATTAAGAAGGGCA-3'; D, 5'-TGA-CATCTTTGAATCGCTG-3'; E, 5'-TCGCTGTACTACAGGAAGA-3'; and F, 5'-GATGTACGGAGACATGGCA-3'. Oligonucleotides were inserted into pSiren vector for retroviral transfection and CCN2 shRNA-expressing cells were puromycin selected. Surviving cells were either pooled or single clones were picked and expanded to make clonal cell populations. CCN2 in the pBabe Puro vector was used to induce CCN2 expression.

For monolayer growth curves, 10^5 cells were plated in 60-mm plates and grown for 3 to 5 d. Cells were trypsinized, counted using a Coulter Z1 particle counter (Beckman Coulter, Inc.), and 10^5 cells were replated and allowed to grow for another 3 to 5 d. For soft agar growth, 0.5% Noble agar

(BD Biosciences) was allowed to solidify in a 12-well plate, and 5×10^3 cells were plated in 0.3% Noble agar on top. Tumor cell colonies were stained with 0.02% Geimsa stain in PBS after 14 to 21 d.

Subcutaneous and orthotopic tumor implants. All animal studies were performed in accordance with the Stanford University Animal Care and Use Committee. For subcutaneous tumor growth, male 8- to 10-wk-old Nu/Nu mice were anesthetized with 2% isoflurane in O_2 , and 10^7 Panc-1 or Su86.86 tumor cells were implanted in 100 μ L of PBS. Tumor volumes were calculated as the volume of an ellipsoid based on three orthogonal caliper measurements ($L \times W \times H \times \pi/6$).

For orthotopic tumor growth, male 8- to 10-wk-old Nu/Nu mice were anesthetized with 100 mg/kg ketamine and 20 mg/kg xylazine administered i.p. before a transverse abdominal incision. The stomach, spleen, and pancreas were exposed and 10^6 tumor cells in 50 μ L PBS were injected into the tail of the pancreas. To prevent leakage of the cell suspension from the injection site, a cotton swab was held in contact with the injection site for 30 s postinjection, and the area was then monitored for evidence of leakage for an additional 30 s (25). A successful injection was determined by the presence of a fluid-filled bleb in the pancreas without i.p. leakage. The incision was then sutured closed. We have found this method to form reproducible orthotopic pancreatic tumors.⁴

Positron emission tomography. Relative volumes of orthotopic pancreatic tumors were estimated by measuring uptake of ^{18}F -2-deoxyglucose (FDG) using positron emission tomography (PET). ^{18}F was produced through bombardment of ^{18}O -enriched H_2O on a PETTrace cyclotron (GE Medical Systems), and FDG was produced on an FX-FN synthesis unit (GE Medical Systems) through nucleophilic substitution of a precursor with ^{18}F . Mice were anesthetized with 2% isoflurane in O_2 , and 200 to 250 μ Ci FDG was administered in ~ 100 μ L via the lateral tail vein. Mice remained anesthetized for 1 h after radiotracer injection to minimize FDG uptake by skeletal muscle and brown fat tissue. Mice were transferred to the bed of a Vista microPET scanner (GE Medical Systems), and microPET data were collected for 10 min over a 4-cm longitudinal field of view. The coincidence events measured were then reconstructed into a three-dimensional image of FDG concentration using an ordered subsets expectation maximization algorithm (26).

Immunofluorescence. Tumors were frozen in optimum cutting temperature and 8- μ m sections were cut and stained with the appropriate antibodies: polyclonal goat anti-CCN2 (Santa Cruz Biotechnology, Inc.), monoclonal anti-pimonidazole (NPI, Inc.), and Alexa 488 or 594 secondary antibodies (Invitrogen). Slides were mounted in Vectashield mounting medium containing 4',6-diamidino-2-phenylindole (DAPI; Vector Laboratories, Inc.). Images were captured at $\times 20$ magnification using a Qimaging Retica EXi camera mounted on a Leica DM6000B microscope (JH Technologies).


Hypoxia treatment and apoptosis assay. Cells were placed in a humidified Invivo₂ 400 hypoxia workstation (Ruskin, Inc.) at 5% CO_2 and the indicated oxygen tensions. For the apoptosis assay, cells were incubated in 0.5% O_2 before cell harvest and staining with annexin-V-conjugated FITC antibody according to manufacturer's instructions. Apoptotic cells were quantified using a FACSCalibur flow cytometer (BD Biosciences) with subsequent data analysis using CellQuest software.

CCN2 Northern blots and Western blots. For Northern blot analysis, cells were harvested in Trizol reagent (Invitrogen), and 5 to 10 μ g of glyoxal-treated total RNA was loaded on a sodium phosphate agarose gel using standard methods. Northern blots were probed with ^{32}P -labeled CCN2 and visualized by exposure to a phosphorscreen with subsequent scanning on a Storm 860 scanner (Molecular Dynamics).

To analyze CCN2 levels *in vivo*, small pieces of excised tumors were stored in RNAlater stabilization reagent (Qiagen, Inc.) at $-20^\circ C$ according to manufacturer's instructions. Tissues were removed from the RNAlater reagent and homogenized in 1 mL Trizol for RNA extraction and subsequent Northern blot.

⁴ C.M. Ham and A.J. Giaccia, unpublished data.

Figure 1. CCN2 knockdown decreases soft agar growth of pancreatic tumor cells. **A**, Northern blot of total RNA (*top*) from wild-type Panc-1 cells and Panc-1 cells stably expressing one of six shRNA sequences targeting *CCN2* (A–F). *Scr*, scrambled control shRNA. *Bottom*, Northern blot of Panc-1 cells stably expressing shRNA D and subsequently transfected with *CCN2*. **B**, 10^5 Panc-1 cells stably expressing the indicated shRNA constructs were plated as monolayers and counted every 4 d. Cells (10^5) were replated after each count and the cumulative cell numbers are plotted. *Points*, mean; *bars*, SE. **C**, 5×10^3 Panc-1 cells stably expressing the indicated shRNA constructs were plated in soft agar and allowed to grow for 2 to 3 wk. Colonies were stained with Giemsa stain for visualization and representative images are shown. **D**, quantification of the soft agar colonies from **C**. *Columns*, mean; *bars*, SE; *, $P < 0.05$ relative to control. *Wt*, wild-type.



For analyzing secreted CCN2 protein, equal numbers of cells were plated 24 h before the experiment and the media was changed to DMEM containing 0.25% bovine serum albumin at time 0. Plates were kept under either 21% or 0.5% O₂ and conditioned medium was collected (and cell lysates harvested) at the indicated time points. The conditioned medium was cleared by centrifugation and the media volume was normalized to 10 mL with water. One hundred eighty microliters of heparin sepharose CL-6B beads (Amersham Biosciences) were added to 4.5 mL of conditioned media and rocked for 24 h at 4°C. Beads were washed thrice with cold PBS, and CCN2 protein was collected by incubating the beads with 200 μ L of radioimmunoprecipitation assay buffer at 95°C for 10 min. Twenty microliters of each sample were loaded for Western blot analysis.

Protein lysates were harvested at the indicated time points using a 9M Urea, 0.075M Tris buffer (pH 7.6). Protein lysates were quantified using the Bradford assay, and subjected to reducing SDS-PAGE using standard methods. Western blots were probed with the following antibodies: HIF-1 α (610959; BD Biosciences), actin (AC-40 A3853; Sigma-Aldrich), CCN2 (SC-14939; Santa Cruz Biotechnology, Inc.).

Statistical analysis. Soft agar colony numbers or apoptotic cell numbers were assessed by Student's *t* test, and survival of orthotopic tumor-bearing mice were assessed by log-rank test. Areas under the curves (AUC) were calculated for each tumor growth curve and assessed by ANOVA or Student's *t* test as appropriate. χ^2 analyses were performed on the clinical data. GraphPad Prism was used for statistical analyses.

Results

To investigate the role of tumor cell-derived CCN2 in pancreatic cancer, we generated a number of stable *CCN2* shRNA-expressing Panc-1 cell lines. Six *CCN2* shRNA sequences were tested, with three shRNAs producing varying degrees of CCN2 knockdown (Fig. 1A, *top*). As controls against potential off-target effects of the *CCN2* shRNA, we used a scrambled shRNA sequence and also added CCN2 back into cells stably expressing shRNA D, which had produced the most potent CCN2 knockdown (Fig. 1A, *bottom*).




Figure 2. Partial knockdown of *CCN2* induces modest delays in s.c. pancreatic tumor growth. **A**, 10^7 wild-type or *CCN2* shRNA-expressing Panc-1 cells were implanted s.c. in nu/nu mice and tumor volumes were monitored over time. Points, mean with at least seven mice per group; bars, SE; no significant difference by Student's *t* test of AUCs. **B**, Northern blots of wild-type or *CCN2* shRNA-expressing Panc-1 cells used for tumor implants compared with cell lines generated from representative tumors excised 15 to 18 wk after tumor implant. **C**, 10^7 wild-type or *CCN2* shRNA-expressing Su86.86 cells were implanted s.c. in nu/nu mice and tumor volumes were monitored over time. Points, mean with at least five mice per group; bars, SE; *, $P < 0.05$ by Student's *t* test of AUCs. *Inset*, Northern blot of total RNA from wild-type and *CCN2* shRNA-expressing Su86.86 cells. **D**, Northern blot (*top*) of cell lines generated from representative subcutaneous *CCN2* shRNA-expressing Su86.86 tumors harvested 8 wk after tumor implant. Input cells represent the original *CCN2* shRNA-expressing Su86.86 cells used for the tumor implants. *Bottom*, Northern blot of total RNA extracted directly from excised s.c. wild-type or *CCN2* shRNA-expressing Su86.86 tumors.

CCN2 levels did not affect the proliferation of Panc-1 tumor cells (Fig. 1*B*) or Su86.86 tumor cells (Supplementary Fig. S1) in monolayer culture in air (or at 2% oxygen; data not shown). However, *CCN2* knockdown induced significant decreases in the ability of Panc-1 cells to grow in soft agar (Fig. 1*C*) with the greatest decrease in soft agar growth induced by shRNA "D" (Fig. 1*D*). Soft agar growth was not decreased in cells expressing either a scrambled control shRNA sequence or shRNA constructs that did not effectively knockdown *CCN2*. Furthermore, adding *CCN2* expression back to cells expressing *CCN2* shRNA D significantly increased the soft agar growth of these cells. These data indicate that the level of *CCN2* knockdown influences the ability of Panc-1 cells to grow in soft agar.

We then s.c. implanted Nude mice with wild-type Panc-1 cells or a pooled population of Panc-1 cells expressing the most potent *CCN2* shRNA construct D. We observed a modest growth delay in tumors derived from the pooled population of *CCN2* shRNA-expressing cells (Fig. 2*A*), but the difference was not statistically significant. Interestingly, the take-rate of shRNA-expressing tumors was only 80%, and cell lines generated from these tumors at the end of the experiment indicated a marked increase in *CCN2* expression relative to the input shRNA-expressing cells (Fig. 2*B*). These data indicate a heterogeneous loss of stable shRNA expression *in vivo* and/or an outgrowth of tumor cells that had incomplete knockdown of *CCN2*. Similarly, cell lines generated from wild-type Panc-1 tumors excised at the end of the experiment had increased *CCN2* expression relative to the initial Panc-1 input cells. These data suggest that cells

expressing high levels of *CCN2* have a growth advantage in solid tumors.

In agreement with these data, pooled populations of Su86.86 cells expressing *CCN2* shRNA-D exhibited a modest statistically significant tumor growth delay (Fig. 2*C*) relative to wild-type Su86.86 tumors and had a 75% take-rate. An increase in *CCN2* expression was again observed when cell lines generated from *CCN2* shRNA-expressing tumors at the end of the experiment were compared with the input shRNA-expressing cells (Fig. 2*D, top*). Furthermore, when excised tumor tissue was directly assayed for *CCN2*, we observed *CCN2* expression levels in shRNA-expressing tumors that were comparable with wild-type Su86.86 tumors (Fig. 2*D, bottom*). Taken with Fig. 2*A* to *B*, these data indicate that the microenvironment in pancreatic tumors promotes the outgrowth of tumor cells that express high levels of *CCN2*.

To determine if clonal populations of cells with more effective knockdown of *CCN2* would result in greater suppression of tumor growth, we screened 40 clones from the heterogeneous Panc-1 *CCN2* shRNA D-expressing population. We observed a wide range of *CCN2* knockdown efficiency (Supplementary Fig. S2) despite puromycin selection of the cells; all clones showed at least some *CCN2* knockdown with several clones showing >90% reductions in *CCN2* levels. We were unable to derive clonal populations of Su86.86 shRNA-expressing cells because Su86.86 cells do not grow from single cells in our hands.

We also tested the efficacy of *CCN2* knockdown in selected clones after exogenous stimulation of *CCN2* overexpression. *CCN2* is

considered an immediate-early response gene, and we have found that serum stimulation of Panc-1 cells induces a transient increase in *CCN2* expression as reported in other cell types (27, 28). Wild-type Panc-1 cells exposed to a change of fresh DMEM containing 10% FBS showed remarkable increases in *CCN2* RNA and protein expression from 30 minutes up to 8 hours, at which time *CCN2* expression returned to control levels (Fig. 3A). Changing the medium to DMEM without serum did not induce changes in *CCN2* RNA or protein expression (data not shown). We chose two clones (#1 and #5) for further study based on sustained knockdown of *CCN2* by >90% despite stimulation by 10% serum (Fig. 3B).


CCN2 shRNA-expressing Panc-1 clones displayed dramatically decreased growth in soft agar, whereas Panc-1 cells that overexpressed *CCN2* showed significantly increased soft agar growth relative to wild-type Panc-1 cells (Fig. 3C–D). We also found that neither complete knockdown of *CCN2* in clonal cell populations nor overexpression of *CCN2* affected monolayer growth of Panc-1 tumor cells (Supplementary Fig. S3). These data suggest that *CCN2*

increases the growth and/or survival of pancreatic tumor cells under nonideal growth conditions as are found in soft agar matrices, and also illustrate the importance of tumor cell-derived *CCN2* in an *in vitro* system devoid of stromal cells.

We then tested our hypothesis that *CCN2* shRNA-expressing clones with more efficient *CCN2* knockdown than the pooled *CCN2* shRNA-expressing cells would provide greater effects on tumor growth. Indeed, the *CCN2* shRNA-expressing Panc-1 clones with the greatest knockdown of *CCN2* did not form tumors when implanted s.c. (Fig. 4A). Furthermore, subcutaneous tumors derived from *CCN2* overexpressing Panc-1 cells grew significantly faster than wild-type Panc-1 tumors. These data indicate an essential role for tumor cell-derived *CCN2* relative to stromal cell-derived *CCN2* in the growth of s.c. pancreatic tumor xenografts.

We wanted to determine the role of tumor cell-derived *CCN2* in the growth of tumors implanted in tissue that is more representative of clinical pancreatic tumors. Orthotopically implanted pancreatic tumors (29) model the local development and distal

Figure 3. Clonal populations of *CCN2* shRNA-expressing Panc-1 cells show decreased growth in soft agar. **A**, kinetics of *CCN2* RNA and protein induction after changing media on the cells to fresh (10% FBS containing) DMEM. **B**, *CCN2* expression in the indicated cell types was stimulated by changing the medium. Clonal populations with high levels of *CCN2* knockdown after serum stimulation were selected for further study. **C**, 5×10^3 wild-type, *CCN2* overexpressing, and *CCN2* shRNA-expressing Panc-1 clones were plated in soft agar and allowed to grow for 2 wk. Colonies were stained with Geimsa stain for visualization, and photographs were taken using a dissecting microscope ($\times 2$). **D**, quantification of soft agar colonies from **C**. Columns, mean; bars, SE; *, $P < 0.05$ relative to control.






Figure 4. CCN2 knockdown decreases subcutaneous and orthotopic tumor growth, and increases the survival of orthotopic tumor-bearing mice. *A*, 10⁷ wild-type, CCN2 overexpressing, and CCN2 shRNA-expressing Panc-1 clones were s.c. implanted in nu/nu mice. Tumor volumes were monitored weekly using calipers. Panc-1 + CCN2 shRNA (pool) curve is included from Fig. 2*A* for visual comparison purposes only. Points, mean with at least five mice per group; bars, SE; *, *P* < 0.05 by ANOVA of AUCs. *B*, 10⁷ wild-type, CCN2 overexpressing, or CCN2 shRNA-expressing Panc-1 cells were orthotopically implanted in nu/nu mice. Tumor volume was monitored by uptake of i.v. administered FDG using PET. Representative maximum intensity projections are shown, with regions of high FDG uptake highlighted. G, Harderian glands; H, heart; K, kidney; B, bladder; T, tumor. *C*, survival plot of mice bearing orthotopic pancreatic tumors. Data from three to nine mice per group; *P* values indicate comparison with mice bearing wild-type tumors by log-rank test.

metastasis of clinical pancreatic adenocarcinomas, and are responsive to stimulation by TGF- β (30), which is an upstream inducer of CCN2. Furthermore, transcription factors such as hypoxia-inducible factor-1 (HIF-1) can have opposite effects on the growth of tumors implanted in relatively poorly vascularized s.c. sites compared with the often better-perfused tissues of origin (31). Thus, the tissue of origin represents an optimal site for tumor xenograft implantation, and we therefore followed up our s.c. studies by testing the effect of CCN2 on the growth of pancreatic tumor xenografts implanted orthotopically.

Mice bearing orthotopically implanted tumors were imaged by PET for FDG uptake (Fig. 4*B*) and monitored for overall survival (Fig. 4*C*). CCN2 overexpression enhanced the growth and metastasis of orthotopic pancreatic tumors as detected by FDG, leading to significantly decreased survival times when compared with wild-type Panc-1 tumors. Mice implanted with CCN2 shRNA-expressing Panc-1 clones exhibited smaller orthotopic tumors by

PET imaging with a concomitant statistically significant increase in survival (Fig. 4*C*). Taken together, these data indicate that tumor cell-derived CCN2 is important for the growth of both s.c. and orthotopic pancreatic tumor xenografts.

CCN2 expression can be induced in some cell types by exposure to low levels of oxygen (32–35), and pancreatic tumors contain relatively large numbers of hypoxic cells. We found that CCN2 colocalized with the hypoxia marker pimonidazole (Hypoxyprobe-1) in orthotopic Panc-1 tumors (Fig. 5*A*), suggesting that CCN2 may be regulated by hypoxia in pancreatic tumors. However, CCN2 is a secreted protein that is capable of binding to the extracellular matrix (36) and the presence of CCN2 protein in hypoxic tumor regions is not necessarily indicative of CCN2 production by the hypoxic cells.


We therefore studied the hypoxia-inducibility of CCN2 in Panc-1 cells *in vitro*. CCN2 protein levels in cell lysates and secreted into the surrounding medium were elevated after exposure of cells to

0.5% O₂ (Fig. 5B). When cell lysates were assayed for HIF-1 α , we found that the increase in CCN2 production and secretion in response to hypoxia occurred when HIF-1 α levels were maximally induced (at 4 hours). These data are consistent with previously published data suggesting a role for HIF-1 in CCN2 regulation in some cell types (32, 33). The levels of HIF-1 and intracellular CCN2 decreased by 16 hours of 0.5% O₂ in relation to death of significant numbers of tumor cells. Increases in secreted CCN2 have been observed without significant changes in intracellular CCN2 levels (37), indicative of rapid and quantitative secretion of CCN2 upon up-regulation (38). Thus, the secreted CCN2 in Fig. 5B is a cumulative indication of CCN2 levels secreted from the cells over time, whereas the intracellular CCN2 levels are more representative of CCN2 production at the time of cell harvest. CCN2 expression and

secretion is therefore induced by hypoxia in Panc-1 tumor cells *in vitro*, and CCN2 protein colocalizes with hypoxic cells in Panc-1 tumors.

Our data indicate that CCN2 derived from tumor cells is required for efficient pancreatic tumor growth, and *in vivo* growth provides a selective pressure for the outgrowth of tumor cells that express high levels of CCN2. CCN2 also enhances the growth and survival of pancreatic tumor cells in soft agar but had no effect on cells grown as monolayers. These observations led us to postulate that CCN2 may provide a growth or survival advantage to tumor cells under nonideal growth conditions. Solid tumor hypoxia can represent a significant microenvironmental stress for cells, and we hypothesized that CCN2 affected tumor growth partly by modulating the survival of tumor cells found in hypoxic regions of

Figure 5. CCN2 expression and secretion is increased by hypoxia and protects pancreatic tumor cells from hypoxia-induced apoptosis. **A**, pimonidazole (Hypoxyprobe-1) was administered 90 min before orthotopic Panc-1 tumor excision, and frozen tumor sections were analyzed for pimonidazole (red) and CCN2 (green). Areas of colocalization are indicated (yellow), and nuclei are stained with DAPI (blue). **B**, Western blots of CCN2 and HIF-1 α in Panc-1 cells incubated at 21% or 0.5% oxygen for the indicated periods of time before collection of cell lysate and conditioned medium. Secreted CCN2 was obtained by contacting conditioned medium with heparin sepharose-coated beads before loading. Actin in the cell lysate was used as a loading control. **C**, flow cytometric quantification of wild-type Panc-1 cells or Panc-1 shRNA-expressing clones exposed to 0.5% O₂ for 24 h. Apoptotic cells were measured by Annexin-V staining and quantified by flow cytometry. *, $P < 0.05$ relative to control cells.






Figure 6. CCN2 is associated with tumor tissue and hypoxia in clinical pancreatic adenocarcinoma samples. *A*, immunohistochemical staining for CCN2 and CAIX in paraffin-embedded samples from clinical pancreatic cancer patients. *B*, CCN2 staining in pancreatic adenocarcinoma versus normal pancreatic tissue from the same patient ($n = 20$). *C*, CCN2 staining in pancreatic adenocarcinoma tissue versus stroma surrounding the tumor in the same section ($n = 30$). *D*, overall CCN2 staining and CAIX staining in step-sections of clinical pancreatic adenocarcinoma samples ($n = 30$).

tumors. We therefore treated Panc-1 cells with 0.5% O₂ for 24 h and monitored cell viability and apoptosis by annexin-V staining. Interestingly, we found that wild-type Panc-1 cells did not survive well in 0.5% O₂ *in vitro*, and that CCN2 inhibition produced statistically significant increases in apoptotic cell death in response to hypoxia (Fig. 5C). Importantly, CCN2 inhibition did not affect the survival of Panc-1 cells in normoxia (data not shown). These data indicate that CCN2 protects Panc-1 cells against hypoxia-induced apoptosis and, when taken with our observations that CCN2 increases the growth of tumor cells in soft agar, provide an explanation for the observed *in vivo* selection for the enhanced survival and outgrowth of tumor cells that express high levels of CCN2.

To put our data in context with clinical pancreatic cancer, we obtained pancreatic adenocarcinoma samples and stained them for CCN2. Pancreatic adenocarcinoma and normal pancreas tissue samples were collected from the same patient where possible (20 of 30 patients). Both the intensity and the degree of CCN2 staining (Fig. 6A) were scored, and an overall rating of CCN2 staining was determined. We found that 14 of 20 patients (70%) had higher CCN2 staining in pancreatic adenocarcinoma

tissue compared with normal pancreatic tissue (Fig. 6B), and that 19 of 30 patients (63%) had more intense CCN2 staining in tumor tissue compared with the stromal tissue surrounding the tumor (Fig. 6C). We also compared CCN2 staining with levels of the classic endogenous hypoxia marker CAIX in step-sections. We found that 20 of 30 patients (67%) showed similar overall levels of staining for both CAIX and CCN2 (Fig. 6D), and there was a considerable degree of overlap between CCN2 and CAIX in individual tumor samples (Fig. 6A). Taken together, these data indicate that CCN2 expression is elevated in primary pancreatic adenocarcinomas and that CCN2 is found preferentially associated with hypoxic pancreatic tumor cells rather than the surrounding stroma.

Discussion

We have identified a critical role for tumor cell-derived CCN2 in the growth of pancreatic tumors. A robust selective pressure is present in pancreatic tumors for cells that express high levels of CCN2 and this selection is driven, at least in part, by tumor hypoxia. CCN2 protects tumor cells against microenvironmental

stresses such as hypoxia by inhibiting apoptosis and enhancing tumor growth. Greater than 90% knockdown of CCN2 by shRNA in clonal populations of cells was required to prevent CCN2-expressing cells from populating the tumors. It is important to note that wild-type Panc-1 cells and CCN2 shRNA-expressing clones exhibited differences in tumor growth despite being implanted in the same tissue type, suggesting that any CCN2 derived from the surrounding stromal cells was not sufficient to affect the differential growth rates of the tumors. Similarly, immunologic inhibition of CCN2 is the most robust in suppressing the growth of tumors that contain CCN2-expressing tumor cells (22, 23), indicating the role of stromal cell-derived CCN2 in pancreatic tumor growth is minimal.

CCN2 is downstream of the ras/MAP-ERK kinase/extracellular signal-regulated kinase pathway (39), and activation of the ras pathway mediates basal overexpression of CCN2 in pancreatic cancer cells in a Smad4-independent manner (40). Thus, activating ras mutations, which are known to occur in the vast majority of pancreatic adenocarcinomas, would induce CCN2 overexpression in the early stages of pancreatic tumor development. When combined with our data illustrating a role for CCN2 in enhancing the growth of pancreatic tumor cells in soft agar (Figs. 1C-D and 3C-D), the combination of an activating ras mutation with the resultant overexpression of CCN2 would lead to enhanced proliferation of pancreatic tumor cells. Interestingly, secreted CCN2 is known to directly interact with TGF- β and enhance binding to the TGF- β receptor (7); CCN2 expression is also induced by TGF- β . These observations suggest a feedforward loop mediated by TGF- β and CCN2 resulting in further increases in CCN2 production and tumor growth. There are therefore multiple mechanisms that can increase CCN2 expression levels in pancreatic tumor cells throughout the development and growth of a solid tumor.

As tumors grow, increasing numbers of tumor cells become hypoxic and induction of CCN2 expression by hypoxia has been observed in a number of normal and neoplastic cell types. Mechanistically, hypoxic induction of CCN2 is mediated by HIF-1 transcriptional activity in murine primary renal tubular epithelial cells (32), and in dermal fibroblasts from systemic sclerosis patients (33). Our data are consistent with these reports in that we observed increased CCN2 expression and secretion with HIF-1 α stabilization in response to hypoxia (Fig. 5B). Interestingly, HIF-2-mediated CCN2 expression has been observed in human 786-0 renal cell carcinoma cells that lack HIF-1 (34), indicating that both HIF-1 and HIF-2 can influence induction of CCN2 by hypoxia in different cell types. Hypoxia has also been shown to increase CCN2 expression in human chondrosarcoma cells by 3'-untranslated region-mediated increases in CCN2 mRNA stability during hypoxia (35). We observed modest increases in intracellular CCN2 protein levels with greater increases in secreted CCN2 protein from hypoxic tumor cells, consistent with reports of hypoxia-induced increases in CCN2 secretion from human fibroblasts with lesser increases observed in intracellular CCN2 levels (37). CCN2 is efficiently secreted through the Golgi from rat hepatic stellate cells and TGF- β -stimulated human dermal fibroblasts (38), and we have observed localization of CCN2 to the Golgi in hypoxic pancreatic tumor cells (data not shown). Taken together, these data indicate that hypoxia induces increased expression and rapid secretion of CCN2 from pancreatic tumor cells, providing a further mechanism for increased CCN2 production with tumor growth.

Hypoxia also represents a microenvironmental stressor for tumor cells, and we observed increased apoptosis of pancreatic tumor cells in response to hypoxia with genetic inhibition of CCN2 (Fig. 5C). These data are consistent with reports of CCN2 inhibition increasing apoptosis of human rhabdomyosarcoma cells (41), and with observations that the lungs of perinatally lethal CCN2 knockout mice contain increased levels of apoptosis compared with wild-type mice (42). CCN2 also enhances TGF- β -mediated apoptosis in human breast cancer cells (43), human aortic smooth muscle cells (44), and rat peritoneal endothelial cells (45), whereas CCN2 has been reported to protect murine endothelial cells against growth factor deprivation-induced apoptosis (10), and impair apoptosis of chondrocytes (46). Whether these conflicting reports are due to cell type-specific differences in CCN2 action and/or differences in CCN2 structure in these systems is an open question, and is the subject of active investigation.

Healthy human pancreatic tissue does not typically express high levels of CCN2. Significant overexpression of CCN2 mRNA has been observed in acute necrotizing pancreatitis (47) and in pancreatic cancer tissue relative to normal pancreas (48, 49), although one group observed only a modest (nonstatistically significant) increase in CCN2 expression in pancreatic cancer tissues (50). CCN2 mRNA overexpression has been shown in fibroblasts, scattered acinar cells, and pancreatic tumor cells, whereas CCN2 protein is associated with tumor cells and fibroblasts surrounding pancreatic tumor tissue (49). We have found that scattered acinar cells and subsets of islet cells in normal pancreatic tissue can stain intensely for CCN2 (data not shown), but that overall CCN2 staining is typically low in the healthy pancreas. We observed more intense and more abundant CCN2 staining in the majority of pancreatic tumor samples when compared with normal pancreatic tissue from the same patient (Fig. 6B). CCN2 also colocalizes with CAIX staining in human pancreatic tumor tissue (Fig. 6A and D) and in fibroblasts adjacent to necrosis. Taken together, these data highlight the relationship between hypoxia and CCN2 expression in clinical pancreatic cancer, and support our preclinical findings that CCN2 derived from tumor cells is important for pancreatic tumor growth and progression.

Taken together, our data indicate that CCN2 derived from tumor cells is up-regulated by hypoxia, enhances the survival of hypoxic tumor cells, and increases tumor growth. CCN2 therefore acts as a protective factor for pancreatic tumor cells, enhancing the growth and progression of pancreatic tumors despite relatively high fractions of pancreatic tumor cells residing in hypoxic tumor regions. The selective pressure exerted by the tumor microenvironment produces tumor cells that express high levels of CCN2, which ultimately leads to enhanced tumor growth. Our data indicate a critical role for CCN2 derived from pancreatic tumor cells in the growth of pancreatic tumors and identify CCN2 as a therapeutic target for the clinical treatment of pancreatic cancer.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

Received 3/14/2008; revised 10/8/2008; accepted 10/17/2008.

Grant support: NIH (A.J. Giaccia), the Blue Dot Fund (A. Koong), and a postdoctoral fellowship from the Canadian Institutes of Health Research (K.L. Bennewith).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

We thank Drs. Scott Welford, Tom Johnson, and Sandra Turcotte for expert technical assistance and helpful discussion, and Drs. Barbara Bedogni, Nadja Dornhofer, and Derek Bingham for helpful discussion.

References

- Jemal A, Siegel R, Ward E, et al. Cancer statistics, 2008. *CA Cancer J Clin* 2008;58:71-96.
- Korc M. Pancreatic cancer-associated stroma production. *Am J Surg* 2007;194:S84-6.
- Brigstock DR. The connective tissue growth factor/cysteine-rich 61/nephroblastoma overexpressed (CCN) family. *Endocr Rev* 1999;20:189-206.
- Leask A, Abraham DJ. All in the CCN family: essential extracellular signaling modulators emerge from the bunker. *J Cell Sci* 2006;119:4803-10.
- Perbal B. CCN proteins: multifunctional signalling regulators. *Lancet* 2004;363:62-4.
- Brigstock DR, Steffen CL, Kim GY, Vegunta RK, Diehl JR, Harding PA. Purification and characterization of novel heparin-binding growth factors in uterine secretory fluids. Identification as heparin-regulated Mr 10,000 forms of connective tissue growth factor. *J Biol Chem* 1997;272:20275-82.
- Abreu JG, Ketpura NI, Reversade B, De Robertis EM. Connective-tissue growth factor (CTGF) modulates cell signalling by BMP and TGF- β . *Nat Cell Biol* 2002;4:599-604.
- Heng EC, Huang Y, Black SA, Jr. Trackman PC. CCN2, connective tissue growth factor, stimulates collagen deposition by gingival fibroblasts via module 3 and α 6- and β 1 integrins. *J Cell Biochem* 2006;98:409-20.
- Gao R, Brigstock DR. A novel integrin α 5 β 1 binding domain in module 4 of connective tissue growth factor (CCN2/CTGF) promotes adhesion and migration of activated pancreatic stellate cells. *Gut* 2006;55:856-62.
- Babic AM, Chen CC, Lau LF. Fisp12/mouse connective tissue growth factor mediates endothelial cell adhesion and migration through integrin α v β 3, promotes endothelial cell survival, and induces angiogenesis *in vivo*. *Mol Cell Biol* 1999;19:2958-66.
- Leask A, Denton CP, Abraham DJ. Insights into the molecular mechanism of chronic fibrosis: the role of connective tissue growth factor in scleroderma. *J Invest Dermatol* 2004;122:1-6.
- Koliopoulos A, Friess H, di Mola FF, et al. Connective tissue growth factor gene expression alters tumor progression in esophageal cancer. *World J Surg* 2002;26:420-7.
- Xie D, Yin D, Wang HJ, et al. Levels of expression of CYR61 and CTGF are prognostic for tumor progression and survival of individuals with gliomas. *Clin Cancer Res* 2004;10:2072-81.
- Xie D, Nakachi K, Wang H, Elashoff R, Koeffler HP. Elevated levels of connective tissue growth factor, WISP-1, and CYR61 in primary breast cancers associated with more advanced features. *Cancer Res* 2001;61:8917-23.
- Liu L, Li Z, Feng G, You W, Li J. Expression of connective tissue growth factor is in agreement with the expression of VEGF, VEGF-C, -D and associated with shorter survival in gastric cancer. *Pathol Int* 2007;57:712-8.
- Sala-Torra O, Gundacker HM, Stirewalt DL, et al. Connective tissue growth factor (CTGF) expression and outcome in adult patients with acute lymphoblastic leukemia. *Blood* 2007;109:3080-3.
- Wong YF, Cheung TH, Tsao GS, et al. Genome-wide gene expression profiling of cervical cancer in Hong Kong women by oligonucleotide microarray. *Int J Cancer* 2006;118:2461-9.
- Deng Y-Z, Chen P-P, Wang Y, et al. Connective tissue growth factor is overexpressed in esophageal squamous cell carcinoma and promotes tumorigenicity through -catenin-t-cell factor/lef signaling. *J Biol Chem* 2007;282:36571-81.
- Zirn B, Hartmann O, Samans B, et al. Expression profiling of Wilms tumors reveals new candidate genes for different clinical parameters. *Int J Cancer* 2006;118:1954-62.
- Shakunaga T, Ozaki T, Ohara N, et al. Expression of connective tissue growth factor in cartilaginous tumors. *Cancer* 2000;89:1466-73.
- Chen PP, Li WJ, Wang Y, et al. Expression of Cyr61, CTGF, and WISP-1 correlates with clinical features of lung cancer. *PLoS ONE* 2007;2:e534.
- Dornhofer N, Spong S, Bennewith K, et al. Connective tissue growth factor-specific monoclonal antibody therapy inhibits pancreatic tumor growth and metastasis. *Cancer Res* 2006;66:5816-27.
- Aikawa T, Gunn J, Spong SM, Klaus SJ, Korc M. Connective tissue growth factor-specific antibody attenuates tumor growth, metastasis, and angiogenesis in an orthotopic mouse model of pancreatic cancer. *Mol Cancer Ther* 2006;5:1108-16.
- Le QT, Kong C, Lavori PW, et al. Expression and prognostic significance of a panel of tissue hypoxia markers in head-and-neck squamous cell carcinomas. *Int J Radiat Oncol Biol Phys* 2007;69:167-75.
- Bruns CJ, Harbison MT, Kuniyasu H, Eue I, Fidler IJ. *In vivo* selection and characterization of metastatic variants from human pancreatic adenocarcinoma by using orthotopic implantation in nude mice. *Neoplasia* 1999;1:50-62.
- Graves EE, Quon A, Loo BW, Jr. RT_Image: an open-source tool for investigating PET in radiation oncology. *Technol Cancer Res Treat* 2007;6:111-21.
- Perbal B. NOV (nephroblastoma overexpressed) and the CCN family of genes: structural and functional issues. *Mol Pathol* 2001;54:57-79.
- Ryseck RP, Macdonald-Bravo H, Mattei MG, Bravo R. Structure, mapping, and expression of fisp-12, a growth factor-inducible gene encoding a secreted cysteine-rich protein. *Cell Growth Differ* 1991;2:225-33.
- Vezeridis MP, Doremus CM, Tibbetts LM, Tzanakakis G, Jackson BT. Invasion and metastasis following orthotopic transplantation of human pancreatic cancer in the nude mouse. *J Surg Oncol* 1989;40:261-5.
- Rowland-Goldsmith MA, Maruyama H, Matsuda K, et al. Soluble type II transforming growth factor- β receptor attenuates expression of metastasis-associated genes and suppresses pancreatic cancer cell metastasis. *Mol Cancer Ther* 2002;1:161-7.
- Blouw B, Song H, Tihan T, et al. The hypoxic response of tumors is dependent on their microenvironment. *Cancer Cell* 2003;4:133-46.
- Higgins DF, Biju MP, Akai Y, Wutz A, Johnson RS, Haase VH. Hypoxic induction of Ctgf is directly mediated by Hif-1. *Am J Physiol Renal Physiol* 2004;287:F1223-32.
- Hong KH, Yoo SA, Kang SS, Choi JJ, Kim WU, Cho CS. Hypoxia induces expression of connective tissue growth factor in scleroderma skin fibroblasts. *Clin Exp Immunol* 2006;146:362-70.
- Chintalapudi MR, Markiewicz M, Kose N, et al. Cyr61/CCN1 and CTGF/CCN2 mediate the pro-angiogenic activity of VHL mutant renal carcinoma cells. *Carcinogenesis* 2008;29:696-703.
- Kondo S, Kubota S, Mukudai Y, et al. Hypoxic regulation of stability of connective tissue growth factor/CCN2 mRNA by 3'-untranslated region interacting with a cellular protein in human chondrosarcoma cells. *Oncogene* 2006;25:1099-110.
- Kireeva ML, Latinkic BV, Kolesnikova TV, et al. Cyr61 and Fisp12 are both ECM-associated signaling molecules: activities, metabolism, and localization during development. *Exp Cell Res* 1997;233:63-77.
- Rimon E, Chen B, Shanks AL, Nelson DM, Sadovsky Y. Hypoxia in human trophoblasts stimulates the expression and secretion of connective tissue growth factor. *Endocrinology* 2008;149:2952-8.
- Chen Y, Segarini P, Raoufi F, Bradham D, Leask A. Connective tissue growth factor is secreted through the Golgi and is degraded in the endosome. *Exp Cell Res* 2001;271:109-17.
- Pickles M, Leask A. Analysis of CCN2 promoter activity in PANC-1 cells: regulation by ras/MEK/ERK. *J Cell Commun Signal* 2007;1:85-90.
- Kwon S, Munroe X, Crawley SC, et al. Expression of connective tissue growth factor in pancreatic cancer cell lines. *Int J Oncol* 2007;31:693-703.
- Croci S, Landuzzi L, Astolfi A, et al. Inhibition of connective tissue growth factor (CTGF/CCN2) expression decreases the survival and myogenic differentiation of human rhabdomyosarcoma cells. *Cancer Res* 2004;64:1730-6.
- Baguma-Nibasheka M, Kablar B. Pulmonary hypoplasia in the connective tissue growth factor (Ctgf) null mouse. *Dev Dyn* 2008;237:485-93.
- Hishikawa K, Oemar BS, Tanner FC, Nakaki T, Luscher TF, Fujii T. Connective tissue growth factor induces apoptosis in human breast cancer cell line MCF-7. *J Biol Chem* 1999;274:37461-6.
- Hishikawa K, Nakaki T, Fujii T. Connective tissue growth factor induces apoptosis via caspase 3 in cultured human aortic smooth muscle cells. *Eur J Pharmacol* 2000;392:19-22.
- Szeto CC, Chow KM, Lai KB, Szeto CY, Kwan BC, Li PK. Connective tissue growth factor is responsible for transforming growth factor- β -induced peritoneal mesothelial cell apoptosis. *Nephron Exp Nephrol* 2006;103:e166-74.
- Fujisawa T, Hattori T, Ono M, et al. CCN family 2/connective tissue growth factor (CCN2/CTGF) stimulates proliferation and differentiation of auricular chondrocytes. *Osteoarthritis Cartilage* 2008;16:787-95.
- di Mola FF, Friess H, Riesle E, et al. Connective tissue growth factor is involved in pancreatic repair and tissue remodeling in human and rat acute necrotizing pancreatitis. *Ann Surg* 2002;235:60-7.
- Wenger C, Ellenrieder V, Alber B, et al. Expression and differential regulation of connective tissue growth factor in pancreatic cancer cells. *Oncogene* 1999;18:1073-80.
- Hartel M, di Mola FF, Gardini A, et al. Desmoplastic reaction influences pancreatic cancer growth behavior. *World J Surg* 2004;28:818-25.
- Aoyagi Y, Oda T, Kinoshita T, et al. Overexpression of TGF- β by infiltrated granulocytes correlates with the expression of collagen mRNA in pancreatic cancer. *Br J Cancer* 2004;91:1316-26.