

Hugh P. Kelly FREE

Judah M. Eisenberg; Akiva Ron



Physics Today **46** (5), 80–81 (1993);

<https://doi.org/10.1063/1.2808917>



View
Online



Export
Citation

CrossMark


E. Bright Wilson

E. Bright Wilson, who died on 12 July 1992, had a highly productive scientific career that spanned 60 years, most of them at Harvard University. He was the model of a university professor, excelling in the two essentially interwoven university functions, research and education. When called upon, he effectively and unstintingly fulfilled his societal responsibilities.

Wilson was born in Tennessee in 1908. He earned his BS in 1930 and his MA at Princeton University and his PhD at Caltech in 1933, each in the field of chemistry. After earning his doctorate, Wilson became a junior fellow in the Society of Fellows at Harvard. He went on to become a professor in the chemistry department.

A founder of chemical physics, Wilson in particular led the development of molecular dynamics and spectroscopy, bringing to the field both a thorough theoretical foundation and a practical bent toward the implementation of spectroscopic methods for the effective treatment of general polyatomic molecules. Wilson was a pioneer who placed nuclear dynamics and spectroscopy of polyatomic molecules on a rigorous foundation of both quantum mechanical ideas and general principles that do not depend on particular mechanical models. To achieve this model-independent basis, he rigorously exploited the dynamical constraints required by the symmetry of molecules. Wilson characterized the electronic, vibrational, rotational and nuclear spin components of the complete molecular wavefunction in a penetrating group theoretical approach. This work, like much of Wilson's, laid a permanent founda-

E. Bright Wilson



tion that has endured to the present.

The fundamental theory of polyatomic spectroscopy was effectively laid out in Wilson's basic works of the mid-1930s. His deep physical insight, coupled to his conviction that molecular spectroscopy must be useful to a broad scientific audience, led him to a relatively complete development of the rotation-vibration theory of molecules for small-amplitude oscillations. His breadth of understanding of molecular structure, in particular the nature of chemical bonds and valence forces, enabled him to formulate molecular dynamics in terms of the internal molecular coordinates, thus allowing him to make use of the regularity of molecular structural units. His classification of molecular vibrations and his reduction of the Hamiltonian using the irreducible representations of the molecular point group then allowed quantitative analysis of complex systems to be made in the precomputer age. Wilson's methods became the standard treatment for the vibrations of polyatomic molecules. In addition, Wilson and his research group at Harvard developed a number of experimental methods for vibrational spectroscopies during the 1930s.

Wilson's work during World War II was on the assessment of underwater naval explosives. He also did much to develop the field of operations research while serving in the Department of Defense in 1952-53. In these activities Wilson demanded valid empirical data and unambiguous analysis, paralleling his rigorous approach to chemical physics.

In the late 1940s and 1950s Wilson worked on microwave spectroscopy of polyatomic molecules, the theory of which was based in large part on his fundamental work of the 1930s. Efficient experimental techniques were required for the application of microwave spectroscopy to the broad variety of polyatomic molecules of chemical interest. The Stark-modulated spectrometer, which Wilson invented with Richard Hughes, became the workhorse of the trade until the recent development of time-domain methods.

Wilson exploited the exceptionally high resolution of microwave spectroscopy for the study not only of overall rotational energy levels but also of large-amplitude tunneling motions that produced spectral line splittings. A frequently occurring example of nonrigidity in molecular systems is the internal rotation about carbon-carbon single bonds. Wilson developed the theory required for the analysis of the anharmonic tunneling

motions in such systems, allowing the quantitative characterization of the factors that determine the potential energy for internal rotation. The results of Wilson's work were applied extensively in chemistry and other fields.

Wilson made no separation between teaching and research. His lectures, textbooks and research papers were models of unpretentious logical clarity. His three textbooks, *Introduction to Quantum Mechanics* (with Linus Pauling), *Molecular Vibrations* (with Jack Decius and Paul Cross) and *Introduction to Scientific Research* are classics that remain in print today.

While Wilson gave public service generously—he served both in the Federal government and on numerous committees concerned with societal consequences of technology—his devotion remained steadfastly with the creation of scientific knowledge. In his work, he was scrupulously honest and totally unselfish. Generous in spirit and in ideas, he was a scientist who helped all in the advancement of science. Much of the pleasure of molecular spectroscopy and chemical physics is a consequence of the style of science that he set.

WILLIAM KLEMPERER
Harvard University

Cambridge, Massachusetts

Hugh P. Kelly

Hugh P. Kelly died on 29 June 1992, at the age of 60, after a long and brave struggle with cancer. Characteristically, he taught and carried on research in physics until his last few days. The wide circle of people who met Hugh through his research activities in theoretical atomic physics and through his administrative tasks at the University of Virginia will remember him as one of the rarest of personalities.

Hugh received his AB from Harvard in 1953 and his MSc from UCLA a year later. He then served in the US Marine Corps for three years. After that, he returned to graduate school at the University of California, Berkeley, where he received his doctorate in 1963. Between 1963 and 1965, Hugh did postdoctoral work with Keith Bruckner at the University of California, San Diego, where he mastered methods that Bruckner had initially developed to deal with the nuclear many-particle problem and that now could be used with great effectiveness in atomic theory. Their work allowed them to handle complex electron correlation effects and thus achieve substantially higher preci-



Hugh P. Kelly

sion in atomic calculations, crucial for the understanding of a variety of effects in atomic physics, as well as in chemistry and astrophysics.

It was in these directions that Hugh later moved throughout his scientific career. Indeed, Hugh in large measure helped to bring about profound changes in many-body perturbation theory. The beginning, however, was not without difficulty: Hugh encountered more than a few problems in getting initial acceptance for the new approach, and only his certainty about the correctness of the basic theory and the validity of his careful calculations, as well as his intrinsic good humor, eventually won for it universal recognition.


In 1965 Hugh joined the faculty at the University of Virginia, where he established a program in atomic theory that for many years rested on his own activities, aided only by those of the large number of graduate students who were drawn to work with him.

Hugh's friends could count on him for sensitive and practical support. Over the years he served as chair of the department of physics, dean of the faculty and provost at Virginia, and his fellow feeling was a crucial ingredient of his administrative style, matched only by his extremely high scientific standards.

Hugh had a large family, and only a man with his prodigious capacity for dedicated work could have combined his absolute devotion to family with his equally great devotion to science. With his death, we mourn a fine scholar and a person of great humanity.


JUDAH M. EISENBERG
Tel Aviv University
Tel Aviv, Israel
AKIVA RON

Hebrew University of Jerusalem, Israel ■



CHARGE SENSITIVE PREAMPLIFIER

A250




RUN SILENT — RUN FAST!!!

A NEW STATE-OF-THE-ART
EXTERNAL FET

FET CAN BE COOLED

NOISE: < 100e-RMS (Room Temp.)
< 20e-RMS (Cooled FET)

POWER: 19 mW typical
SLEW RATE: > 475 V/μs
GAIN-BANDWIDTH f_T > 1.5 GHz



If you are using: Solid State Detectors, Proportional counters, Photodiodes, PM tubes, CEMS or MCPs and want the best performance, try an AMPTEK CHARGE SENSITIVE PREAMPLIFIER


Send for Complete Catalog

Features:

- Low noise (less than 100 electrons RMS)
- Low power (5 milliwatts)
- Small size (Hybrids)
- High Reliability
- Radiation hardened (as high as 10⁷ Rads)
- One year warranty

Applications:

- Aerospace
- Portable Instrumentation
- Nuclear Plant Monitoring
- Imaging
- Research Experiments
- Medical and Nuclear Electronics
- Electro-Optical Systems and others.



AMPTEK INC.

6 DE ANGELO DRIVE, BEDFORD, MA 01730 U.S.A. (617) 275-2242

AUSTRALIA: Austeknis PTY Ltd, Kingswood 2763533; AUSTRIA: Item Beratung, Vienna 975958; BELGIUM: Landre Intechmij, Aartselaar 8875382; BRAZIL: Domex Comercio Exterior Ltda, Sao Jose Dos Campos-SP 234235; DENMARK: Ellime, Slangerup 780303; ENGLAND: Teknis Ltd., Crowthorne, Berkshire 780022; FRANCE: Leversan, Rousset, 42290019; GERMANY: Teknis & Co. Sauerlach, 81049543; HONG KONG: Idealand Electronics Ltd, Kowloon, 7443516-9; INDIA: Bakubhai Ambalal Bombay 6323303; ISRAEL: Giveon Agencies Ltd, Tel Aviv, 5612171; ITALY: C.I.E.R. Roma 856814; JAPAN: Jepico, Tokyo 3480623; KOREA: Hongwood International, Seoul, 5551010; NETHERLANDS: Hollinda B.V. The Hague 512801; NORWAY: Ingener Harald Benestad A/S, Lierskogen 850295; PAKISTAN: Fabricon, Karachi 412266; PHILIPPINES: QV Philippines Co. Ltd Metro Manila, 8193365.

Circle number 30 on Reader Service Card

CONTINUOUS FLOW CRYOSTATS FROM

JANIS

Systems offer:

- Compact, light weight cryostats to fit most applications
- Interchangeable, high efficiency, flexible transfer line
- Lowest ultimate temperature
- VP series is top loading for fast sample cooldown and interchange
- Basic SuperTran series usable with either LHe or LN₂

SuperTran-VP

1.4 - 325 K

SuperTran

1.4 - 325 K
(600 K optional)



JANIS RESEARCH COMPANY, Inc.

2 Jewel Drive, P.O. Box 696
Wilmington, MA 01887-0696
TEL: (508) 657-8750 FAX: (508) 658-0349 TELEX: 200079
IN EUROPE: Cryophysics SA, Geneva TEL: 41 22 732 95 20

Circle number 31 on Reader Service Card