

Some don't like it hot

Daniela Schmidt

University of Bristol, Wills Memorial Building, Queens Road, Bristol BS8 1RJ, UK

The oceans are experiencing vast environmental changes that are predicted to accelerate in the future (Stocker et al., 2013). Warming, acidification, deoxygenation, and increased stratification act on a global scale, whereas other factors such as eutrophication, the effect of changes in run-off on the carbonate system, and pollution act more locally to regionally. It is very hard to predict whether these drivers will act synergistically, antagonistically, or additively on marine organisms (Pörtner et al., 2014). Individual drivers can be tested in laboratory experiments (see Pörtner et al. [2014] for a recent assessment), but most of these experiments are too short, with little acclimation or possibility of adaptation. Species interact (Munday et al., 2009; Sanford et al., 2014) and evolve (Collins and Bell, 2004; Lohbeck et al., 2012), which is proving to be highly challenging to test in laboratory settings and mesocosms.

The geological record provides an archive of the integrated, long-term effects of climate change and ocean acidification, as well as other factors, on marine ecosystems (Hönisch et al., 2012). The Paleocene–Eocene Thermal Maximum (PETM), ca. 56 Ma, is a key interval for such comparisons because the effects of the climate perturbation can be found all over the globe and in all ecosystems (see McInerney and Wing [2011] and Sluijs et al. [2007] for reviews). A substantial negative carbon isotope excursion suggests the addition of between 2000–6000 Gt of isotopically light carbon to the atmosphere (Cui et al., 2011; Dickens, 2003). The global surface ocean warmed on average by 4–5 °C, and the subsurface ocean by 5–6 °C (Dunkley Jones et al., 2013). The observation of global shoaling of the carbonate compensation depth (Zachos et al., 2005), combined with recent modeling (Ridgwell and Schmidt, 2010) and boron isotope analysis (Penman et al., 2014) support that ocean acidification occurred in both the surface and the deep ocean. Ecosystem changes have been widely documented (Foster et al., 2013; Gibbs et al., 2006; Scheibner et al., 2005; Kelly et al., 1996; Thomas, 2007; Webb et al., 2009) showing, amongst others, migration toward higher latitudes, changes in ecosystem composition, extinction amongst deep sea species, and calcification responses.

The usefulness of the geological record in improving our understanding of the impacts of future climate change and ocean acidification, though, depends on accurate regional climate reconstructions, which allow a differentiated assessment of the impact on marine biota. Papers in this issue of *Geology*, by Aze et al. (2014, p. 739) and Frieling et al. (2014, p. 767), increase our knowledge in two critical areas: the tropical western Indian Ocean (19°S) and the subpolar West Siberian Seaway (~55°N), with the first PETM temperature reconstructions for these regions. Aze et al.'s novel tropical peak PETM values range from 32 °C to 43 °C, with a warming of 3 °C above background, depending on calibration and whether average or maximum values are considered. Similarly, warming is documented by Frieling et al. to have been ~7 °C, so that temperatures of 27 °C were reached in the West Siberian Seaway, combined with seasonal anoxia.

Both papers contain provocative novel ideas. For example, a complete lack of temperature difference between the Arctic and the West Siberian Seaway provides new targets for climate models. These papers also point to the challenges of working in comparatively shallower-water, near-coastal sections. Sediments at shallow-water sites are often subjected to reworking and unconformities, both of which make identifying base-lines of pre-event climate variability, and hence the relative amplitude of the warming, very difficult.

More importantly, though, both records point at our limitations to calculate absolute temperatures for deep-time records. Using oxygen isotopes (as in Aze et al.) meets the limits of our knowledge because seawater $\delta^{18}\text{O}$ is not well constrained, resulting in a several-degree uncertainty in temperature reconstructions (Tindall et al., 2010), as large as the climate signal in the event. This is especially true in settings with strong evaporation near the coast, and likely a high variability in the carbonate system, by analogy to modern shelf seas (Artioli et al., 2014). Additional effects such as unknown calibration equations for extinct species and the effect of the surface-water acidification on isotope incorporation just add to the problem (Spero et al., 1997). Given the very recent quantification of the surface-water pH values prior to the PETM and the change within (Penman et al., 2014), the most likely average sea-surface temperature for the PETM in Tanzania was between 33.9 °C and 35.9 °C, which agrees well with temperature ranges in model simulations (Huber and Caballero, 2011; Tindall et al., 2010) for pre-PETM background values, combined with the 3 °C warming found by Aze et al.

So if we take these data at face value, what are the consequences for biology, and what does this tell us about the future? These papers highlight the migration of phytoplankton to follow their niche, and suggest that the extreme warmth led to an absence of calcifiers in tropical waters. Intriguingly, though, this abiotic zone appears several tens of thousands of years after the onset of the extreme temperatures and the acidification, is associated with changes in lithology, and follows a gap in the record. This potentially slow response contradicts everything we know about ecosystem response to decadal temperature variability; for example, the North Atlantic Oscillation (Beaugrand et al., 2009; Beaugrand et al., 2002) or the California upwelling system (Chavez et al., 2003; Chavez et al., 1999). Aze et al. explain the abiotic zone by comparing it to the temperature adaptation of modern foraminifers. One would expect, though, that Paleogene foraminifers which evolved in an ~15 °C warmer environment than today (Huber and Caballero, 2011) were generally adapted to these warmer temperatures. As so often, new papers ask more questions than they answer, such as: why are these abiotic zones not found at other open ocean sites nearer the equator? If the high-end temperatures are reasonable estimates, these might point to physiological limits at which enzymes start denaturalizing. Given the high metabolic rates in response to these high temperatures, the size of the supply of food needed to sustain the organisms is a pressing question and might have played a role in a regional exclusion. More work is needed, though, to move from assessments of past climates to predictive models for policy makers of the impact of future climate change on marine ecosystems, such as the cascading effects of these potential abiotic zones on food webs.

ACKNOWLEDGMENTS

I thank the Royal Society for funding via a University Research Fellowship.

REFERENCES CITED

- Artioli, Y., Blackford, J.C., Nondal, G., Bellerby, R.G.J., Wakelin, S.L., Holt, J.T., Butenschön, M., and Allen, J.I., 2014, Heterogeneity of impacts of high CO₂ on the North Western European shelf: *Biogeosciences*, v. 11, p. 601–612, doi:10.5194/bg-11-601-2014.
- Aze, T., Pearson, P.N., Dickson, A.J., Badger, M.P.S., Bown, P.R., Pancost, R.D., Gibbs, S.J., Huber, B.T., Leng, M.J., Coe, A.L., Cohen, A.S., and Foster, G.L., 2014, Extreme warming of tropical waters during the Paleocene–Eocene Thermal Maximum: *Geology*, v. 42, p. 739–742, doi:10.1130/G35637.1.

- Beaugrand, G., Luczak, C., and Edwards, M., 2009, Rapid biogeographical plankton shifts in the North Atlantic Ocean: *Global Change Biology*, v. 15, p. 1790–1803, doi:10.1111/j.1365-2486.2009.01848.x.
- Beaugrand, G., Reid, P.C., Ibanez, F., Lindley, J.A., and Edwards, M., 2002, Reorganization of North Atlantic marine copepod biodiversity and climate: *Science*, v. 296, p. 1692–1694, doi:10.1126/science.1071329.
- Chavez, F.P., Ryan, J., Lluch-Cota, S.E., and Niquen, C.M., 2003, From anchovies to sardines and back: Multidecadal change in the Pacific Ocean: *Science*, v. 299, p. 217–221, doi:10.1126/science.1075880.
- Chavez, F.P., Strutton, P.G., Friederich, G.E., and Feely, R.A., 1999, Biological and chemical response of the equatorial Pacific Ocean to the 1997/98 El Niño: *Science*, v. 286, p. 2126–2131, doi:10.1126/science.286.5447.2126.
- Collins, S., and Bell, G., 2004, Phenotypic consequences of 1,000 generations of selection at elevated CO₂ in a green alga: *Nature*, v. 431, p. 566–569, doi:10.1038/nature02945.
- Cui, Y., Kump, L.R., Ridgwell, A.J., Charles, A.J., Junium, C.K., Diefendorf, A.F., Freeman, K.H., Urban, N.M., and Harding, I.C., 2011, Slow release of fossil carbon during the Palaeocene-Eocene Thermal Maximum: *Nature Geoscience*, v. 4, p. 481–485, doi:10.1038/ngeo1179.
- Dickens, G.R., 2003, Rethinking the global carbon cycle with a large, dynamic and microbially mediated gas hydrate capacitor: *Earth and Planetary Science Letters*, v. 213, p. 169–183, doi:10.1016/S0012-821X(03)00325-X.
- Dunkley Jones, T., Lunt, D.J., Schmidt, D.N., Ridgwell, A., Sluijs, A., Valdes, P.J., and Maslin, M., 2013, Climate model and proxy data constraints on ocean warming across the Paleocene-Eocene Thermal Maximum: *Earth-Science Reviews*, v. 125, p. 123–145, doi:10.1016/j.earscirev.2013.07.004.
- Foster, L.C., Schmidt, D.N., Thomas, E., Arndt, S., and Ridgwell, A., 2013, Surviving rapid climate change in the deep sea during the Paleogene hyperthermals: *Proceedings of the National Academy of Sciences of the United States of America*, v. 110, p. 9273–9276, doi:10.1073/pnas.1300579110.
- Frieling, J., Iakovleva, A.I., Reichert, G.J., Aleksandrova, G.N., Gnibidenko, Z.N., Schouten, S., and Sluijs, A., 2014, Paleocene–Eocene warming and biotic response in the epicontinental West Siberian Sea: *Geology*, v. 42, p. 767–770, doi:10.1130/G35724.1.
- Gibbs, S.J., Bown, P.R., Sessa, J.A., Bralower, T.J., and Wilson, P.A., 2006, Nanoplankton extinction and origination across the Paleocene-Eocene Thermal Maximum: *Science*, v. 314, p. 1770–1773, doi:10.1126/science.1133902.
- Hönisch, B., et al., 2012, The geological record of ocean acidification: *Science*, v. 335, p. 1058–1063, doi:10.1126/science.1208277.
- Huber, M., and Caballero, R., 2011, The early Eocene equable climate problem revisited: *Climate of the Past*, v. 7, p. 603–633, doi:10.5194/cp-7-603-2011.
- Kelly, D.C., Bralower, T.J., Zachos, J.C., Premoli-Silva, I., and Thomas, E., 1996, Rapid diversification of planktonic foraminifera in the tropical Pacific (ODP Site 865) during the late Paleocene thermal maximum: *Geology*, v. 24, p. 423–426, doi:10.1130/0091-7613(1996)024<0423:RDOPFI>2.3.CO;2.
- Lohbeck, K.T., Riebesell, U., and Reusch, T.B.H., 2012, Adaptive evolution of a key phytoplankton species to ocean acidification: *Nature Geoscience*, v. 5, p. 346–351, doi:10.1038/ngeo1441.
- McInerney, F.A., and Wing, S.L., 2011, The Paleocene-Eocene Thermal Maximum: A perturbation of carbon cycle, climate, and biosphere with implications for the future: *Annual Review of Earth and Planetary Sciences*, v. 39, p. 489–516, doi:10.1146/annurev-earth-040610-133431.
- Munday, P.L., Dixson, D.L., Donelson, J.M., Jones, G.P., Pratchett, M.S., Devitsina, G.V., and Døving, K.B., 2009, Ocean acidification impairs olfactory discrimination and homing ability of a marine fish: *Proceedings of the National Academy of Sciences of the United States of America*, v. 106, p. 1848–1852, doi:10.1073/pnas.0809996106.
- Penman, D.E., Hönisch, B., Zeebe, R.E., Thomas, E., and Zachos, J.C., 2014, Rapid and sustained surface ocean acidification during the Paleocene-Eocene Thermal Maximum: *Paleoceanography*, v. 29, doi:10.1002/2014PA002621.
- Pörtner, H.O., Karl, D., Boyd, P.W., Cheung, W.W.L., Lluch-Cota, S.E., Nojiri, Y., Schmidt, D.N., and Zavialov, P., 2014, Ocean Systems, in Field, C. B., and Barros, V., eds., *Impacts, Adaptation and Vulnerability, Volume II: Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*: Cambridge, UK, Cambridge University Press.
- Ridgwell, A., and Schmidt, D.N., 2010, Past constraints on the vulnerability of marine calcifiers to massive carbon dioxide release: *Nature Geoscience*, v. 3, p. 196–200, doi:10.1038/ngeo755.
- Sanford, E., Gaylord, B., Hettinger, A., Lenz, E.A., Meyer, K., and Hill, T.M., 2014, Ocean acidification increases the vulnerability of native oysters to predation by invasive snails: *Proceedings of the Royal Society B*, v. 281, no. 1778, doi:10.1098/rspb.2013.2681.
- Scheibner, C., Speijer, R.P., and Marzouk, A.M., 2005, Turnover of larger foraminifera during the Paleocene–Eocene Thermal Maximum and paleoclimatic control on the evolution of platform ecosystems: *Geology*, v. 33, p. 493–496, doi:10.1130/G21237.1.
- Sluijs, A., Bowen, G.J., Brinkhuis, H., Lourens, L.J., and Thomas, E., 2007, The Palaeocene-Eocene Thermal Maximum super greenhouse: Biotic and geochemical signatures, age models and mechanisms of global change, in Williams, M., et al., *Deep Time Perspectives on Climate Change—Marrying the Signal from Computer Models and Biological Proxies: The Micropalaeontological Society, Special Publications*, The Geological Society of London, p. 323–351.
- Spero, H.J., Bijma, J., Lea, D.W., and Bemis, B.E., 1997, Effect of seawater carbonate concentration on foraminiferal carbon and oxygen isotopes: *Nature*, v. 390, p. 497–500, doi:10.1038/37333.
- Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P.M., 2013, *Climate Change 2013: The Physical Science Basis: Working Group I Contribution to the Fifth Assessment Report of the IPCC*: Cambridge, UK, Cambridge University Press, 1535 p.
- Thomas, E., 2007, Cenozoic mass extinctions in the deep sea: What perturbs the largest habitat on earth? in Monechi, S., et al., eds., *Large Ecosystem Perturbations: Causes and Consequences*: Boulder, Colorado, Geological Society of America Special Paper 424, p. 1–23.
- Tindall, J., Flecker, R., Valdes, P., Schmidt, D.N., Markwick, P., and Harris, J., 2010, Modeling the oxygen isotope distribution of ancient seawater using a coupled ocean-atmosphere GCM: Implications for reconstructing early Eocene climate: *Earth and Planetary Science Letters*, v. 292, p. 265–273, doi:10.1016/j.epsl.2009.12.049.
- Webb, A.E., Leighton, L.R., Schellenberg, S.A., Landau, E.A., and Thomas, E., 2009, Impact of the Paleocene-Eocene Thermal Maximum on deep-ocean microbial community structure: Using rank-abundance curves to quantify paleoecological response: *Geology*, v. 37, p. 783–786, doi:10.1130/G30074A.1.
- Zachos, J.C., et al., 2005, Rapid acidification of the ocean during the Paleocene-Eocene Thermal Maximum: *Science*, v. 308, p. 1611–1615, doi:10.1126/science.1109004.

Printed in USA