

# Titanium Dioxide Nanoparticles Induce DNA Damage and Genetic Instability *In vivo* in Mice

Benedicte Trouiller,<sup>1</sup> Ramune Reliene,<sup>1,3</sup> Aya Westbrook,<sup>1,4</sup>  
Parrisa Solaimani,<sup>1,4</sup> and Robert H. Schiestl<sup>1,2,5</sup>

Departments of <sup>1</sup>Pathology and Laboratory Medicine, <sup>2</sup>Radiation Oncology, and <sup>3</sup>Medicine, Center for Human Nutrition, David Geffen School of Medicine; <sup>4</sup>Interdepartmental Program in Molecular Toxicology; <sup>5</sup>Department of Environmental Health Sciences, School of Public Health, University of California-Los Angeles, Los Angeles, California

## Abstract

**Titanium dioxide (TiO<sub>2</sub>) nanoparticles are manufactured worldwide in large quantities for use in a wide range of applications including pigment and cosmetic manufacturing. Although TiO<sub>2</sub> is chemically inert, TiO<sub>2</sub> nanoparticles can cause negative health effects, such as respiratory tract cancer in rats. However, the mechanisms involved in TiO<sub>2</sub>-induced genotoxicity and carcinogenicity have not been clearly defined and are poorly studied *in vivo*. The present study investigates TiO<sub>2</sub> nanoparticles–induced genotoxicity, oxidative DNA damage, and inflammation in a mice model. We treated wild-type mice with TiO<sub>2</sub> nanoparticles in drinking water and determined the extent of DNA damage using the comet assay, the micronuclei assay, and the  $\gamma$ -H2AX immunostaining assay and by measuring 8-hydroxy-2'-deoxyguanosine levels and, as a genetic instability endpoint, DNA deletions. We also determined mRNA levels of inflammatory cytokines in the peripheral blood. Our results show that TiO<sub>2</sub> nanoparticles induced 8-hydroxy-2'-deoxyguanosine,  $\gamma$ -H2AX foci, micronuclei, and DNA deletions. The formation of  $\gamma$ -H2AX foci, indicative of DNA double-strand breaks, was the most sensitive parameter. Inflammation was also present as characterized by a moderate inflammatory response. Together, these results describe the first comprehensive study of TiO<sub>2</sub> nanoparticles–induced genotoxicity *in vivo* in mice possibly caused by a secondary genotoxic mechanism associated with inflammation and/or oxidative stress. Given the growing use of TiO<sub>2</sub> nanoparticles, these findings raise concern about potential health hazards associated with TiO<sub>2</sub> nanoparticles exposure. [Cancer Res 2009;69(22):8784–9]**

## Introduction

Titanium dioxide (TiO<sub>2</sub>) accounts for 70% of the total production volume of pigments worldwide (1). It is widely used to provide whiteness and opacity to products such as paints, plastics, papers, inks, food colorants, and toothpastes. TiO<sub>2</sub> is also used in cosmetic and skin care products, particularly in sunblocks, where it helps to protect the skin from UV light, especially in the case of nanosized particles (<100 nm). Nevertheless, TiO<sub>2</sub> has recently been reclassified by the IARC as group 2B carcinogen: “possibly carcinogenic to


humans”<sup>6</sup>. The reason for this new classification stems from the fact that high concentrations of pigment-grade (<2.5  $\mu$ m) and ultrafine (<100 nm) TiO<sub>2</sub> dust can cause respiratory tract cancer in exposed rats (2, 3). However, it should be noted that epidemiologic studies of workers exposed to pigment-grade TiO<sub>2</sub> conducted thus far have not been able to detect an association between occupational exposure to TiO<sub>2</sub> and an increased risk for lung cancer (4, 5). Genotoxicity studies that measure different types of DNA damage (e.g., gene mutations, chromosomal damage, and DNA strand break formation) are an important part of cancer research and risk assessment of potential carcinogens. These studies help to understand possible mechanisms causing tumor induction. As such, *in vivo* mechanisms underlying TiO<sub>2</sub> nanoparticles tumor induction are still unclear.

Because nanoparticle diameter does not exceed a hundred nanometers at maximum, they are able to penetrate cells (6) and interfere with several subcellular mechanisms. Indeed, some studies show that some nanoparticles can penetrate into cell nuclei and hence may directly interfere with the structure and function of genomic DNA (7). Additionally, after oral administration in mice, TiO<sub>2</sub> particles were shown to translocate to systemic organs such as liver and spleen as well as lung and peritoneal tissues (8). Genotoxicity studies have been done to understand the carcinogenic potential of TiO<sub>2</sub> nanoparticles using assays that measure mutations in genes (e.g., Ames/*Salmonella* and hypoxanthine guanine phosphoribosyl transferase (Hprt) assays; refs. 9–11), chromosomal damage representing possible clastogenic activity of the particles (e.g., micronuclei; refs. 10, 12–15), and DNA strand breakage (e.g., alkaline comet assay; refs. 10, 13). Except for one, these studies were conducted *in vitro* in cultured cells but conflict in their results. Half of the studies show that TiO<sub>2</sub> nanoparticles are genotoxic in cell lines (10, 12, 13, 15), whereas the other half show that TiO<sub>2</sub> nanoparticles are not (9, 11, 14). The rationale for these conflicting results is not clear because different cell types, doses, and nanoparticle sizes have been used. Some studies suggest possible mechanisms for TiO<sub>2</sub> nanoparticles genotoxicity. TiO<sub>2</sub> nanoparticles might damage DNA directly or indirectly via oxidative stress and/or inflammatory responses. Two recent studies show a direct chemical interaction between TiO<sub>2</sub> nanoparticles and DNA, through the DNA phosphate group, but a link to mutagenesis has not been proven (16, 17). On the other hand, other studies show that TiO<sub>2</sub> nanoparticles can cause DNA damage indirectly through inflammation (18–21) and generation of reactive oxygen species (12, 13, 22, 23).

**Requests for reprints:** Robert H. Schiestl, Department of Pathology, University of California-Los Angeles School of Medicine, 650 Charles E. Young Drive South, Los Angeles, CA 90095. Phone: 310-267-2087; Fax: 310-267-2578; E-mail: rschiestl@mednet.ucla.edu.

©2009 American Association for Cancer Research.  
doi:10.1158/0008-5472.CAN-09-2496

<sup>6</sup> IARC. Monographs on the evaluation of carcinogenic risks to humans. Carbon black, titanium dioxide and non-asbestiform talc. Lyon (France): IARC. In press.



**Figure 1.** Frequency of DNA deletions in control and TiO<sub>2</sub> nanoparticles-treated mice. One RPE corresponds to one eye. Mice were treated with nanoparticles during embryonic development at a total dose of 500 mg/kg. Mean ± SE numbers of eyespots per RPE with  $n = 42$  eyes for control and  $n = 53$  eyes for TiO<sub>2</sub> nanoparticles-treated mice. \*,  $P < 0.05$ .

Thus far, most nanoparticle genotoxicity studies have focused on cell culture systems, but confirmation from animal experiments, more relevant to human exposure, is required. To further understand TiO<sub>2</sub> nanoparticles toxicity *in vivo*, we studied the effect of TiO<sub>2</sub> nanoparticles exposure on genotoxicity, DNA damage, and inflammation in mice. To evaluate inflammation in mice, we determined mRNA expression of both proinflammatory and anti-inflammatory cytokines. To assess DNA damage, we used the  $\gamma$ -H2AX and the comet assays to evaluate DNA strand breaks, the micronuclei assay to estimate chromosomal damage, and the measure of 8-hydroxy-2'-deoxyguanosine (8-OHdG) levels using high-performance liquid chromatography to determine oxidative DNA damage. We also used an *in vivo* DNA deletion assay, which allows visual detection of DNA deletion events within the *pink-eyed unstable* ( $p^{un}$ ) locus in developing mouse embryos (24), which can detect environmental as well as genetic cancer predisposing factors (24). Our results show that TiO<sub>2</sub> P25 nanoparticles can induce 8-OHdG,  $\gamma$ -H2AX foci, micronuclei, DNA deletions, and inflammation markers in a mice model. Therefore, this study suggests that TiO<sub>2</sub> nanoparticles are genotoxic *in vivo*.

## Materials and Methods

**Mouse care and breeding.** C57Bl/6J $p^{un}/p^{un}$  mice were obtained from The Jackson Laboratory. The C57Bl/6J $p^{un}/p^{un}$  background is essentially identical to C57Bl/6J, with the exception of a naturally occurring 70-kb internal duplication in the *pink-eyed dilution* ( $p$ ) gene, termed the  $p^{un}$  allele. Mice were housed and cared for under standard specific pathogen-free conditions and according to the Animal Rescue Coalition and Institutional Animal Care and Use Committee regulations. Mice were given a standard, autoclaved diet from Harlan Teklad (Harlan Teklad No. 8656) and sterilized water *ad libitum*. Mice were housed in a 12-h light/dark cycle. Pregnancy was timed by checking for vaginal plugs, with noon of the day of discovery counted as 0.5 days post-coitum. Four- to 5-month-old mice were used for all experiments.

**TiO<sub>2</sub> nanoparticles preparation and exposure.** "Aeroxide" P25 TiO<sub>2</sub> (Degussa, now Evonik) nanoparticles were chosen for this study. The crystal structure is a mixture of 75% anatase and 25% rutile TiO<sub>2</sub>, purity was at least 99.5% TiO<sub>2</sub>, and primary particle size was 21 nm with a specific surface area of  $50 \pm 15$  m<sup>2</sup>/g. These nanoparticles have been used in many of the previous mammalian studies (13, 14, 22, 25–28). Using dynamic light scattering in water revealed that the size of TiO<sub>2</sub> nanoparticles agglomerates ranged from 21 to 1,446 nm and the mean size was


160 ± 5 nm. About 70% of particles have a size of 160 nm. Solutions of dispersed TiO<sub>2</sub> nanoparticles were prepared by ultrasonication (Solid State/Ultrasonic FS-14; Fisher Scientific) for 15 min in drinking water at 60, 120, 300, and 600  $\mu$ g/mL concentrations just before use. We measured TiO<sub>2</sub> nanoparticles-supplemented water intake at the end of experiments in each cage, which housed 2 to 3 mice, and calculated an average daily water intake per mouse. Daily TiO<sub>2</sub> nanoparticles-supplemented water intake ranged from 3 to 7 mL/mouse, consistent with normal daily water intake. Doses were calculated using a 30 g average weight per mouse, and an average of 5 mL water intake per day. The exposure was 5 days in adult males. For *in utero* exposure, pregnant dams were given nanoparticles-supplemented drinking water for 10 days from 8.5 to 18.5 days post-coitum at a concentration of 300  $\mu$ g/mL. Water was used as negative control.

***In vivo* DNA deletion assay.** To evaluate genotoxicity of TiO<sub>2</sub> nanoparticles, we employed an intrachromosomal duplication of 70-kb fragment spanning exons 6 to 18 of the  $p$  gene in mice (termed  $p^{un}$  mutation). When a DNA deletion event occurs between these duplications, the  $p^{un}$  allele reverts to the wild-type  $p$  gene. Reconstitution of the wild-type  $p$  gene can be seen as a single pigmented cell or a clone of pigmented cells on the unpigmented retinal pigment epithelium (RPE) in the transgenic mice and represents a DNA deletion as a permanent genotoxic event (29). Pregnant mice were treated with TiO<sub>2</sub> nanoparticles, and offspring were sacrificed at age 20 days. Their eyes were extracted and dissected to display the RPE for the deletion/eyespot assay as described previously (24). One RPE corresponds to one eye.


**Alkaline comet assay.** Peripheral blood was collected by submandibular vein puncture (before treatment and after treatment) in an EDTA-coated tube. The comet assay was done as described previously (30). On average, from three slides, 150 to 200 randomly captured comets per sample were analyzed. Results were expressed as average ± SE tail moment.

**Micronuclei assay.** The micronuclei assay was done as described elsewhere (31). Three microliter aliquots of the peripheral blood were collected as described above and smeared on slides and stained into Giemsa stain for 1.5 min. Approximately 2,000 erythrocytes were scored per animal to estimate the frequency of micronucleated erythrocytes.

**Bone marrow preparation.** Animals were sacrificed with an overdose of isoflurane after 5 days of treatment with TiO<sub>2</sub> nanoparticles in drinking water. Both femora were dissected, and marrow cells were flushed out with 1 mL PBS and pipetted several times. The cell suspension was centrifuged at 1,000 rpm for 5 min, the supernatant was withdrawn, and the cell pellet was resuspended and placed on a clean glass slide.



**Figure 2.** Percentage of  $\gamma$ -H2AX-positive cells in bone marrow in untreated and TiO<sub>2</sub> nanoparticles-treated mice and a picture of a  $\gamma$ -H2AX-positive cell with more than four foci. Columns, mean of 5 mice; bars, SE. \*\*\*,  $P < 0.001$ , TiO<sub>2</sub> nanoparticles-treated versus control.



**Figure 3.** Frequency of DNA strand breaks in mice before and after treatment with 500 mg/kg TiO<sub>2</sub> nanoparticles. DNA damage is represented by the tail moment. Mean ± SE ( $n = 5$  mice/group). \*,  $P < 0.05$ , compared with untreated mice.

**$\gamma$ -H2AX assay, RNA isolation, and quantitative real-time PCR.** The  $\gamma$ -H2AX assay was done with bone marrow cells, and the RNA isolation for quantitative real-time PCR was carried out on peripheral blood. These assays were done as described elsewhere (31).

**Determination of oxidative DNA damage by measuring 8-OHdG.** Mouse livers were isolated just after 5 days of treatment with nanoparticles and immediately frozen in liquid nitrogen and homogenized under liquid nitrogen. 8-OHdG level was measured using high-performance liquid chromatography with electron capture detection system as described previously (32).

**Statistical analysis.** For the deletion assay, the comet assay, the micronuclei assay, and 8-OHdG and mRNA levels of cytokines, we used the Student's  $t$  test to compare untreated mice with treated mice. For the  $\gamma$ -H2AX experiment, the percentage of positive cells for control groups versus treated groups was compared via  $\chi^2$  test. In addition to the  $t$  test, for comet assay data, the Wilcoxon test for matched paired data was also used to compare the effect of TiO<sub>2</sub> nanoparticles on the tail moment before and after nanoparticle treatment. The difference was considered significant at the 95% confidence level ( $P < 0.05$ ) and highly significant at the 99% confidence level ( $P \leq 0.01$ ).

## Results

**TiO<sub>2</sub> nanoparticles increased the frequency of DNA deletions.** We used the DNA deletion assay to evaluate *in vivo* genotoxicity of TiO<sub>2</sub> nanoparticles. We quantified the number of eyespots per RPE as a measure of DNA deletions in *in utero* exposed mice. TiO<sub>2</sub> nanoparticles-treated mice had an average of  $8.13 \pm 1.70$  eyespots per RPE versus  $6.42 \pm 1.47$  eyespots per RPE in nontreated mice (Fig. 1). TiO<sub>2</sub> nanoparticles-exposed mice displayed a significant increase in eyespots (27%) compared with unexposed mice ( $P = 0.019$ ), suggesting that, after maternal oral exposure, TiO<sub>2</sub> nanoparticles increased DNA deletion frequency in fetuses.

**TiO<sub>2</sub> nanoparticles induced  $\gamma$ -H2AX foci.** Phosphorylation of histone H2AX on serine 139 occurs at sites flanking DNA double-strand breaks (DSB), providing a measure of the number of DSBs within a cell (33). We used this assay to compare DSB formation in bone marrow of mice with and without TiO<sub>2</sub> nanoparticles treatment.

The  $\gamma$ -H2AX foci formation increased by ~10%, 20%, 25%, and 30% following treatment with 50, 100, 250, and 500 mg/kg TiO<sub>2</sub> nanoparticles, respectively, compared with untreated mice (Fig. 2). Percentage of  $\gamma$ -H2AX-positive cells increased with TiO<sub>2</sub> nanoparticles concentration in a clear dose-dependent manner ( $P < 0.001$ ).


These data provided evidence that, after oral administration, TiO<sub>2</sub> nanoparticles induce DSBs in bone marrow cells.

**TiO<sub>2</sub> nanoparticles increased DNA strand breaks.** DNA strand breaks (DSBs, single-strand breaks, and/or strand breaks induced by alkali-labile sites) were measured by the alkaline comet assay in mice peripheral blood before and after treatment. Tail moment significantly increased after TiO<sub>2</sub> nanoparticles treatment (Fig. 3). The average tail moment was  $0.0102 \pm 0.001$  before treatment and  $0.0137 \pm 0.0011$  after TiO<sub>2</sub> nanoparticles treatment. TiO<sub>2</sub> nanoparticles increased DNA strand breaks in WBCs from peripheral blood by 34% ( $P = 0.001$ ,  $t$  test, and  $P = 0.04$ , Wilcoxon test).


**TiO<sub>2</sub> nanoparticles induced micronuclei.** The micronuclei assay was used to detect chromosomal damage in erythrocytes from peripheral blood. The incidence of micronuclei serves as an index of clastogenicity. Micronuclei frequency increased significantly only at the highest (500 mg/kg) dose of TiO<sub>2</sub> nanoparticles used ( $P = 0.009$ ; Fig. 4). At this dose, the average micronuclei frequencies for untreated mice were  $4.3 \pm 0.93$  versus  $9.2 \pm 1.07$  per 2,000 RBC for TiO<sub>2</sub> nanoparticles-treated mice, which resulted in a 2.1-fold increase in micronuclei formation. This result showed that, at high dose, TiO<sub>2</sub> nanoparticles induced detectable clastogenicity in mice peripheral blood.

**TiO<sub>2</sub> nanoparticles induced oxidative DNA damage.** We examined the degree of oxidative DNA damage by measuring the level of 8-OHdG in DNA isolated from TiO<sub>2</sub> nanoparticles-treated and untreated mouse livers. The level of 8-OHdG was significantly higher in TiO<sub>2</sub> nanoparticles-treated than untreated mice ( $P = 0.04$ ; Fig. 5). The average number of 8-OHdG per  $10^6$  dG was  $4.25 \pm 0.66$  for untreated mice and  $6.43 \pm 0.58$  for TiO<sub>2</sub> nanoparticles-treated mice resulting in a 1.5-fold increase at 500 mg/kg TiO<sub>2</sub> nanoparticles. This suggested that TiO<sub>2</sub> nanoparticles induced oxidative DNA damage in liver.

**TiO<sub>2</sub> nanoparticles induced a proinflammatory response.** We quantified mRNA transcripts of three Th1/proinflammatory cytokines (T-helper cell type 1) and three Th2/anti-inflammatory cytokines (T-helper cell type 2) in the peripheral blood. After treatment, the proinflammatory cytokines tumor necrosis factor- $\alpha$ , IFN- $\gamma$ , and the mouse orthologue of interleukin-8 (KC) were significantly upregulated ( $P = 0.01$ , 0.02, and 0.05, respectively; Fig. 6A).



**Figure 4.** Frequency of micronuclei in mice before and after TiO<sub>2</sub> nanoparticles treatment in peripheral blood erythrocytes. Open columns, untreated controls; gray columns, TiO<sub>2</sub> nanoparticles-treated mice. Columns, mean of 5 mice; bars, SE. \*,  $P < 0.01$ , compared with untreated mice.



**Figure 5.** Level of 8-OHdG in untreated and 500 mg/kg TiO<sub>2</sub> nanoparticles-treated mouse livers. Mean ± SE ( $n = 5$  mice/group). \*,  $P < 0.05$ , compared with untreated mice.

A general upregulation of these cytokines may be due to the effects of circulating TiO<sub>2</sub> nanoparticles directly in the peripheral blood, suggesting systemic distribution, and direct activation of a proinflammatory response. In the contrary, anti-inflammatory cytokines with generally opposing function were not upregulated, including transforming growth factor- $\beta$ , interleukin-10, and interleukin-4 (Fig. 6B). TiO<sub>2</sub> nanoparticles did not induce an anti-inflammatory response, which mean they did not inhibit the production and release of proinflammatory mediators.


## Discussion

Here, we report for the first time that TiO<sub>2</sub> nanoparticles are genotoxic and clastogenic *in vivo* in mice. We showed that TiO<sub>2</sub> nanoparticles (500 mg/kg) induce not only DNA single-strand breaks and DSBs but also chromosomal damage. The formation of  $\gamma$ -H2AX foci, which show DSB formation, was the most sensitive parameter and showed a consistent dose-dependent response. Concerning health relevance, DSBs are much more damaging in terms of genetic instability than single-strand breaks and oxidative DNA damage, which are transient. Our results extend previous *in vitro* findings with the micronuclei and comet assays in several human cells and Syrian hamster embryo cells (10, 12, 13, 15),

although they have not been detected in some studies (11, 14). Differences in response between studies may be due to how TiO<sub>2</sub> nanoparticles differ in terms of TiO<sub>2</sub> production, particle size, degree of aggregation, preparation method (sonication), incubation conditions, dose, and susceptibility between cell types (34, 35), implying that more studies are needed to determine the conditions in which TiO<sub>2</sub> nanoparticles genotoxicity occurs.

To date, very few *in vivo* genotoxicity studies have been carried out with nanoparticles. A chronic exposure to TiO<sub>2</sub> nanoparticles at concentrations that produce chronic pulmonary inflammation was associated with an increased incidence of tumors in rat lungs (3). Thus far, only two *in vivo* genotoxicity studies have been reported, which showed that *in vivo* TiO<sub>2</sub> nanoparticles increased *Hprt* mutation frequency in alveolar epithelial cells (9) but did not induce DNA adduct formation in rat lungs (26). Our study showed for the first time that, *in vivo* after oral exposure, TiO<sub>2</sub> nanoparticles induce DNA strand breaks and chromosomal damage in bone marrow and/or peripheral blood, which may help to further understand potential mechanisms of TiO<sub>2</sub> nanoparticles carcinogenicity. We also found that maternal exposure to 500 mg/kg TiO<sub>2</sub> nanoparticles during gestation results in significantly elevated frequencies of DNA deletions in offspring. This result is a major finding because it shows for the first time that *in utero* exposure of fetuses via the mother causes an increase in large deletions in offspring. Taken together, our findings show that TiO<sub>2</sub> nanoparticles, orally administrated, induce genotoxicity systemically in organs, such as blood, bone marrow, and even the embryo.

Surprisingly, human studies have not been able to detect any relation between TiO<sub>2</sub> occupational exposure and cancer risk (4, 5, 36), but these studies have methodologic and epidemiologic limitations as reviewed by the National Institute of Occupational Safety and Health (37). In addition, these studies were not designed to investigate the relationship between TiO<sub>2</sub> particle size and lung cancer risk, an important question for assessing the potential occupational carcinogenicity of TiO<sub>2</sub>. Indeed, Dankovic and colleagues, comparing several tumor studies, pointed out that TiO<sub>2</sub> nanoparticles produced lung tumors in rats at a much lower dose than fine particles (<250 nm; 10 and 250 mg/m<sup>3</sup> for nanoparticles and fine particles, respectively; ref. 3). Although TiO<sub>2</sub> nanoparticles are prone to forming agglomerates of >100 nm in solution, these agglomerates are apparently not stable and appear to dissociate in



**Figure 6.** TiO<sub>2</sub> nanoparticles at 500 mg/kg induce proinflammatory but not anti-inflammatory cytokines. Open columns, untreated controls; black columns, TiO<sub>2</sub> nanoparticles-treated mice. A, expression of proinflammatory cytokine panel relative to TBP, the internal control gene. \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ , Student's *t* test for treatment comparisons. B, expression of anti-inflammatory cytokine panel relative to TBP. Student's *t* test revealed no significant differences. For each graph,  $n = 5$  mice/group.

bodily fluids and tissues, which could be an explanation for TiO<sub>2</sub> nanoparticles higher toxicity. It has also been reported that TiO<sub>2</sub> nanoparticles surface interactions are weak (20). In addition, an inhalation study showed that TiO<sub>2</sub> nanoparticles agglomerates of ~700 nm dissociate into smaller units after deposition in the lung (21). Even if nanoparticles aggregate into larger-sized agglomerates, their primary particle sizes remain a significant trait that affects their toxicity. Further human studies would be necessary to understand nanoparticle health effects. For instance, one could use blood-based assays similar to those done in this study in a future molecular epidemiology study in occupational settings. Furthermore, TiO<sub>2</sub> is also used in food colorants and toothpaste. This supports the notion that nanoparticle ingestion constitutes a relevant route of exposure in humans and underscores the significance of the findings of our study. In addition, given the capability of nanoparticles to enter the systemic blood circulation, nanoparticles may pose hazard to a variety of other organs as we have shown here.

As suggested previously, a possible mechanism for nanoparticles-caused genotoxicity might be via oxidative stress (38). Indeed, previous studies showed that TiO<sub>2</sub> nanoparticles have hydroxyl radical activity (39–41) and can also trigger reactive oxygen species production in different cell lines (13, 42) on interaction with the cell membrane or even in cell-free environment (23, 43). We confirmed these results in our *in vivo* experiment by showing oxidative DNA damage (8-OHdG) increase in mouse livers after TiO<sub>2</sub> nanoparticles treatment. Also, TiO<sub>2</sub> nanoparticles–increased DNA deletions during fetal development might be a result of oxidative genome damage. As discussed previously (44), oxidative stress is particularly hazardous in replicating cells. For instance, oxidative DNA lesions (e.g., 8-OHdG, single-strand breaks, or stalled replication forks) can lead to DSBs after replication, which can result in recombination, thus producing permanent genome rearrangements. As shown in yeast, oxidative mutagens might be powerful inducers of DNA deletions (45). Because embryonic cells are generally characterized by a high replication index, they might be particularly susceptible to oxidative genome damage.

We have also observed an inflammatory reaction as shown by changes in cytokine expression in peripheral blood, in which TiO<sub>2</sub> nanoparticles could be exerting direct inflammatory effects on circulating innate cells and Th1 effector cells. The inflammatory response involving recruitment and activation of phagocytes is

capable of causing oxidative bursts that may serve as a possible explanation for the observed genotoxicity to peripheral leukocytes, micronuclei formation, oxidative DNA damage in liver cells, and DNA deletion induction in fetal RPE. Because we showed that in mice TiO<sub>2</sub> nanoparticles induce an inflammatory reaction and oxidative DNA damage, it is tempting to speculate that the mechanism underlying TiO<sub>2</sub> nanoparticles genotoxicity might be a secondary (indirect) genotoxicity pathway as suggested by Dankovic and colleagues (3). Secondary genotoxicity is considered to be the key aspect of some particle toxicity (e.g., quartz and silica) because of their ability to elicit persistent inflammation *in vivo* (9, 46). This implies that particles have prooxidant and proinflammatory activity, leading to genotoxicity.

In summary, our study showed for the first time that TiO<sub>2</sub> nanoparticles induce clastogenicity, genotoxicity, oxidative DNA damage, and inflammation *in vivo* in mice. These results have been observed after only 5 days of treatment via drinking water and in multiple organs suggesting a systemic effect. We also showed that *in utero* exposure to TiO<sub>2</sub> nanoparticles results in an increased frequency in DNA deletions in the fetus. These results represent the first comprehensive *in vivo* genotoxicity study of TiO<sub>2</sub> nanoparticles. These data suggest that we should be concerned about a potential risk of cancer or genetic disorders especially for people occupationally exposed to high concentrations of TiO<sub>2</sub> nanoparticles and that it might be prudent to limit ingestion of TiO<sub>2</sub> nanoparticles through nonessential drug additives, food colors, etc.

## Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

## Acknowledgments

Received 7/20/09; revised 8/22/09; accepted 9/10/09; published OnlineFirst 11/3/09.

**Grant support:** NIH grants RO1 ES09519 (R.H. Schiestl) and 1RO3 CA133928-01 (R. Reliene).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

We thank Susanne Henning (University of California-Los Angeles Center for Human Nutrition) for performing high-performance liquid chromatography; Dr. Gopal Iyer (Department of Chemistry and Biochemistry, University of California-Los Angeles) for dynamic light scattering measurements; and Drs. Fabien Nogué, Norman Arnheim, and, especially, Jean-Louis Plouhinec for very interesting comments on the article.

## References

- Baan R, Straif K, Grosse Y, et al. Carcinogenicity of carbon black, titanium dioxide, and talc. *Lancet Oncol* 2006;7:295–6.
- Born PJ, Schins RP, Albrecht C. Inhaled particles and lung cancer, part B: paradigms and risk assessment. *Int J Cancer* 2004;110:3–14.
- Dankovic D, Kuempel E, Wheeler M. An approach to risk assessment for TiO<sub>2</sub>. *Inhal Toxicol* 2007;19 Suppl 1: 205–12.
- Boffetta P, Soutar A, Cherrie JW, et al. Mortality among workers employed in the titanium dioxide production industry in Europe. *Cancer Causes Control* 2004;15:697–706.
- Fryzek JP, Chadda B, Marano D, et al. A cohort mortality study among titanium dioxide manufacturing workers in the United States. *J Occup Environ Med* 2003;45:400–9.
- Park S, Lee YK, Jung M, et al. Cellular toxicity of various inhalable metal nanoparticles on human alveolar epithelial cells. *Inhal Toxicol* 2007;19:59–65.
- Chen M, von Mikecz A. Formation of nucleoplasmic protein aggregates impairs nuclear function in response to SiO<sub>2</sub> nanoparticles. *Exp Cell Res* 2005;305:51–62.
- Jani PU, McCarthy DE, Florence AT. Titanium dioxide (rutile) particle uptake from the rat GI tract and translocation to systemic organs after oral administration. *Int J Pharm* 1994;105:157–68.
- Driscoll KE, Deyo LC, Carter JM, et al. Effects of particle exposure and particle-elicited inflammatory cells on mutation in rat alveolar epithelial cells. *Carcinogenesis* 1997;18:423–30.
- Wang JJ, Sanderson BJ, Wang H. Cyto- and genotoxicity of ultrafine TiO<sub>2</sub> particles in cultured human lymphoblastoid cells. *Mutat Res* 2007;628:99–106.
- Warheit DB, Hoke RA, Finlay C, et al. Development of a base set of toxicity tests using ultrafine TiO<sub>2</sub> particles as a component of nanoparticle risk management. *Toxicol Lett* 2007;171:99–110.
- Gurr JR, Wang AS, Chen CH, Jan KY. Ultrafine titanium dioxide particles in the absence of photoactivation can induce oxidative damage to human bronchial epithelial cells. *Toxicology* 2005;213:66–73.
- Kang SJ, Kim BM, Lee YJ, Chung HW. Titanium dioxide nanoparticles trigger p53-mediated damage response in peripheral blood lymphocytes. *Environ Mol Mutagen* 2008;49:399–405.
- Linnainmaa K, Kivipensas P, Vainio H. Toxicity and cytogenetic studies of ultrafine titanium dioxide in cultured rat liver epithelial cells. *Toxicol In Vitro* 1997;11:329–35.
- Rahman Q, Lohani M, Dopp E, et al. Evidence that ultrafine titanium dioxide induces micronuclei and apoptosis in Syrian hamster embryo fibroblasts. *Environ Health Perspect* 2002;110:797–800.
- Li S, Zhu H, Zhu R, et al. Impact and mechanism of TiO<sub>2</sub> nanoparticles on DNA synthesis *in vitro*. *Sci China Ser B Chem* 2008;51:367–72.
- Zhu R-R, Wang S-L, Zhang R, Sun X-Y, Yao S-D. A novel toxicological evaluation of TiO<sub>2</sub> nanoparticles on DNA structure. *Chin J Chem* 2007;25:958–61.
- Chen HW, Su SF, Chien CT, et al. Titanium dioxide nanoparticles induce emphysema-like lung injury in mice. *FASEB J* 2006;20:2393–5.
- de Haar C, Hassing I, Bol M, Bleumink R, Pieters R. Ultrafine but not fine particulate matter causes airway

- inflammation and allergic airway sensitization to co-administered antigen in mice. *Clin Exp Allergy* 2006; 36:1469–79.
20. Grassian VH, O'Shaughnessy PT, Adamcakova-Dodd A, Pettibone JM, Thorne PS. Inhalation exposure study of titanium dioxide nanoparticles with a primary particle size of 2 to 5 nm. *Environ Health Perspect* 2007;115: 397–402.
21. Oberdorster G, Ferin J, Lehnert BE. Correlation between particle size, *in vivo* particle persistence, and lung injury. *Environ Health Perspect* 1994;102 Suppl 5:173–9.
22. Federici G, Shaw BJ, Handy RD. Toxicity of titanium dioxide nanoparticles to rainbow trout (*Oncorhynchus mykiss*): Gill injury, oxidative stress, and other physiological effects. *Aquat Toxicol* 2007;84:415–30.
23. Veranth JM, Kaser EG, Veranth MM, Koch M, Yost GS. Cytokine responses of human lung cells (BEAS-2B) treated with micron-sized and nanoparticles of metal oxides compared to soil dusts. *Part Fibre Toxicol* 2007;4:2.
24. Reliene R, Bishop AJ, Aubrecht J, Schiestl RH. *In vivo* DNA deletion assay to detect environmental and genetic predisposition to cancer. *Methods Mol Biol* 2004;262: 125–39.
25. Dick CA, Brown DM, Donaldson K, Stone V. The role of free radicals in the toxic and inflammatory effects of four different ultrafine particle types. *Inhal Toxicol* 2003;15:39–52.
26. Gallagher J, Heinrich U, George M, et al. Formation of DNA adducts in rat lung following chronic inhalation of diesel emissions, carbon black and titanium dioxide particles. *Carcinogenesis* 1994;15:1291–9.
27. Rehn B, Seiler F, Rehn S, Bruch J, Maier M. Investigations on the inflammatory and genotoxic lung effects of two types of titanium dioxide: untreated and surface treated. *Toxicol Appl Pharmacol* 2003;189:84–95.
28. Xia T, Kovochich M, Brant J, et al. Comparison of the abilities of ambient and manufactured nanoparticles to induce cellular toxicity according to an oxidative stress paradigm. *Nano Lett* 2006;6:1794–807.
29. Reliene R, Schiestl RH. Mouse models for induced genetic instability at endogenous loci. *Oncogene* 2003;22: 7000–10.
30. Reliene R, Pollarda J, Sobol Z, et al. *N*-acetyl cysteine protects against ionizing radiation-induced DNA damage but not against cell killing in yeast and mammals. *Mutat Res* 2009;665:37–43.
31. Westbrook AM, Wei B, Braun JB, Schiestl RH. Intestinal mucosal inflammation leads to systemic genotoxicity in mice. *Cancer Res* 2009;69:4827–34.
32. Reliene R, Fischer E, Schiestl RH. Effect of *N*-acetyl cysteine on oxidative DNA damage and the frequency of DNA deletions in atm-deficient mice. *Cancer Res* 2004;64:5148–53.
33. Ismail IH, Wadhwa TI, Hammarsten O. An optimized method for detecting  $\gamma$ -H2AX in blood cells reveals a significant interindividual variation in the  $\gamma$ -H2AX response among humans. *Nucleic Acids Res* 2007;35:e36.
34. Tao F, Kobzik L. Lung macrophage-epithelial cell interactions amplify particle-mediated cytokine release. *Am J Respir Cell Mol Biol* 2002;26:499–505.
35. Thevenot P, Cho J, Wavhal D, Timmons RB, Tang L. Surface chemistry influences cancer killing effect of TiO<sub>2</sub> nanoparticles. *Nanomed Nanotechnol Biol Med* 2008;4:226–36.
36. Garabrant DH, Fine LJ, Oliver C, Bernstein L, Peters JM. Abnormalities of pulmonary function and pleural disease among titanium metal production workers. *Scand J Work Environ Health* 1987;13:47–51.
37. National Institute of Occupational Safety and Health. Evaluation of health hazard and recommendations for occupational exposure to titanium dioxide. NIOSH Current Intelligence Bulletin. Cincinnati: National Institute of Occupational Safety and Health, DHHS publication; 2005.
38. Schins RP, Knaapen AM. Genotoxicity of poorly soluble particles. *Inhal Toxicol* 2007;19 Suppl 1:189–98.
39. Donaldson K, Beswick PH, Gilmour PS. Free radical activity associated with the surface of particles: a unifying factor in determining biological activity? *Toxicol Lett* 1996;88:293–8.
40. Gilmour P, Brown DM, Beswick PH, et al. Surface free radical activity of PM10 and ultrafine titanium dioxide: a unifying factor in their toxicity? *Ann Occup Hyg* 1997; 41:32–8.
41. Zhang Q, Kusaka Y, Sato K, et al. Differences in the extent of inflammation caused by intratracheal exposure to three ultrafine metals: role of free radicals. *J Toxicol Environ Health Part A* 1998;53:423–38.
42. Helfenstein M, Miragoli M, Rohr S, et al. Effects of combustion-derived ultrafine particles and manufactured nanoparticles on heart cells *in vitro*. *Toxicology* 2008;253:70–8.
43. Sayes CM, Wahi R, Kurian PA, et al. Correlating nanoscale titania structure with toxicity: a cytotoxicity and inflammatory response study with human dermal fibroblasts and human lung epithelial cells. *Toxicol Sci* 2006;92:174–85.
44. Reliene R, Schiestl RH. Antioxidant *N*-acetyl cysteine reduces incidence and multiplicity of lymphoma in Atm deficient mice. *DNA Repair (Amst)* 2006;5: 852–9.
45. Brennan RJ, Swoboda BE, Schiestl RH. Oxidative mutagens induce intrachromosomal recombination in yeast. *Mutat Res* 1994;308:159–67.
46. Schins RPF. Mechanisms of genotoxicity of particles and fibers. *Inhal Toxicol* 2002;14:57–78.