

Negative Regulation by Interleukin-3 (IL-3) of Mouse Early B-Cell Progenitors and Stem Cells in Culture: Transduction of the Negative Signals by β c and β IL-3 Proteins of IL-3 Receptor and Absence of Negative Regulation by Granulocyte-Macrophage Colony-Stimulating Factor

By Takuya Matsunaga, Fumiya Hirayama, Yuji Yonemura, Richard Murray, and Makio Ogawa

The receptors for interleukin-3 (IL-3), granulocyte-macrophage colony-stimulating factor (GM-CSF), and IL-5 share a common signaling subunit β c. However, in the mouse, there is an additional IL-3 signaling protein, β IL-3, which is specific for IL-3. We have previously reported that IL-3 abrogates the lymphoid potentials of murine lymphohematopoietic progenitors and the reconstituting ability of hematopoietic stem cells. We used bone marrow cells from β c- and β IL-3-knock-out mice to examine the relative contributions of the receptor proteins to the negative regulation by IL-3. First, we tested the effects of IL-3 on lymphohematopoietic progenitors by using lineage-negative (Lin⁻) marrow cells of 5-fluorouracil (5-FU)-treated mice in the two-step methylcellulose culture we reported previously. Addition of IL-3 to the combination of steel factor (SF, c-kit ligand) and IL-11 abrogated the B-lymphoid potential of the marrow cells of both types of knock-out mice as well as wild-type mice. Next, we

investigated the effects of IL-3 on in vitro expansion of the hematopoietic stem cells. We cultured Lin⁻Sca-1-positive, c-kit-positive marrow cells from 5-FU-treated mice in suspension in the presence of SF and IL-11 with or without IL-3 for 7 days and tested the reconstituting ability of the cultured cells by transplanting the cells into lethally irradiated Ly-5 congenic mice together with "compromised" marrow cells. Presence of IL-3 in culture abrogated the reconstituting ability of the cells from both types of knock-out mice and the wild-type mice. In contrast, addition of GM-CSF to the suspension culture abrogated neither B-cell potential nor reconstituting abilities of the cultured cells of wild-type mice. These observations may have implications in the choice of cytokines for use in in vitro expansion of human hematopoietic stem cells and progenitors.

© 1998 by The American Society of Hematology.

INTERLEUKIN-3 (IL-3) supports the development of multiple hematopoietic lineages by interacting with multipotential and lineage-committed progenitors in culture.¹⁻³ Studies in our laboratory indicated that IL-3, as a single factor supports proliferation of the progenitors after they exit from the cell-cycle dormant state (G0).^{3,4} IL-3 also synergizes with IL-6,⁵ IL-11,^{6,7} granulocyte colony-stimulating factor (G-CSF),⁸ leukemia inhibitory factor,⁹ thrombopoietin (TPO),^{10,11} and steel factor (SF, c-kit ligand)^{12,13} in triggering cell divisions of the multipotential progenitors in G0.

In contrast to the positive regulation of myeloid lineages, IL-3 seems to exert negative effects on the early stages of lymphopoiesis. In our laboratory, we have established a two-step methylcellulose culture assay for murine lymphohematopoietic progenitors and characterized their cytokine requirement.¹⁴ SF-based cytokine combinations supported the proliferation and differentiation of lymphohematopoietic progenitors whereas addition of IL-3 to the permissive cytokine combinations abrogated the B-lymphoid potential of the progenitors.¹⁵ We subsequently observed that T-cell¹⁶ and natural killer-cell¹⁷ potential of the progenitors is also inhibited by IL-3. These observations raised the possibility that IL-3 may be a stage-specific negative regulator and that it may suppress the earliest process of hematopoiesis, ie, self-renewal of the stem cells. This hypothesis was confirmed later by our observation that IL-3 abrogates reconstituting ability of hematopoietic stem cells with long-term engraftment capability.¹⁸

Both mouse and human IL-3 receptors are heterodimers consisting of α and β subunits.¹⁹⁻²² The high-affinity receptors for human IL-3, granulocyte-macrophage CSF (GM-CSF), and IL-5 share common β subunit (β c).¹⁹⁻²² The α subunits are specific for each cytokine and bind their ligand with low affinity. Whereas there is only one type of human β subunit, β c, mouse has two closely related β subunits, β c and β IL-3.²³⁻²⁵ They have 56% homology with human β c. Like human β c, mouse β c is the common β subunit of the receptors for mouse IL-3, GM-CSF, and IL-5. Although β IL-3 has extensive se-

quence homology with mouse β c (91% at the amino acid level), β IL-3 does not form a high-affinity receptor with mouse IL-5 or mouse GM-CSF α receptors. When transfected into mouse T-cell line, CTLL-2, both β c and β IL-3 interacted equally well with the α subunit of mouse IL-3 receptor and transmitted proliferation signals in the presence of IL-3.²⁵ Independently, investigators in two laboratories²⁶⁻²⁸ reported generation and analysis of the mice lacking either of the two β receptor genes. Mice lacking the β c showed pulmonary alveolar proteinosis-like disease, a phenotype similar to that of GM-CSF-deficient mice.^{29,30} The number of peripheral blood eosinophils of β c-deficient mice was markedly reduced, and the mice failed to produce eosinophils in response to parasitic infections because of absence of transduction of IL-5 signals.²⁶ The colony-forming ability of the bone marrow cells from β c- or β IL-3-deficient mice was also examined.²⁶⁻²⁸ Cells from β c-deficient mice did not respond to GM-CSF or IL-5 but responded normally to IL-3. In contrast, cells from β IL-3-deficient mice responded normally to IL-3, GM-CSF, or IL-5. These results clearly show the unique redundancy of mouse IL-3 receptor

From the Department of Veterans Affairs Medical Center and the Department of Medicine, Medical University of South Carolina, Charleston, SC; and DNAX Research Institute of Molecular and Cellular Biology, Palo Alto, CA.

Submitted November 24, 1997; accepted April 1, 1998.

Supported by the Office of Research and Development, Medical Research Service, Department of Veterans Affairs; NIH grants DK32294 and DK/HL48714; and a contribution from Amgen.

Address reprint requests to Makio Ogawa, MD, PhD, Ralph H. Johnson Medical Center, 109 Bee St, Charleston, SC 29401-5799.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. section 1734 solely to indicate this fact.

© 1998 by The American Society of Hematology.

0006-4971/98/9203-0028\$3.00/0

system, which is not present for the human IL-3 receptor. In this study, we used cells from β c and β IL-3-gene knock-out mice to examine the relative contributions of β c and β IL-3 to the negative regulation of the early B lymphopoiesis and the long-term reconstituting ability of stem cells.

MATERIALS AND METHODS

Cytokines. Purified recombinant murine IL-3 and GM-CSF were purchased from R&D Systems (Minneapolis, MN). Purified recombinant murine SF was obtained from Immunex (Seattle, WA). Purified recombinant human IL-6 was a gift from M. Naruto of Toray Industries (Kamakura, Japan). Purified recombinant human IL-7 was a gift from Sterling Winthrop Inc (Collegeville, PA). Purified recombinant human IL-11 was a gift from P. Schendel, Genetics Institute (Cambridge, MA). Purified recombinant human TPO was prepared by the Cytokine Production Group of Kirin Brewery (Takasaki, Japan). Recombinant human FLT3/FLK-2 ligand (FL) was provided by S.D. Lyman of Immunex. Purified recombinant human erythropoietin (EPO) was provided by the Genetics Institute Clinical Manufacturing Group (Cambridge, MA). Recombinant human G-CSF was a gift from A. Shimosaka of Kirin Brewery, Co, Ltd. Unless otherwise specified, the concentrations of cytokines used were as follows: IL-3, 10 ng/mL; SF, 100 ng/mL; IL-6, 100 ng/mL; IL-7, 200 U/mL; IL-11, 100 ng/mL; TPO, 100 ng/mL; FL, 100 ng/mL; EPO, 2 U/mL; G-CSF, 100 ng/mL; GM-CSF, 160 ng/mL.

Monoclonal antibodies (MoAbs). Hybridoma D7 (anti-Ly-6A/E [anti-Sca-1]); rat immunoglobulin G [IgG]2a) was a gift from P. Kincaid of Oklahoma Medical Research Foundation (Oklahoma City, OK). MoAb ACK4 (anti-c-kit; rat IgG2a) was provided by S.I. Nishikawa of Kyoto University (Kyoto, Japan). Hybridoma RB6-8C5 (anti-mouse granulocytes; rat IgG2b) was provided by R.L. Coffman of DNAX (Palo Alto, CA). MoAb TER119 (anti-erythrocytes; rat IgG2b) was a gift from T. Kina of Kyoto University. Hybridomas 14.8 (anti-B220; rat IgG2b), M1/70.15.11.5 (anti-macrophages; rat IgG2b), GK1.5 (anti-CD4; rat IgG2b), and 53-6.72 (anti-CD8; rat IgG2a) were purchased from American Type Culture Collection (Rockville, MD). 53-2.1 (Biotin-conjugated-anti-Thy-1.2; rat IgG2a), RA3-6B2 (Biotin-conjugated-anti-CD45R/B220; rat IgG2a), RB6-8C5 (Biotin-conjugated-anti-Gr-1; rat IgG2b), and M1/70 (Biotin-conjugated-anti-Mac-1; rat IgG2b) were purchased from Pharmingen (San Diego, CA). AL1-4A2 (fluorescein isothiocyanate [FITC]-conjugated anti-Ly-5.2; mouse IgG1) and A20-1.7 (FITC-conjugated anti-Ly-5.1; mouse IgG1) were provided by H. Fleming of Emory University.

Cell preparations. Cells from 10- to 15-week-old male and female β c-deficient,²⁶ β IL-3-deficient,²⁶ and wild-type littermates (C57B1/6)²⁶ were used in clonal cultures and transplantation experiments. 5-fluorouracil (5-FU; Adria Laboratories, Columbus, OH) was administered intravenously through the tail vein at 150 mg/kg body weight, and bone marrow cells were obtained 2 days later. Cells prepared from pooled femurs and tibiae were washed twice and then subjected to density gradient separation by using Nycodenz (Accurate Chemical and Scientific Corp, Westbury, NY) solution. Cells with densities ranging from 1.063 g/mL to 1.077 g/mL were collected.³¹ Cells reacting to a cocktail of lineage-specific rat MoAbs (RB6-8C5, 14.8, M1/70.15.11.5, GK1.5, TER119, and 53-6.72) were removed twice by using immunomagnetic beads (Dynabeads M-450 coupled to sheep anti-rat IgG; DYNAL, Great Neck, NY). The resulting lineage-negative cells (Lin^-) were treated with normal rat IgG (Jackson ImmunoResearch Laboratories, West Grove, PA) at 20 μ g/ 10^6 cells to prevent nonspecific binding of MoAbs to Fc receptors, and were then stained with FITC-conjugated rat MoAb D7 (anti-Sca-1)³² and biotin-conjugated rat MoAb ACK4 (anti-c-kit).³³ Cells were washed twice before staining with streptavidin-conjugated R-phycoerythrin (PE) (Jackson ImmunoResearch Laboratories). Both FITC-conjugated rat IgG2a and biotin-labeled rat IgG2a (Caltag

Laboratories, San Francisco, CA) were used as isotype controls. Sca-1⁺ c-kit⁺ cells were collected by sorting on FACS Vantage (Becton Dickinson Immunocytometry Systems).

Two-step methylcellulose culture for lymphohematopoietic progenitors. Four thousand Lin^- cells or 100 Lin^- Sca-1⁺ c-kit⁺ cells of 5-FU-treated mice were plated in 35-mm suspension culture dishes (Falcon, Lincoln Park, NJ) containing α -medium (ICN, Irvine, CA), 1.2% 1,500-cp methylcellulose (Shinetsu Chemical, Tokyo, Japan), 30% (vol/vol) fetal calf serum (FCS) (Intergen, Purchase, NY), 1% deionized Fraction V bovine serum albumin (BSA) (Sigma Chemical, St Louis, MO), 1×10^{-4} mol/L 2-mercaptoethanol (2-ME) (Sigma), and designated cytokines. Dishes were incubated at 37°C in a humidified atmosphere flushed with 5% CO₂. On the designated day of incubation, 20 colonies were picked, pooled, and washed and 1/20 of the pooled cells were plated in secondary methylcellulose culture containing SF and IL-7. The number of pre-B-cell colonies was counted on day 10 of the secondary culture by using criteria previously described.¹⁴

Suspension and clonal cell cultures. Two hundred Lin^- Sca-1⁺ c-kit⁺ cells were incubated in each well of a 6-well plate (Falcon) in 5 mL suspension culture. The culture medium contained α -medium, 20% (vol/vol) FCS, 1% deionized BSA, 1×10^{-4} mol/L 2-ME, and designated cytokines. On day 7 of incubation, aliquots were analyzed for colony formation and in vivo reconstituting capabilities. Clonal culture was performed in 35-mm suspension culture dishes containing α -medium; 1.2% 1,500-cp methylcellulose; 30% FCS; 1% BSA; and 1×10^{-4} mol/L 2-ME, SF, IL-3, IL-6, FL, TPO, and EPO. Colonies were scored on day 8 of incubation by in situ observation of the plates on an inverted microscope according to the criteria described previously.³⁴ Megakaryocyte colonies were scored when the colony contained four or more megakaryocytes. Abbreviations for colony types are as follows: GM, granulocyte/macrophage colonies; GEM, granulocyte/erythrocyte/macrophage colonies; GMM, granulocyte/macrophage/megakaryocyte colonies; GEMM, granulocyte/erythrocyte/macrophage/megakaryocyte colonies³⁴; Meg, megakaryocyte colonies.

In vivo reconstitution experiments. In studies of knock-out mice, 10- to 12-week-old male C57B1/6-Ly-5.1 mice were administered with single 850-cGy total-body irradiation via a 4×10^6 V linear accelerator. After irradiation of the recipient mice, freshly sorted Lin^- Sca-1⁺ c-kit⁺ marrow cells (Ly-5.2 cells) from female wild-type, β c^{-/-}, and β IL-3^{-/-} mice were injected into the tail vein of the recipients together with 4×10^5 "compromised" marrow cells of male C57B1/6-Ly-5.1 mice. "Compromised" cells had been subjected to two previous rounds of transplantation and regeneration in male mice.³⁵ Cells cultured in suspension were also tested for reconstituting capabilities

Table 1. Effects of IL-3 on Lymphohematopoietic Progenitors

Mice	Cytokines	Replated on Day 10		Replated on Day 12
		No. of 1 ⁺ Colonies	No. of Pre-B-Cell Colonies	No. of Pre-B-Cell Colonies
Wild-type	SF, IL-11	34 ± 5	33 ± 4	15 ± 2
(β c ^{+/+} , β IL-3 ^{+/+})	SF, IL-11, IL-3	32 ± 6	0	0
β c ^{-/-}	SF, IL-11	36 ± 2	34 ± 2	14 ± 2
	SF, IL-11, IL-3	35 ± 2	0	0
β IL-3 ^{-/-}	SF, IL-11	35 ± 5	32 ± 2	16 ± 1
	SF, IL-11, IL-3	36 ± 4	0	0

Four thousand Lin^- cells from the marrow of 5-FU-treated mice were cultured in methylcellulose with SF and IL-11 in the presence or absence of IL-3. Ten and 12 days later, 20 primary colonies were pooled and 5% of the cells were replated in methylcellulose with SF and IL-7. The pre-B-cell colony formation was examined on day 10 of secondary culture. Similar results were obtained in an additional experiment.

Table 2. Expansion of Total Cells and Progenitors in Suspension Culture

Mice	Cytokines	Cells	Total CFU	CFU-GEMM	CFU-Meg
Wild-type	No culture*	200	67	16	8
(β c +/+)	SF, IL-11	1.1×10^5	2.7×10^4	687	137
β IL-3 +/+)	SF, IL-11, IL-3	3.2×10^6	4.0×10^4	2.0×10^3	400
β c -/-	No culture*	200	68	13	13
	SF, IL-11	6.7×10^4	1.3×10^4	335	84
	SF, IL-11, IL-3	2.5×10^6	4.9×10^4	3.3×10^3	650
β IL-3 -/-	No culture*	200	64	13	8
	SF, IL-11	1.2×10^5	1.1×10^4	450	150
	SF, IL-11, IL-3	2.6×10^6	2.9×10^4	2.0×10^3	650

200 Lin⁻ Sca-1⁺ c-kit⁺ cells prepared from the marrow of 5-FU-treated mice were cultured in the presence of designated cytokines. On day 7 of culture, cells were analyzed for CFU, CFU-GEMM, and CFU-Meg in methylcellulose culture. Similar results were obtained in an additional experiment.

*Freshly sorted Lin⁻ Sca-1⁺ c-kit⁺ cells.

after 7 days' incubation with designated cytokines; all of the cells in each well were injected into male C57B1/6-Ly-5.1 mice together with "compromised" cells. Peripheral blood was obtained from the retro-orbital venous plexus using heparin-coated micropipettes (Drummond Scientific Co, Broomall, PA) 2, 4, and 6 months after transplantation. Red blood cells were lysed by 0.15 mol/L NH₄Cl. The samples were then used for flow cytometric analysis of donor-derived cells by staining with FITC-conjugated anti-Ly-5.2 (AL1-4A2). In studies of GM-CSF effects, we used C57B1/6-Ly-5.1 male mice as donors and C57B1/6-Ly-5.2 female mice as recipients. The lineage phenotype of the donor cells at 6 months posttransplantation was determined by staining with biotin-conjugated anti-Thy-1.2, biotin-conjugated anti-CD45R/B220, biotin-conjugated anti-Gr-1, and biotin-conjugated anti-Mac-1. For indirect staining of cells with biotin-conjugated antibodies, cells were first incubated with biotin-conjugated antibodies, then followed by staining with streptavidin-conjugated PE.

RESULTS


Effects of IL-3 on lymphohematopoietic progenitors. The results of the studies of lymphohematopoietic progenitors are presented in Table 1. Regardless of the origin of the cells,

colonies supported by the combination of SF and IL-11 possessed B-cell potential. Addition of IL-3 to the combination of SF and IL-11 strongly inhibited the B-cell potential of the primary colonies of both types of knock-out mice as well as wild-type mice. As we reported previously,^{14,15} the number of primary colonies was unaffected by IL-3. These results indicated that β c and β IL-3 are redundant regarding signal transduction in negative regulation of early B lymphopoiesis by IL-3.

Effects of IL-3 on expansion of total cells and progenitors. Next we studied the effects of IL-3 on the expansion of cells and colony-forming cells by plating 200 enriched cells in 7-day suspension culture in the presence of 100 ng/mL SF and 100 ng/mL IL-11 with or without 100 ng/mL IL-3. As shown in Table 2, the combination of SF and IL-11 increased the total cell counts by 335- to 600-fold, total colony-forming units (total CFU) by 172- to 402-fold, CFU-GEMM by 26- to 43-fold, and CFU-Meg by 6- to 19-fold, as compared with freshly enriched cells. Addition of IL-3 resulted in about 30-fold enhancement of total cell counts and several-fold increase in the total CFU, CFU-GEMM, and CFU-Meg. These results confirmed that the myelostimulatory effects of IL-3 are transduced by both β c and β IL-3.

Effects of IL-3 on long-term repopulating cells (LTRC). We then tested the in vivo reconstituting ability of the cultured cells. The suspension cultures were initiated with 200 Lin⁻ Sca-1⁺ c-kit⁺ cells in the presence of 100 ng/mL SF and 100 ng/mL IL-11 with or without 100 ng/mL IL-3. After 7 days of incubation, cells in a well were obtained and injected into a lethally irradiated Ly-5.1 recipient. As a control group, we also transplanted 200 freshly prepared Lin⁻ Sca-1⁺ c-kit⁺ cells from wild-type, β IL-3 -/-, and β c -/- mice. The results of the analyses of peripheral blood nucleated cells are presented in Fig 1. The enriched cells from all mice incubated with SF and IL-11 had similar reconstituting ability as freshly prepared cells. In contrast, incubation in the presence of IL-3 significantly reduced the reconstituting abilities of the cells of all types of mice. A few repopulating cells survived in SF, IL-11, and IL-3 at 2 months posttransplantation, as evinced by the appearance of small percentages of donor cells in five, three, and three mice

Fig 1. Repopulating abilities of freshly sorted bone marrow cells and cultured cells. (A) Mice transplanted with wild-type cells. (B) Mice transplanted with β IL-3 -/- cells. (C) Mice transplanted with β c -/- cells. (○), 2 months posttransplantation; (●), 4 months posttransplantation; (◻), 6 months posttransplantation.



out of a total of seven in Fig 1A, B, and C, respectively. However, reconstitution was absent at 4 and 6 months posttransplantation. These observations confirmed that the negative effects of IL-3 are directed only to LTRC and that the effects are transduced by both βc and $\beta IL-3$.

At 6 months posttransplantation, the proportions of donor blood nucleated cells in each of T-cell, B-cell, and myeloid (granulocyte and monocyte/macrophage) compartments were determined (Table 3). Multi-lineage cells were detected in the peripheral blood of engrafted recipients. An example of analysis

Table 3. Lineage Expression by Engrafted Donor (Ly-5.2) Cells in Individual Mice at 6 Months Posttransplantation

Mice	Cytokines	Mice	% B220 ⁺ Cells	% Thy-1.2 ⁺ Cells	% Gr-1 ⁺ Mac-1 ⁺ Cells
Wild-type (βc +/+, $\beta IL-3$ +/+)	No culture	1	35.4	16.9	22.9
		2	43.8	12.1	20.9
		3	19.3	10.8	11.6
		4	29.2	13.2	20.5
		5	16.6	14.3	15.2
		6	36.2	18.0	16.8
		7	33.9	22.0	24.6
	SF, IL-11	1	11.4	4.8	2.4
		2	21.3	15.5	12.4
		3	38.0	22.2	14.5
		4	22.6	12.5	4.9
		5	39.4	23.7	23.6
		6	34.1	19.7	19.3
		7	22.4	16.6	14.6
$\beta IL-3$ -/-	No culture	1	27.6	17.7	15.4
		2	34.2	18.1	15.9
		3	39.1	20.8	18.9
		4	11.9	6.5	8.7
		5	15.7	8.1	3.1
		6	14.8	10.9	3.9
		7	30.9	26.1	18.1
	SF, IL-11	1	20.7	16.1	10.0
		2	8.6	3.9	2.7
		3	24.6	17.4	10.5
		4	13.8	13.5	11.3
		5	24.3	22.3	7.4
		6	26.1	19.6	16.7
		7	14.9	4.3	3.2
βc -/-	No culture	1	10.6	5.1	2.9
		2	46.1	22.9	15.8
		3	36.3	12.0	6.6
		4	26.0	11.2	6.6
		5	17.3	6.4	4.2
		6	23.5	13.7	23.2
		7	25.7	13.0	11.9
	SF, IL-11	1	30.0	23.7	11.6
		2	24.7	11.7	4.9
		3	19.5	9.7	6.8
		4	25.5	10.0	9.4
		5	18.5	16.4	12.0
		6	10.7	7.6	5.8
		7	16.7	11.6	2.9

Irradiated recipients (seven mice per group) were transplanted with freshly sorted 200 Lin⁻ Sca-1⁺ c-kit⁺ cells or with all of the cultures expanded from 200 Lin⁻ Sca-1⁺ c-kit⁺ cells. Background (peripheral blood of nontransplanted male mice) staining by anti-Ly-5.2 MoAb (AL1-4A2) was less than 1.0%.

of a mouse transplanted with βc -/- cells cultured with SF and IL-11 is shown in Fig 2.

Effects of GM-CSF. The observation that the βc chain transduces the negative effects of IL-3 on the stem cells and lymphohematopoietic progenitors raised the possibility that βc may also transduce negative signals of GM-CSF. Earlier we reported that, in addition to IL-11, IL-6 and G-CSF can interact with SF in stimulating proliferation of cell cycle dormant progenitors^{3,12,13} and the lymphohematopoietic progenitors.^{14,15} In the next two experiments, we tested the effects of GM-CSF and IL-3 in variable concentrations on the progenitors and stem cells using cells from wild-type mice. Specifically, we tested the effects of addition of these cytokines to the cultures supported by SF and one of IL-11, IL-6, and G-CSF. The results are presented in Table 4. As reported previously,^{14,15} the primary colonies supported by SF plus IL-11, SF plus IL-6, and SF plus G-CSF possessed B-lymphoid potential. Addition of IL-3 at as low as 1 ng/mL concentration completely abrogated the B-cell potential of the primary colonies. In contrast, addition of GM-CSF in concentrations ranging from 10 to 1,000 ng/mL failed to suppress the lymphohematopoietic progenitors.

We then tested the effects of GM-CSF on repopulating abilities of the bone marrow cells. The enriched marrow cells of Ly-5.1 wild-type mice were cultured in suspension for 1 week under permissive cytokine conditions with or without additional GM-CSF and transplanted to lethally irradiated Ly-5.2 recipients. As presented in Table 5, GM-CSF did not negatively affect the repopulating abilities of cultured cells.

DISCUSSION

There is significant current interest in hematology/oncology fields regarding in vitro expansion of hematopoietic stem cells and progenitors.³⁶⁻⁵¹ A number of investigators have already shown that it is possible to increase the number of hematopoietic progenitors in culture by using combinations of early-acting cytokines.³⁶⁻⁵¹ Because of the well-known myelopoietic effects of IL-3, the majority of preclinical protocols for murine³⁷⁻⁴³ and human^{36,44-51} cells included IL-3. We previously noted negative effects of IL-3 on the ability of cultured cells to engraft the marrow of recipient mice.¹⁸ Our observation was in agreement with the report from Peters et al⁵² that suspension culture of murine marrow cells in the presence of IL-3, IL-6, IL-11, and SF results in impairment of the engrafting capability of the cultured cells.

The negative effects of IL-3 observed in murine models may be relevant to in vitro manipulation of human stem cells. Ten patients with advanced cancers were transplanted with peripheral blood progenitors that had been expanded in cultures containing IL-3.⁵³ Recently, Williams et al⁵⁴ reported transplantation of peripheral blood CD34⁺ cells expanded in liquid culture with PIXY321 (the fusion product of IL-3 and GM-CSF) to 8 patients with advanced breast cancer. No information was provided in these reports regarding long-term effects on the recipient's hematopoiesis. Donahue et al⁵⁵ studied transduction of glucocerebrosidase genes to primate CD34⁺ Thy-1⁺ cells using 7-day culture in the presence of IL-3, IL-6, and SF.⁵⁵ After autologous transplantation, provirus was detected at all time points in both B-cell and T-cell lineages, but long-term gene transfer was not observed in the granulocyte population.⁵⁵ It is




Fig 2. An example of hematopoietic reconstitution by $\beta c^{-/-}$ cells cultured with SF and IL-11. Nucleated blood cells of a recipient mouse were analyzed using flow cytometry 6 months after transplantation. Thy-1.2⁺ cells, B220⁺ cells, and Gr-1⁺ Mac-1⁺ cells of donor (Ly-5.2) origin are seen.

Table 4. Effects of GM-CSF on Lymphohematopoietic Progenitors

Cytokines	Replated on Day 8		Replated on Day 11
	No. of 1 ⁺ Colonies	No. of Pre-B-Cell Colonies	No. of Pre-B-Cell Colonies
	SF, IL-11	27 ± 2	44 ± 2
SF, IL-11, IL-3 (1 ng/mL)	26 ± 3	0	0
SF, IL-11, IL-3 (10 ng/mL)	26 ± 4	0	0
SF, IL-11, IL-3 (100 ng/mL)	25 ± 3	0	0
SF, IL-11, GM-CSF (10 ng/mL)	27 ± 1	45 ± 2	31 ± 5
SF, IL-11, GM-CSF (100 ng/mL)	28 ± 4	44 ± 5	32 ± 8
SF, IL-11, GM-CSF (1,000 ng/mL)	27 ± 4	45 ± 2	29 ± 4
SF, IL-6	25 ± 3	33 ± 4	51 ± 3
SF, IL-6, IL-3 (1 ng/mL)	28 ± 3	0	0
SF, IL-6, IL-3 (10 ng/mL)	27 ± 5	0	0
SF, IL-6, IL-3 (100 ng/mL)	28 ± 3	0	0
SF, IL-6, GM-CSF (10 ng/mL)	27 ± 3	31 ± 2	53 ± 5
SF, IL-6, GM-CSF (100 ng/mL)	28 ± 6	30 ± 3	51 ± 7
SF, IL-6, GM-CSF (1,000 ng/mL)	24 ± 4	31 ± 4	49 ± 7
SF, G-CSF	23 ± 6	55 ± 6	21 ± 3
SF, G-CSF, IL-3 (1 ng/mL)	25 ± 2	0	0
SF, G-CSF, IL-3 (10 ng/mL)	26 ± 2	0	0
SF, G-CSF, IL-3 (100 ng/mL)	25 ± 4	0	0
SF, G-CSF, GM-CSF (10 ng/mL)	23 ± 7	53 ± 9	20 ± 3
SF, G-CSF, GM-CSF (100 ng/mL)	25 ± 2	46 ± 6	21 ± 2
SF, G-CSF, GM-CSF (1,000 ng/mL)	25 ± 5	53 ± 5	21 ± 3

One hundred Lin⁻ Sca-1⁺ c-kit⁺ cells from the marrow of 5-FU-treated wild-type mice were cultured in methylcellulose with designated cytokines. Eight and 11 days later, 20 primary colonies were pooled and 5% of the cells were replated in methylcellulose with SF and IL-7. The pre-B-cell colony formation was examined on day 10 of secondary culture. Similar results were obtained in an additional experiment.

possible that IL-3 abrogated the long-term reconstitution capability of the cultured primate stem cells.

Both βc and $\beta IL-3$ transduced the negative signals of IL-3. βc and $\beta IL-3$ have been shown to have redundant functions. Coexpression of one of the two β proteins and IL-3 receptor α protein in an IL-2-dependent mouse T-cell line, CTLL-2, resulted in identical, high-affinity binding for mouse IL-3 and IL-3-dependent proliferation of the cell line.²⁵ Neither βc -null nor $\beta IL-3$ -null mice showed apparent hematopoietic defects.²⁶ However, careful analysis of the $\beta IL-3$ -null mice revealed hyporesponsiveness of the hematopoietic progenitors to IL-3, indicating probable quantitative differences between βc and $\beta IL-3$.²⁸

Table 5. Repopulating Abilities of Freshly Prepared Bone Marrow Cells and Cultured Cells

Additional Cytokines	Cells	Fractions of Cells Injected	% Donor PB Cells*	
			2 mo	6 mo
No culture	100	1	51.4 ± 17.2	65.2 ± 26.3
		1/10	16.7 ± 26.7	25.7 ± 30.3
None	1.4 × 10 ⁵	1	65.0 ± 3.6	48.5 ± 2.2
		1/10	19.0 ± 8.0	4.9 ± 3.2
GM-CSF	3.5 × 10 ⁵	1	64.3 ± 12.1	50.4 ± 29.0
		1/10	8.8 ± 6.0	5.5 ± 4.4

Irradiated recipients (five mice per group) were transplanted with 100 or 10 freshly prepared cells or with fractions of day-7 cultures that had been initiated with 100 enriched cells. These cells had been expanded in culture in the presence of SF, IL-6, IL-11, EPO, and the designated additional cytokines. Background (peripheral blood of nontransplanted female mice) staining was less than 1.0%.

Abbreviation: PB, peripheral blood.

*Mean ± SD of percent donor-derived nucleated peripheral blood cells tested with anti-Ly-5.1 MoAb (A20-1.7).

Although receptors for IL-3 and GM-CSF share common signal-transducing protein βc , neither lymphohematopoietic progenitors nor long-term reconstituting cells were negatively affected by GM-CSF. These observations are in agreement with the previous reports on the differences between IL-3 and GM-CSF. Earlier, we documented that GM-CSF is significantly weaker than IL-3 in support of murine⁵⁶ and human⁵⁷ blast cell colony formation. Recently, McKinstry et al⁵⁸ reported that GM-CSF receptor is not expressed by mouse Rhodamine-123^{lo} Lin⁻Ly6A/E (Sca-1)⁺ c-kit⁺ cells that are highly enriched for long-term repopulating cells. These observations together with our observations presented in this paper indicate that the α protein of GM-CSF receptor is expressed by neither the stem cells nor the most primitive progenitors. Further, these observations may have a significant implication in "in vitro expansion" of human stem cells and progenitors. It is possible that the use of GM-CSF may be preferable to IL-3 in cytokine mixtures because GM-CSF may not abrogate long-term reconstituting stem cells while it supports expansion of some multipotential progenitors.

ACKNOWLEDGMENT

We thank Dr Haiqun Zeng for assistance in cell sorting; Dr Pamela N. Pharr and Anne G. Leary for assistance in preparation of this manuscript; and the staff of Radiation Oncology Department of the Medical University of South Carolina for assistance in irradiation of mice.

REFERENCES

- Ihle JN, Keller J, Oroszlan S, Henderson LE, Copeland TD, Fitch F, Prystowsky MB, Goldwasser E, Schrader JW, Palaszynski E, Dy M, Lebel B: Biologic properties of homogenous interleukin 3: I. Demonstration of WEHI-3 growth factor activity, mast cell growth factor activity, P cell-stimulating factor activity, colony-stimulating factor activity, and histamine-producing cell-stimulating factor activity. *J Immunol* 131: 282, 1983
- Metcalf D: Hematopoietic regulators: Redundancy or subtlety? *Blood* 82:3515, 1993
- Ogawa M: Differentiation and proliferation of hematopoietic stem cells. *Blood* 81:2844, 1993
- Suda T, Suda J, Ogawa M, Ihle JN: Permissive role of interleukin 3 (IL-3) in proliferation and differentiation of multipotential hemopoietic progenitors in culture. *J Cell Physiol* 124:182, 1985
- Ikebuchi K, Wong GG, Clark SC, Ihle JN, Hirai Y, Ogawa M: Interleukin 6 enhanced of interleukin 3-dependent proliferation of multipotential hemopoietic progenitors. *Proc Natl Acad Sci USA* 84:9035, 1987
- Musashi M, Clark SC, Sudo T, Urdal DL, Ogawa M: Synergistic interactions between interleukin-11 and interleukin-4 in support of proliferation of primitive hematopoietic progenitors of mice. *Blood* 78:1448, 1991
- Musashi M, Yang YC, Paul SR, Clark SC, Sudo T, Ogawa M: Direct and synergistic effects of interleukin 11 on murine hemopoiesis in culture. *Proc Natl Acad Sci USA* 88:765, 1991
- Ikebuchi K, Clark SC, Ihle JN, Souza LM, Ogawa M: Granulocyte colony-stimulating factor enhances interleukin 3-dependent proliferation of multipotential hemopoietic progenitors. *Proc Natl Acad Sci USA* 85:3445, 1998
- Leary AG, Wong GG, Clark SC, Smith AG, Ogawa M: Leukemia inhibitory factor differentiation-inhibiting activity/human interleukin for DA cells augments proliferation of human hematopoietic stem cells. *Blood* 75:1960, 1990
- Ku H, Yonemura Y, Kaushansky K, Ogawa M: Thrombopoietin, the ligand for the Mpl receptor, synergizes with steel factor and other early acting cytokines in supporting proliferation of primitive hematopoietic progenitors of mice. *Blood* 87:4544, 1996
- Kobayashi M, Laver JH, Kato T, Miyazaki H, Ogawa M: Thrombopoietin supports proliferation of human primitive hematopoietic cells in synergy with steel factor and/or interleukin-3. *Blood* 88:429, 1996
- Tsuji K, Zsebo KM, Ogawa M: Enhancement of murine blast cell colony formation in culture by recombinant rat stem cell factor, ligand for c-kit. *Blood* 78:1223, 1991
- Tsuji K, Lyman SD, Sudo T, Clark SC, Ogawa M: Enhancement of murine hematopoiesis by synergistic interactions between steel factor (ligand for c-kit), interleukin-11, and other early acting factors in culture. *Blood* 79:2855, 1992
- Hirayama F, Shih JP, Awguelewitsh A, Warr GW, Clark SC, Ogawa M: Clonal proliferation of murine lymphohematopoietic progenitors in culture. *Proc Natl Acad Sci USA* 89:5907, 1992
- Hirayama F, Clark SC, Ogawa M: Negative regulation of early B lymphopoiesis by interleukin 3 and interleukin 1 α . *Proc Natl Acad Sci USA* 91:469, 1994
- Hirayama F, Ogawa M: Negative regulation of early T lymphopoiesis by interleukin-3 and interleukin-1 α . *Blood* 86:4527, 1995
- Aiba Y, Ogawa M: Development of natural killer (NK) cells, B lymphocytes, macrophages and mast cells from single hematopoietic progenitors in culture of murine fetal liver cells. *Blood* 90:3923, 1997
- Yonemura Y, Ku H, Hirayama F, Souza LM, Ogawa M: Interleukin 3 or interleukin 1 abrogates the reconstituting ability of hematopoietic stem cells. *Proc Natl Acad Sci USA* 93:4040, 1996
- Kitamura T, Sato N, Arai K, Miyajima A: Expression cloning of the human IL-3 receptor cDNA reveals a shared β subunit for the human IL-3 and GM-CSF receptors. *Cell* 66:1165, 1991
- Tavernier J, Devos R, Cornelis S, Tuypens T, Van der Heyden J, Fiers W, Plaetinck G: A human high affinity interleukin-5 receptor (IL5R) is composed of an IL-5-specific α chain and a β chain shared with the receptor for GM-CSF. *Cell* 66:1175, 1991
- Hayashida K, Kitamura T, Gorman DM, Arai K, Yokota T, Miyajima A: Molecular cloning of a second subunit of the receptor for human granulocyte-macrophage colony-stimulating factor (GM-CSF): Reconstitution of a high-affinity GM-CSF receptor. *Proc Natl Acad Sci USA* 87:9655, 1990
- Kitamura T, Hayashida K, Sakamaki K, Yokota T, Arai K, Miyajima A: Reconstitution of functional receptors for human granulocyte/macrophage colony-stimulating factor (GM-CSF): Evidence that AIC2B cDNA is a subunit of the murine GM-CSF receptor. *Proc Natl Acad Sci USA* 88:5082, 1991
- Itoh N, Yonehara S, Schreurs J, Gorman DM, Maruyama K, Ishii A, Yahara I, Arai K, Miyajima A: Cloning of an interleukin-3 receptor: A member of a distinct receptor gene family. *Science* 247:324, 1990
- Gorman DM, Itoh N, Kitamura T, Schreurs J, Yonehara S, Yahara I, Arai K, Miyajima A: Cloning and expression of a gene encoding an interleukin 3 receptor-like protein: Identification of another member of the cytokine receptor gene family. *Proc Natl Acad Sci USA* 87:5459, 1990
- Hara T, Miyajima A: Two distinct functional high affinity receptors for mouse interleukin-3 (IL-3). *EMBO J* 11:1875, 1992
- Nishinakamura R, Nakayama N, Hirabayashi Y, Inoue T, Aud D, McNeil T, Azuma S, Yoshida S, Toyoda Y, Arai K, Miyajima A, Murray R: Mice deficient for the IL-3/GM-CSF/IL-5 βc receptor exhibit lung pathology and impaired immune response, while β IL-3 receptor-deficient mice are normal. *Immunity* 2:211, 1995
- Robb L, Drinkwater CC, Metcalf D, Li R, Köntgen F, Nicola NA, Begley CG: Hematopoietic and lung abnormalities in mice with a null mutation of the common β subunit of the receptors for granulocyte-macrophage colony-stimulating factor and interleukins 3 and 5. *Proc Natl Acad Sci USA* 92:9565, 1995

28. Nicola NA, Robb L, Metcalf D, Cary D, Drinkwater CC, Begley CG: Functional inactivation in mice of the gene for the interleukin-3 (IL-3)-specific receptor β chain: Implications of IL-3 function and the mechanism of receptor transmodulation in hematopoietic cells. *Blood* 87:2665, 1996
29. Dranoff G, Crawford AD, Sadelain M, Ream B, Rashid A, Bronson RT, Dickersin GR, Bachurski CJ, Mark EL, Whitsett JA, Mulligan RC: Involvement of granulocyte-macrophage colony-stimulating factor in pulmonary hemostasis. *Science* 264:713, 1994
30. Stanley E, Lieschke GJ, Grail D, Metcalf D, Hodgson G, Gall JA, Maher DW, Cebon J, Sinickas V, Dunn AR: Granulocyte/macrophage colony-stimulating factor-deficient mice show no major perturbation of hematopoiesis but develop a characteristic pulmonary pathology. *Proc Natl Acad Sci USA* 91:5592, 1994
31. Shih JP, Zeng HQ, Ogawa M: Enrichment of murine marrow cells for progenitors of multilineage hematopoietic colonies. *Leukemia* 6:193, 1992
32. Ortega G, Korty PE, Shevach EM, Malek TR: Role of Ly-6 in lymphocyte activation. I. Characterization of a monoclonal antibody to a nonpolymorphic Ly-6 specificity. *J Immunol* 137:3240, 1986
33. Nishikawa S, Kusakabe M, Yoshinaga K, Ogawa M, Hayashi S, Kunisada T, Nishikawa S-I: In utero manipulation of coat color formation by a monoclonal anti-c-kit antibody: Two distinct waves of c-kit-dependency during melanocyte development. *EMBO J* 10:2111, 1991
34. Nakahata T, Ogawa M: Clonal origin of murine hemopoietic colonies with apparent restriction to granulocyte-macrophage-megakaryocyte (GMM) differentiation. *J Cell Physiol* 111:239, 1982
35. Harrison DE, Astle CM, Delaitre JA: Loss of proliferative capacity in immunohemopoietic stem cells caused by serial transplantation rather than aging. *J Exp Med* 147:1526, 1978
36. Coutinho LH, Will A, Radford J, Schiro R, Testa NG, Dexter TM: Effects of recombinant human granulocyte colony-stimulating factor (CSF), human granulocyte macrophage-CSF, and gibbon interleukin-3 on hematopoiesis in human long-term bone marrow culture. *Blood* 75:2118, 1990
37. Heimfeld S, Hudak S, Weissman I, Rennick D: The in vitro response of phenotypically defined mouse stem cells and myeloid progenitors to single or multiple growth factors. *Proc Natl Acad Sci USA* 88:9902, 1991
38. Bodine DM, Crosier PS, Clark SC: Effects of hematopoietic growth factors on the survival of primitive stem cells in liquid suspension culture. *Blood* 78:914, 1991
39. Miura N, Okada S, Zsebo KM, Miura Y, Suda T: Rat stem cell factor and IL-6 preferentially support the proliferation of c-kit-positive murine hematopoietic cells rather than their differentiation. *Exp Hematol* 21:143, 1993
40. Muench MO, Firpo MT, Moore MAS: Bone marrow transplantation with interleukin-1 plus kit-ligand ex vivo expanded bone marrow accelerates hematopoietic reconstitution in mice without the loss of stem cell lineage and proliferative potential. *Blood* 81:3463, 1993
41. Rebel VI, Dragowska W, Eaves CJ, Humphries RK, Lansdorp PM: Amplification of Sca-1⁺ Lin⁻WGA⁺ cells in serum-free cultures containing steel factor, interleukin-6, and erythropoietin with maintenance of cells with long-term in vivo reconstituting potential. *Blood* 83:128, 1994
42. Knobel KM, McNally MA, Berson AE, Rood D, Chen K, Kilinski L, Tran K, Okarma TB, Lebkowski JS: Long-term reconstitution of mice after ex vivo expansion of bone marrow cells: Differential activity of cultured bone marrow and enriched stem cell populations. *Exp Hematol* 22:1227, 1994
43. Peters SO, Kittler ELW, Ramshaw HS, Quesenberry PJ: Murine marrow cells expanded in culture with IL-3, IL-6, IL-11, and SCF acquire an engraftment defect in normal hosts. *Exp Hematol* 23:461, 1995
44. Brandt J, Briddell RA, Srour EF, Leemhuis TB, Hoffman R: Role of c-kit ligand in the expansion of human hematopoietic progenitor cells. *Blood* 79:634, 1992
45. Haylock DN, To LB, Dowse TL, Juttner CA, Simmons PJ: Ex vivo expansion and maturation of peripheral blood CD34⁺ cells into the myeloid lineage. *Blood* 80:1405, 1992
46. Sato N, Sawada K, Koizumi K, Tarumi T, Ieko M, Yasukouchi T, Yamaguchi M, Takahashi TA, Sekiguchi S, Koike T: In vitro expansion of human peripheral blood CD34⁺ cells. *Blood* 82:3600, 1993
47. Lansdorp PM, Dragowska W, Mayani H: Ontogeny-related changes in proliferative potential of human hematopoietic cells. *J Exp Med* 178:787, 1993
48. Srour EG, Brandt JE, Briddell RA, Grigsby S, Leemhuis T, Hoffman R: Long-term generation and expansion of human primitive hematopoietic progenitor cells in vitro. *Blood* 81:661, 1993
49. Brugger W, Möcklin W, Heimfeld S, Berenson RJ, Mertelsmann R, Kanz L: Ex vivo expansion of enriched peripheral blood CD34⁺ progenitor cells by stem cell factor, interleukin-1 β (IL-1 β), IL-6, IL-3, interferon- γ , and erythropoietin. *Blood* 81:2579, 1993
50. Flashshove M, Banerjee D, Mineishi S, Li MX, Bertino JR, Moore MAS: Ex vivo expansion and selection of human CD34⁺ peripheral blood progenitor cells after introduction of a mutated dihydrofolate reductase cDNA via retroviral gene transfer. *Blood* 85:566, 1995
51. Rice A, Boiron JM, Barbot C, Dupouy M, Dubosc-Marchenay N, Dumain P, Lacombe F, Reiffers J: Cytokine-mediated expansion of 5-FU-resistant peripheral blood stem cells. *Exp Hematol* 23:303, 1995
52. Peters SO, Kittler ELW, Ramshaw HS, Quesenberry PJ: Ex vivo expansion of murine marrow cells with interleukin-3 (IL-3), IL-6, IL-11, and stem cell factor leads to impaired engraftment in irradiated hosts. *Blood* 87:30, 1996
53. Brugger W, Heimfeld S, Berenson RJ, Mertelsmann R, Kanz L: Reconstitution of hematopoiesis after high-dose chemotherapy by autologous progenitor cells generated ex vivo. *N Engl J Med* 333:283, 1995
54. Williams SF, Lee WJ, Bender JG, Zimmerman T, Swinney P, Blake M, Carreon J, Schilling M, Smith S, Williams DE, Oldham F, Epps DV: Selection and expansion of peripheral blood CD34⁺ cells in autologous stem cell transplantation for breast cancer. *Blood* 87:1687, 1996
55. Donahue RE, Byrne ER, Thomas TE, Kirby MR, Agricola BA, Sellers SE, Gaudernack G, Karlsson S, Lansdorp PM: Transplantation and gene transfer of the human glucocerebrosidase gene into immunoselected primate CD34⁺ Thy-1⁺ cells. *Blood* 88:4166, 1996
56. Koike K, Ogawa M, Ihle JN, Miyake T, Shimizu T, Miyajima A, Yokota T, Arai K: Recombinant murine granulocyte-macrophage (GM) colony-stimulating factor supports formation of GM and multipotential blast cell colonies in culture: Comparison with the effects of interleukin-3. *J Cell Physiol* 131:458, 1987
57. Leary AG, Yang Y-C, Clark SC, Gasson JC, Golde DW, Ogawa M: Recombinant gibbon interleukin-3 supports formation of human multilineage colonies and blast cell colonies in culture: Comparison with recombinant human granulocyte-macrophage colony-stimulating factor. *Blood* 70:1343, 1987
58. McKinstry WJ, Li C-L, Rasko JEJ, Nicola NA, Johnson GR, Metcalf D: Cytokine receptor expression on hematopoietic stem and progenitor cells. *Blood* 89:65, 1997