

Incidence of Type 2 Diabetes Using Proposed HbA_{1c} Diagnostic Criteria in the European Prospective Investigation of Cancer–Norfolk Cohort

Implications for preventive strategies

PARINYA CHAMNAN, MD¹
REBECCA K. SIMMONS, PHD¹
NITA G. FOROUHI, MD¹
ROBERT N. LUBEN, BSC²

KAY-TEE KHAW, PHD³
NICHOLAS J. WAREHAM, PHD¹
SIMON J. GRIFFIN, DM¹

OBJECTIVE—To evaluate the incidence and relative risk of type 2 diabetes defined by the newly proposed HbA_{1c} diagnostic criteria in groups categorized by different baseline HbA_{1c} levels.

RESEARCH DESIGN AND METHODS—Using data from the European Prospective Investigation of Cancer (EPIC)–Norfolk cohort with repeat HbA_{1c} measurements, we estimated the prevalence of known and previously undiagnosed diabetes at baseline (baseline HbA_{1c} \geq 6.5%) and the incidence of diabetes over 3 years. We also examined the incidence and corresponding odds ratios (ORs) by different levels of baseline HbA_{1c}. Incident diabetes was defined clinically (self-report at follow-up, prescribed diabetes medication, or inclusion on a diabetes register) or biochemically (HbA_{1c} \geq 6.5% at the second health assessment), or both.

RESULTS—The overall prevalence of diabetes was 4.7%; 41% of prevalent cases were previously undiagnosed. Among 5,735 participants without diabetes at baseline (identified clinically or using HbA_{1c} criteria, or both), 72 developed diabetes over 3 years (1.3% [95% CI 1.0–1.5]), of which 49% were identified using the HbA_{1c} criteria. In 6% of the total population, the baseline HbA_{1c} was 6.0–6.4%; 36% of incident cases arose in this group. The incidence of diabetes in this group was 15 times higher than in those with a baseline HbA_{1c} of $<$ 5.0% (OR 15.5 [95% CI 7.2–33.3]).

CONCLUSIONS—The cumulative incidence of diabetes defined using a newly proposed HbA_{1c} threshold in this middle-aged British cohort was 1.3% over 3 years. Targeting interventions to individuals with an HbA_{1c} of 6.0–6.4% might represent a feasible preventive strategy, although complementary population-based preventive strategies are also needed to reduce the growing burden of diabetes.

Diabetes Care 34:950–956, 2011

Type 2 diabetes is a major public health concern worldwide. An estimated 439 million people will have the disease by 2030 (1). It is possible to halve the incidence of type 2 diabetes among individuals at high risk through

lifestyle and pharmacologic interventions (2–4). However, it is unlikely that population screening for impaired glucose tolerance using an oral glucose tolerance test (OGTT) is a feasible method of identifying those at high risk in clinical practice

(5) because it is time- and resource-consuming and has poor reproducibility (6). If a measure of blood glucose were to be used to define the risk of developing diabetes, then it would seem logical to use the same test for diagnosis and informing treatment decisions (7).

Glycated hemoglobin (HbA_{1c}) is a reliable measure of long-term glycemic exposure (8) that correlates well with the risk of microvascular and macrovascular complications of diabetes (9,10). It does not necessitate fasting or timed blood samples. Previous concerns regarding the standardization of assays have largely been resolved (11). Consequently, the American Diabetes Association (ADA) recently recommended that HbA_{1c} be included as a diagnostic test for diabetes, with a diagnostic threshold of \geq 6.5% (12).

To better estimate the burden of the disease and potential benefits of preventive interventions, it is necessary to have accurate data on incidence. Reported estimates of incidence in adult populations have varied considerably, from 2 to 25 per 1,000 person-years (13–16). However, many of the studies were restricted to high-risk populations (13) and defined diabetes using a clinical rather than a biochemical diagnosis (14). Fewer studies have investigated the incidence of diabetes based on longitudinal repeat blood glucose measurements in population-based samples (15,16), and none have examined diabetes incidence using repeated measures of HbA_{1c}.

In this study we estimated the prevalence and incidence of diabetes defined using the newly proposed HbA_{1c} cutoff of 6.5% in a population-based British cohort. To inform the choice of appropriate HbA_{1c} thresholds to identify individuals at high risk to whom preventive interventions might be offered, we used data on longitudinal repeat HbA_{1c} values at baseline and after 3 years of follow-up to examine the incidence and relative risk of clinically diagnosed diabetes and diabetes

From the ¹MRC Epidemiology Unit, Institute of Metabolic Science, Addenbrooke's Hospital, Cambridge, U.K.; the ²Department of Public Health and Primary Care, Institute of Public Health, University of Cambridge, Cambridge, U.K.; and the ³Clinical Gerontology Unit, University of Cambridge School of Clinical Medicine, Addenbrooke's Hospital, Cambridge, U.K.

Corresponding author: Simon J. Griffin, simon.griffin@mrc-epid.cam.ac.uk.

Received 19 December 2009 and accepted 2 July 2010.

DOI: 10.2337/dc09-2326

This article contains Supplementary Data online at <http://care.diabetesjournals.org/lookup/suppl/doi:10.2337/dc09-2326/-/DC1>.

© 2011 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

defined using HbA_{1c} diagnostic criteria in groups defined by different baseline HbA_{1c} values.

RESEARCH DESIGN AND METHODS

Study design and population

The European Prospective Investigation of Cancer (EPIC)-Norfolk is a population-based prospective study that monitors 25,639 men and women aged 40–74 years residing in the Norfolk region, U.K. Details of the study have been described elsewhere (17). In brief, between 1993 and 1997, 77,630 individuals were recruited from general practice to participate in the study, and 25,639 (33%) consented and attended a baseline health assessment.

Participants completed questionnaires about their personal and family history of disease, medication, and lifestyle factors, including smoking habits. They were asked whether a physician had ever told them that they had any of the conditions in a list that included diabetes, heart attack, and stroke. Baseline diabetes status was ascertained by 1) self-report of diabetes medication, 2) diabetes medication brought to the baseline examination, 3) participants indicating modification of their diet in the past year because of diabetes, or 4) participants indicating adherence to a diabetic diet. Anthropometric and blood pressure measurements, as well as nonfasting blood samples were taken at the health assessment.

Because funding for measurement of HbA_{1c} only became available in 1995, about 50% of all participants had information on this measure at baseline. HbA_{1c} was measured on fresh EDTA blood samples using high-performance liquid chromatography (Diamat Automated Glycated Hemoglobin Analyzer; Bio-Rad Laboratories Ltd., Hemel Hemstead, U.K.), which was standardized to the Diabetes Control and Complications Trial (DCCT) assay.

Participants were invited to attend a second health assessment after 3 years (1998–2001), at which identical measurements were taken, and 15,028 participants (59%) attended. General practitioners of participants whose HbA_{1c} test results exceeded $\geq 7.0\%$ were notified so that they could assume responsibility for confirming diagnosis and arranging treatment. The study was approved by the Norwich District Health Authority Ethics Committee. All participants gave signed informed consent.

Individuals who live the Norfolk area are slightly healthier than those in the general U.K. population, with a standardized mortality ratio of 93 (Office for National Statistics death registration data, 2008). However, EPIC-Norfolk participants are similar to a nationally representative sample regarding anthropometric indices, blood pressure, and serum lipid levels (17).

We report results for follow-up at the second health assessment, a median of 3 years. We limited our analyses to the 6,372 individuals with HbA_{1c} measurements at baseline and at the second health assessment. We used this study sample to estimate the prevalence of known (clinically diagnosed diabetes, self-reported physician-diagnosed diabetes, and diabetes medication) and previously undiagnosed diabetes at baseline (baseline HbA_{1c} $\geq 6.5\%$). After excluding those with diabetes at baseline (clinically diagnosed diabetes and diabetes defined using HbA_{1c} diagnostic criteria), we further excluded 335 individuals with missing data for other metabolic risk factors, including age, sex, a family history of diabetes, smoking, the use of corticosteroids and antihypertensive drugs, BMI, waist circumference, systolic blood pressure, cholesterol, and triglyceride, leaving 5,735 individuals for analyses of the incidence and risk of diabetes (Fig. 1).

Ascertainment of incident diabetes

Participants were identified as having incident diabetes if 1) they reported physician-diagnosed diabetes or diabetes medication, or brought diabetes medication to the second health assessment (clinical diagnosis), 2) they were identified on medical records, diabetes registers, or death certificates (clinical diagnosis) or 3) they had an HbA_{1c} of $\geq 6.5\%$ at the second health assessment (HbA_{1c}-defined diabetes). Participants were identified through their general practice diabetes register or the Norfolk and Norwich Hospital diabetes register. Participants admitted to a hospital with a diabetes-related condition were identified by their National Health Service number. Hospitals were linked to the East Norfolk Health Authority database, which identifies all hospital contacts throughout England and Wales for Norfolk residents. Vital status for all EPIC-Norfolk participants was obtained through death certification at the Office for National Statistics, and death certification with coding for diabetes was identified. Previous validation studies in this

cohort using capture–recapture analysis indicated that the use of multiple sources of ascertainment information for diabetes detected 99% of incident cases when comparing with diagnostic information from a comprehensive review of medical records (18).

Statistical analyses

In 5,735 participants free of diabetes at baseline with data on HbA_{1c} for the baseline and second health assessments, we calculated the incidence of diabetes defined clinically and by using HbA_{1c} diagnostic criteria in the whole cohort and separately for different categories of baseline HbA_{1c}. Baseline characteristics were summarized for groups defined by different categories of baseline HbA_{1c} (<5.0, 5.0–5.4, 5.5–5.9, and 6.0–6.4%). We tested for differences between groups using the χ^2 test for categorical variables and ANOVA or Kruskal-Wallis tests for normally or nonnormally distributed continuous variables, respectively.

We used logistic regression to estimate the risk of developing diabetes as measured by the odds ratios (ORs) for every 0.5% increase in HbA_{1c} as well as for different categories of HbA_{1c} compared with the lowest HbA_{1c} category of <5.0%. We examined ORs adjusted for age only, age and sex only, and multiple risk factors, including age, sex, self-reported family history of diabetes, smoking, the use of antihypertensive drugs or corticosteroids, BMI, waist circumference, systolic blood pressure, and total cholesterol, HDL cholesterol, and triglyceride values. To inform alternative screening strategies, we investigated risk factors associated with incident diabetes in those with a baseline HbA_{1c} of <6.0%.

We also performed a sensitivity analysis using a more restricted definition of incident diabetes in which participants were not classified as having incident diabetes unless a self-reported diagnosis was supported by information on diabetes-specific medication or confirmed by information from clinical records, death certificates, or HbA_{1c}.

RESULTS—Table 1 summarizes baseline characteristics of participants in the EPIC-Norfolk cohort by different HbA_{1c} categories. Participants (45% men) had a mean age of 57.4 (SD 9.4) years. Participants with a higher HbA_{1c} value were older, more likely to be male, obese, current smokers, and to come from a lower socioeconomic class than those with a




Figure 1—Schematic diagram demonstrates the numbers and percentages of individuals with prevalent and incidence diabetes in a cohort of 6,372 men and women over 3 years. Individuals with clinically diagnosed diabetes and HbA_{1c} ≥6.5% were considered to have clinically diagnosed diabetes in this diagram. *Self-reported diabetes, evidence of diabetes medications, and dietary modification due to diabetes. †Self-reported diabetes, evidence of diabetes medication, diabetes registers, hospitalizations with diabetes, and diabetes codes on death certificates.

lower HbA_{1c} value. They were also more likely to have higher blood pressure, higher total cholesterol and triglyceride values, and lower HDL cholesterol values. There was no difference in family history of diabetes and the use of corticosteroids among the groups.

Prevalence of diabetes at baseline

Among 6,372 individuals with HbA_{1c} measurements at both health assessments, 302 (4.7%) had prevalent diabetes at baseline (Fig. 1). Among these individuals, 178 (2.8%) had known diabetes (those identified clinically), and 124 (1.9%) had previously undiagnosed diabetes (those identified using HbA_{1c} criteria).

Incidence of diabetes over 3 years

Among 5,735 participants free of diabetes at baseline, 72 developed diabetes over 3 years (Fig. 1). The cumulative incidence was 1.3% (95% CI 1.0–1.5) over 3 years, an annual incidence of 0.4%. Among these new cases of diabetes, 37 individuals (51%) were identified clinically (e.g.,

by their response to the questionnaire at the second health check or through linkage to clinical records or diabetes registers), with an incidence of 0.6% (0.4–0.9). The remaining 35 individuals (49%) with incident diabetes were identified by their HbA_{1c} results at the second health assessment.

Risk of developing diabetes in groups defined by different HbA_{1c} levels

Table 2 reports the incidence of diabetes by baseline HbA_{1c} levels for clinical diagnosis only and for clinical or HbA_{1c}-defined diagnosis, or both. The incidence of diabetes increased progressively with increasing baseline HbA_{1c} levels. In those with a baseline HbA_{1c} of 6.0–6.4%, the incidence of clinically diagnosed or HbA_{1c}-defined diabetes was three times higher than that of clinically diagnosed diabetes, at 7.0 (95% CI 4.8–10.1) and 2.4 (1.3–4.6), respectively. Thirty-six percent of incident cases of diabetes arose from individuals with a baseline HbA_{1c} of 6.0–6.4% (6% of the total population), and just over 35% of incident cases arose

among individuals with a baseline HbA_{1c} of <5.5% (69% of the total population).

Significant positive associations were found between HbA_{1c} and the risk of developing diabetes (Table 2). A 0.5% increase in baseline HbA_{1c} was associated with more than a twofold increase in the risk of clinically diagnosed or HbA_{1c}-defined diabetes, or both (age-adjusted OR 2.7 [95% CI 2.1–3.5]). Participants with a baseline HbA_{1c} of 6.0–6.4% had about a sevenfold higher risk of clinically diagnosed diabetes than those with an HbA_{1c} of <5.0%. The highest risk of clinically diagnosed or HbA_{1c}-defined diabetes was observed in the highest baseline HbA_{1c} category compared with those with a baseline HbA_{1c} of <5.0% (OR 15.5 [7.2–33.3]). These ORs remained unchanged after adjustment for other risk factors.

Among individuals with a baseline HbA_{1c} <6.0%, a family history of diabetes and waist circumference were the strongest nonlaboratory predictors of incident diabetes over 3 years.

Table 1—Comparison of baseline characteristics across categories of baseline HbA_{1c} in 5,735 participants in the EPIC-Norfolk cohort

Variable	Total	HbA _{1c} level				P for difference*
		<5.0%	5.0–5.4%	5.5–5.9%	6.0–6.4%	
N (%)	5,735 (100)	1,849 (32.2)	2,119 (36.9)	1,397 (24.4)	370 (6.5)	
Age (years)	57.4 (9.4)	54.1 (9.2)	57.4 (9.1)	60.3 (8.7)	62.4 (8.2)	<0.001
Men, n (%)	2,481 (43.3)	746 (40.4)	932 (44.0)	634 (45.4)	169 (45.7)	0.016
Social class, † n (%)						<0.001
Class I–IIIa	3,694 (64.4)	1,255 (67.9)	1,358 (64.1)	861 (61.6)	220 (59.5)	
Class IIIb–V	2,041 (35.6)	594 (32.1)	761 (35.9)	536 (38.4)	150 (40.5)	
Current smokers, n (%)	525 (9.2)	141 (7.6)	165 (7.8)	165 (11.8)	54 (14.6)	<0.001
Family history of diabetes, n (%)	697 (12.2)	217 (11.7)	250 (11.8)	177 (12.7)	53 (14.3)	0.469
Medication use, n (%)						
Corticosteroids	156 (2.7)	44 (2.4)	61 (2.9)	37 (2.7)	14 (3.8)	0.454
Antihypertensive drugs	811 (14.1)	188 (10.2)	301 (14.2)	243 (17.4)	79 (21.4)	<0.001
BMI (kg/m ²)	25.9 (3.7)	25.4 (3.5)	25.9 (3.6)	26.3 (3.8)	26.6 (4.0)	<0.001
BMI category, n (%)						<0.001
<25 kg/m ²	2,528 (44.1)	935 (50.6)	906 (42.8)	553 (39.6)	134 (36.2)	
25–29.9 kg/m ²	25,24 (44.0)	738 (39.9)	956 (45.1)	657 (47.0)	173 (46.8)	
≥30 kg/m ²	683 (11.9)	176 (9.5)	257 (12.1)	187 (13.4)	63 (17.0)	
Waist circumference, cm	86.9 (12.2)	84.8 (12.2)	87.1 (11.9)	88.6 (11.9)	90.0 (13.0)	<0.001
Systolic blood pressure, mmHg	133.4 (17.5)	130.6 (17.1)	133.4 (17.1)	135.7 (18.1)	138.9 (17.3)	<0.001
Diastolic blood pressure, mmHg	82.0 (10.8)	80.9 (10.6)	82.1 (10.8)	82.9 (11.1)	84.1 (10.9)	<0.001
Total cholesterol, mmol/L	6.1 (1.1)	5.8 (1.1)	6.1 (1.1)	6.3 (1.1)	6.4 (1.2)	<0.001
HDL cholesterol, mmol/L	1.5 (0.4)	1.5 (0.4)	1.5 (0.4)	1.4 (0.4)	1.4 (0.4)	0.001
TG, median (IQR), mmol/L	1.4 (1.0–2.1)	1.3 (0.9–1.8)	1.5 (1.0–2.1)	1.6 (1.1–2.2)	1.7 (1.2–2.5)	<0.001

Data are presented as the mean (SD), unless specified otherwise. IQR, interquartile range; TG, triglyceride. *Differences between groups using χ^2 tests for categorical variables, and ANOVA or Kruskal-Wallis tests for normally or nonnormally distributed continuous variables. †Registrar General's Social Class: class I = professional, etc. occupations; II = managerial and technical occupations; IIIa = skilled occupations (nonmanual); IIIb = skilled occupations (manual); IV = partly-skilled occupations; V = unskilled occupations.

In the sensitivity analysis using a more restricted definition of incident diabetes, 59 individuals developed diabetes (Supplementary Table 1). The incidence was 1.0 (95% CI 0.8–1.3) over 3 years, an annual incidence of 0.3%. Incident cases in 37 individuals (63%) were identified using HbA_{1c} diagnostic criteria. Approximately 40% of incident cases of diabetes developed in individuals with a high baseline HbA_{1c} of 6.0–6.4% (6% of total population). A 27-fold higher risk of diabetes was observed in those with a baseline HbA_{1c} of 6.0–6.4% compared with those with a baseline HbA_{1c} of <5.0%.

CONCLUSIONS—We used data from a large population-based British prospective cohort to estimate the prevalence and incidence of diabetes defined clinically or by HbA_{1c}, or both, over 3 years. The HbA_{1c} diagnostic threshold of 6.5% identified 50% of new cases. The incidence of diabetes increased progressively across baseline HbA_{1c} levels, with 36% of incident cases developing in individuals with a baseline HbA_{1c} of 6.0–6.4%.

To the best of our knowledge, this study is the first to report the incidence of

diabetes based on HbA_{1c} diagnostic criteria using repeated assessment of HbA_{1c}. Our prevalence estimate is comparable with the prevalence of diabetes in England (4.4%) estimated from an epidemiologic model in which individuals with known and previously undiagnosed diabetes were included (19). We also found that approximately 40% of prevalent cases were identified using HbA_{1c} diagnostic criteria and, hence, were previously undiagnosed. This is consistent with previous studies using an OGTT as a screening test (20).

A number of studies have estimated diabetes incidence by using longitudinal repeat OGTT or fasting plasma glucose measurements (15,16). The incidence of diabetes in these studies varied from 6 to 10% over 9 to 10 years. Data from a British population in Ely, Cambridgeshire, showed that the cumulative incidence of diabetes using repeat OGTT measurements was 5.9% over 10 years, corresponding to an annual incidence of 0.6% (16). This is comparable with the low annual incidence of 0.4% in the current study. The higher incidence in the earlier study may be explained by enhanced case

detection from repeated testing by OGTT over a longer period (OGTT testing at baseline, 4.5, and 10 years in the earlier Ely study vs. HbA_{1c} at baseline and after 3 years of follow-up in the EPIC-Norfolk study) and the different contributions of the “healthy volunteer effect” in each study (response rates of 74% in the Ely study and 33% in the EPIC-Norfolk study).

Few studies have examined the incidence and relative risk of diabetes in individuals or groups defined by different baseline HbA_{1c} levels. Selvin et al. (21) examined the incidence of self-reported diabetes in American men and women with different baseline HbA_{1c} values. The 15-year cumulative incidence of diabetes was 6, 12, 21, and 44% in individuals with an HbA_{1c} of <5.0, 5.0–5.4, 5.5–5.9, and 6.0–6.4%, respectively. The estimated annual incidence was higher than those observed across all HbA_{1c} categories in our study. This might be explained by the differences in levels of other risk factors (higher BMI, smoking, and family history of diabetes in the Selvin study), follow-up time (15 vs. 3 years, if incidence rates are not consistent across different durations of

Table 2—Incidence and risk (OR) of diabetes over 3 years by baseline HbA_{1c} categories in the EPIC-Norfolk cohort (N = 5,735)

Variable	Total	HbA _{1c} level				P for trend
		<5.0%	5.0–5.4%	5.5–5.9%	6.0–6.4%	
Clinically diagnosed diabetes						
N (% of total participants)	5,735	1,849 (32)	2,119 (37)	1,397 (24)	370 (6)	—
Incident cases (% of total cases)	37	7 (19)	6 (16)	15 (41)	9 (24)	—
Three-year cumulative incidence, %	0.6 (0.4–0.9)	0.4 (0.2–0.8)	0.3 (0.1–0.6)	1.1 (0.6–1.8)	2.4 (1.3–4.6)	—
Unadjusted OR		1.0	0.7 (0.3–2.2)	2.9 (1.2–7.0)	6.6 (2.4–17.7)	<0.001
Age-adjusted OR		1.0	0.8 (0.3–2.4)	3.3 (1.3–8.4)	8.0 (2.8–22.7)	<0.001
Age- and sex-adjusted OR		1.0	0.8 (0.3–2.4)	3.3 (1.3–8.3)	7.9 (2.8–22.4)	<0.001
Multivariable-adjusted OR*		1.0	0.8 (0.3–2.3)	3.0 (1.2–7.8)	6.8 (2.3–20.1)	<0.001
Clinically diagnosed and/or HbA _{1c} -defined diabetes						
N (% of total participants)	5,735	1,849 (32)	2,119 (37)	1,397 (24)	370 (6)	—
Incident cases (% of total cases)	72	9 (13)	16 (22)	21 (29)	26 (36)	—
Three-year cumulative incidence, %	1.3 (1.0–1.5)	0.5 (0.3–0.9)	0.8 (0.5–1.2)	1.5 (1.0–2.3)	7.0 (4.8–10.1)	—
Unadjusted OR		1.0	1.6 (0.7–3.5)	3.1 (1.4–6.8)	15.5 (7.2–33.3)	<0.001
Age-adjusted OR		1.0	1.6 (0.7–3.8)	3.5 (1.6–7.8)	18.0 (8.1–40.0)	<0.001
Age- and sex-adjusted OR		1.0	1.6 (0.7–3.7)	3.4 (1.5–7.7)	17.7 (8.0–39.5)	<0.001
Multivariable-adjusted OR*		1.0	1.6 (0.7–3.6)	3.3 (1.5–7.4)	15.6 (6.9–35.7)	<0.001

*Adjusted for age, sex, social class, self-reported family history of diabetes, smoking, use of corticosteroids and antihypertensive drugs, BMI, waist circumference, systolic blood pressure, cholesterol, HDL cholesterol, and triglyceride.

follow-up), and in particular, the different definitions of diabetes used in each study.

HbA_{1c} has been shown to be a useful tool for the early detection of diabetes (21). A few studies have demonstrated that HbA_{1c} predicts future risk of diabetes in high-risk individuals with glucose intolerance (22). We have shown that HbA_{1c} predicts risk of diabetes in healthy middle-aged men and women. An HbA_{1c} of 6.0–6.4% identified 24% of clinically incident diabetes, and 36% of clinically incident or HbA_{1c}-defined diabetes. These figures were even higher when a more restricted definition of incident diabetes was used. Recent evidence shows that this predictive ability also holds true in low-risk nondiabetic men and women (21) and in elderly individuals (23). However, given that the associations between HbA_{1c} and the risk of diabetes were hardly changed after adjustment for multiple risk factors, there may not be much to gain from including data on multiple risk factors alongside HbA_{1c} for prediction of diabetes risk.

ADA suggested that there was no specific threshold that defines individuals who might be offered preventive interventions and that any such threshold would vary between countries with different health care priorities (12). However, ADA suggested that individuals with an HbA_{1c} between 6.0 and 6.4% might represent a group in whom the risk of

development of diabetes was very high and who could therefore be targeted for individual prevention interventions (12). ADA also suggested that this range should not be considered an absolute threshold and that interventions may be appropriate in other individuals on the basis of other risk information. Our findings support this statement by demonstrating that most new cases of diabetes developed in those with a baseline HbA_{1c} of <6.0%.

The selection of a population for a high-risk prevention strategy is by the level of risk identified, the proportion of the population to be targeted, and the proportion of future cases that might therefore be prevented. Our study showed that 36% of new cases of diabetes arose from the 6% of the study population who had the highest glycemic levels (HbA_{1c} 6.0–6.4%). Indeed, if previously proven intensive prevention interventions (2,3) were targeted at this middle-aged population, approximately 20% of new cases of diabetes could be prevented over 3 years. Strategies for identifying which individuals should have an HbA_{1c} measurement, including simple risk scores using easily measured or routinely available risk factors, are needed.

Although the category of people with an HbA_{1c} of 6.0–6.4% identifies a high-risk group, most new cases of diabetes developed in individuals whose baseline HbA_{1c} values were <6%. Complementary

strategies to identify high-risk individuals among those without raised HbA_{1c} may therefore be necessary. Our subgroup analysis in individuals with an HbA_{1c} of <6.0% suggested that those with central obesity and a family history of diabetes might represent another relatively easily identifiable subgroup to whom preventive interventions could be targeted. This also suggests that in addition to high-risk approaches, we need to develop a complementary population-based strategy aimed at shifting the whole distribution of HbA_{1c} in the population to reduce the risk of both diabetes and its complications (24). However, although there is some evidence for the cost-effectiveness of prevention interventions among high risk individuals (25), evidence on the cost-effectiveness of population-based strategies is very limited, making judgments about the balance of investment in high-risk and population-based approaches difficult.

We have reported the incidence of diabetes in a large prospective British cohort using clinical ascertainment and newly proposed HbA_{1c} diagnostic criteria. Participants included in this analysis were healthier than those excluded; hence, our findings are likely to underestimate the incidence of diabetes in the whole cohort. Given the 33% recruitment rate in this study, it is possible that participants might be more health-conscious and more likely to engage in healthy behaviors and to take

up existing preventive services compared with nonparticipants. We might therefore have underestimated the overall incidence of diabetes in the population, which might consequently influence estimates of the relative risk for HbA_{1c}.

Because the EPIC-Norfolk study collected nonfasting blood samples, it is not possible to compare the predictive ability for incident diabetes of different measures of glycemia. Similar to other diagnostic tests, the use of a single measure of HbA_{1c} for diagnosing diabetes might lead to some degree of misclassification. However, given that HbA_{1c} is more reliable compared with an OGTT and the fasting plasma glucose test, the misclassification is likely to be only modest. A relatively short follow-up period and relatively small number of events mean that our finding should be interpreted with caution. However, our study is one of the largest incidence studies reported.

Furthermore, the follow-up of 3 years is still a plausible and important time-frame for identifying those at high risk of diabetes, because in addition to long-term risk information, one might also be interested in and more persuaded for behavior modification by information on the short-term risk of diabetes. Our findings were specific to a population aged 40–74 years, and may not represent the burden and risk of diabetes in relation to HbA_{1c} levels in younger people. Lastly, most of the EPIC-Norfolk participants are of European descent, which limits the generalizability of our findings to other ethnic groups and populations.

In conclusion, the cumulative incidence of diabetes defined using a newly proposed HbA_{1c} threshold in this middle-aged British cohort was 1.3% over 3 years (0.4% per year). HbA_{1c} independently predicted the risk of incident diabetes, with each 0.5% difference in HbA_{1c} being associated with more than doubling of the risk of diabetes. Because 36% of incident cases of diabetes came from the 6% of the population with a baseline HbA_{1c} of between 6 and 6.5%, this may be an easily identifiable subgroup to whom preventive interventions could be targeted. Alternative strategies to identify high-risk individuals may be necessary, however, and complementary population-based approaches need to be developed to shift the underlying distribution of glycemia.

Acknowledgments—Funding support was provided by the Medical Research Council,

Cancer Research UK, British Heart Foundation, European Union (Europe Against Cancer Programme), Stroke Association, Wellcome Trust, Research into Ageing, and Academy of Medical Science. P.C. is supported by a Royal Thai Government scholarship. S.J.G. receives support from the National Institute for Health Research programme grant funding scheme (RP-PG-0606-1259). The views expressed in this publication are those of the authors and not necessarily those of the NHS, National Institute for Health Research, or Department of Health. The sponsors did not participate in the design or conduct of this study; in the collection, management, analysis, or interpretation of data; in the writing of the manuscript; or in the preparation, review, approval, or decision to submit this manuscript for publication.

No potential conflicts of interest relevant to this article were reported.

P.C. researched data, contributed to discussion, wrote the manuscript, and reviewed and edited the manuscript. R.K.S., N.G.F., and R.N.L. researched data, contributed to discussion, and reviewed and edited the manuscript. K.-T.K. researched data and reviewed and edited the manuscript. N.J.W. and S.J.G. researched data, contributed to discussion, and reviewed and edited the manuscript.

Parts of this study were presented at the 45th Annual Meeting of the European Diabetes Epidemiology Group, Porto Heli, Greece, 15–18 May 2010.

The authors thank the EPIC-Norfolk participants and the EPIC-Norfolk team for their contributions.

References

- International Diabetes Federation. *The Diabetes Atlas*. 4th ed. Brussels, International Diabetes Federation, 2009
- Tuomilehto J, Lindström J, Eriksson JG, et al.; Finnish Diabetes Prevention Study Group. Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med* 2001;344:1343–1350
- Knowler WC, Barrett-Connor E, Fowler SE, et al.; Diabetes Prevention Program Research Group. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002;346:393–403
- Li G, Zhang P, Wang J, et al. The long-term effect of lifestyle interventions to prevent diabetes in the China Da Qing Diabetes Prevention Study: a 20-year follow-up study. *Lancet* 2008;371:1783–1789
- Ruge T, Nyström L, Lindahl B, et al. Recruiting high-risk individuals to a diabetes prevention program: how hard can it be? *Diabetes Care* 2007;30:e61
- Ko GT, Chan JC, Woo J, et al. The reproducibility and usefulness of the oral glucose tolerance test in screening for diabetes and other cardiovascular risk factors. *Ann Clin Biochem* 1998;35:62–67
- Colagiuri S, Borch-Johnsen K, Wareham NJ. Back to the future—do IGT and IFG have value as clinical entities? *Diabetes Res Clin Pract* 2008;81:131–133
- Nathan DM, Turgeon H, Regan S. Relationship between glycated haemoglobin levels and mean glucose levels over time. *Diabetologia* 2007;50:2239–2244
- Sabanayagam C, Liew G, Tai ES, et al. Relationship between glycated haemoglobin and microvascular complications: is there a natural cut-off point for the diagnosis of diabetes? *Diabetologia* 2009;52:1279–1289
- Khaw KT, Wareham N, Bingham S, Luben R, Welch A, Day N. Association of hemoglobin A1c with cardiovascular disease and mortality in adults: the European prospective investigation into cancer in Norfolk. *Ann Intern Med* 2004;141:413–420
- Kahn R, Fonseca V. Translating the A1C Assay. *Diabetes Care* 2008;31:1704–1707
- American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2010;33(Suppl. 1):S62–S69
- Söderberg S, Zimmet P, Tuomilehto J, et al. High incidence of type 2 diabetes and increasing conversion rates from impaired fasting glucose and impaired glucose tolerance to diabetes in Mauritius. *J Intern Med* 2004;256:37–47
- Geiss LS, Pan L, Cadwell B, Gregg EW, Benjamin SM, Engelgau MM. Changes in incidence of diabetes in U.S. adults, 1997–2003. *Am J Prev Med* 2006;30:371–377
- Bonora E, Kiechl S, Willeit J, et al.; Bruneck study. Population-based incidence rates and risk factors for type 2 diabetes in white individuals: the Bruneck study. *Diabetes* 2004;53:1782–1789
- Forouhi NG, Luan J, Hennings S, Wareham NJ. Incidence of Type 2 diabetes in England and its association with baseline impaired fasting glucose: the Ely study 1990–2000. *Diabet Med* 2007;24:200–207
- Day N, Oakes S, Luben R, et al. EPIC-Norfolk: study design and characteristics of the cohort. European Prospective Investigation of Cancer. *Br J Cancer* 1999;80(Suppl. 1):95–103
- Harding AH, Day NE, Khaw KT, et al. Dietary fat and the risk of clinical type 2 diabetes: the European prospective investigation of Cancer-Norfolk study. *Am J Epidemiol* 2004;159:73–82
- Forouhi NG, Merrick D, Goyder E, et al. Diabetes prevalence in England, 2001—estimates from an epidemiological model. *Diabet Med* 2006;23:189–197
- King H, Rewers M; WHO Ad Hoc Diabetes Reporting Group. Global estimates for prevalence of diabetes mellitus and impaired glucose tolerance in adults. *Diabetes Care* 1993;16:157–177

HbA_{1c}-defined incidence of diabetes

21. Selvin E, Steffes MW, Zhu H, et al. Glycated hemoglobin, diabetes, and cardiovascular risk in nondiabetic adults. *N Engl J Med* 2010;362:800–811
22. Little RR, England JD, Wiedmeyer HM, et al. Glycated haemoglobin predicts progression to diabetes mellitus in Pima Indians with impaired glucose tolerance. *Diabetologia* 1994;37:252–256
23. Motta M, Bennati E, Cardillo E, Ferlito L, Malaguarnera M. The value of glycosylated hemoglobin (HbA_{1c}) as a predictive risk factor in the diagnosis of diabetes mellitus (DM) in the elderly. *Arch Gerontol Geriatr* 2010;50:60–64
24. Simmons RK, Griffin SJ, Wareham NJ. Researching how to realize the potential of diabetes prevention. *Diabet Med* 2007;24:1055–1057
25. Herman WH, Hoerger TJ, Brandle M, et al.; Diabetes Prevention Program Research Group. The cost-effectiveness of lifestyle modification or metformin in preventing type 2 diabetes in adults with impaired glucose tolerance. *Ann Intern Med* 2005;142:323–332