

Molecular Pathways: Linking Tumor Microenvironment to Epithelial–Mesenchymal Transition in Metastasis

Hae-Yun Jung¹, Laurent Fattet¹, and Jing Yang^{1,2}

Abstract

During tumor development, tumor cells constantly communicate with the surrounding microenvironment through both biochemical and biophysical cues. In particular, the tumor microenvironment can instruct carcinoma cells to undergo a morphogenesis program termed epithelial-to-mesenchymal transition (EMT) to facilitate local invasion and metastatic dissemination. Growing evidence uncovered a plethora of microenvironmental factors in promoting EMT, including proinflammatory cytokines secreted by locally activated stromal cells, hypoxia conditions, extracellular matrix components, and mechanical properties. Here, we review various biochem-

ical and biophysical factors in the tumor microenvironment that directly impinge upon the EMT program. Specifically, cytokines such as TGF β , TNF α , and IL6 and hypoxia are capable of inducing EMT in various tumors. Several extracellular matrix (ECM) proteins, including collagen-I, fibronectin, and hyaluronan, and ECM remodeling via extracellular lysyl oxidase are also implicated in regulating EMT. In preclinical studies and ongoing clinical trials, targeting these tumor microenvironmental signals has shown promises in halting tumor progression in various human cancers. *Clin Cancer Res*; 21(5): 962–8. ©2014 AACR.

Background

During tumor metastasis, the epithelial-to-mesenchymal transition (EMT) program has been indicated in giving rise to the dissemination of single tumor cells from primary epithelial tumors (1). EMT refers to a global cellular and molecular transition by which polarized epithelial cells gain mesenchymal properties to migrate. During EMT, epithelial cells reorganize cytoskeleton and resolve cell–cell junctions, which are accompanied with switching off the expression of epithelial markers and turning on mesenchymal genes. Although changes in epithelial and mesenchymal markers during EMT can vary significantly in different biologic contexts, a network of transcription factors, including TWIST1/2, SNAIL1/2, ZEB1/2, and FOXC2, are consistently required to orchestrate the EMT program (2). Numerous studies have shown that the expression of these transcription factors is associated with poor prognosis and distant metastasis in various human cancers (3). Besides its role in promoting tumor cell invasion, EMT is shown to confer tumor cells with resistance to apoptosis (4) and anoikis (5), thus allowing cell survival in the blood stream after intravasation. EMT could also facilitate tumor cells' escape from the senescence program, especially through TWIST1 and ZEB1 (6, 7). Furthermore, EMT has been shown to

endow cancer cells with cancer stem cell (CSC)–like features, which further aid tumor dormancy and chemoresistance (8, 9).

Studies with tumor samples or experimental tumor xenograft models have provided convincing evidence for the activation of EMT in various primary epithelial tumors. Interestingly, more recent studies reveal a dynamic requirement of EMT in tumor metastasis: activation of EMT promotes local tumor invasion, intravasation, and extravasation of the systemic circulation, whereas reversion of EMT is essential to establish macrometastases in distant organs (1, 10). The "reversible" EMT model implies that EMT is unlikely to be regulated by permanent genetic and epigenetic changes in tumor cells; instead, EMT is dynamically controlled by various proinvasion signals from the tumor microenvironment (TME).

The TME is defined as the cellular and physical environment surrounding the primary tumor—including endothelial, inflammatory and immune cells, fibroblasts, extracellular matrix (ECM) components, and soluble factors. In this review, we discuss the most relevant and direct connections between TME signals and the EMT-inducing transcription factors in cancer. On the basis of the properties of the TME signals, we divide our discussion into four major categories: inflammatory signals, hypoxia, ECM components, and ECM mechanical properties (Fig. 1).

Inflammatory cytokines

An association between cancer development and inflammation has long been observed. During tumor progression, tumor cells recruit activated fibroblasts and immune cells that in turn secrete many cytokines to affect tumor development and metastasis (11). Interestingly, such cytokines have been shown to directly regulate the EMT program. Transforming growth factor- β (TGF β), abundantly secreted by cancer-associated fibroblasts, platelets, and tumor cells, is the best-characterized EMT inducer. TGF β has been

¹Department of Pharmacology, University of California, San Diego, La Jolla, California. ²Department of Pediatrics, University of California, San Diego, La Jolla, California.

Note: H.-Y. Jung and L. Fattet contributed equally to this article.

Corresponding Author: Jing Yang, Moores UCSD Cancer Center, University of California, San Diego, 3855 Health Sciences Drive, MC0819, La Jolla, CA 92093. Phone: 858-534-1994; Fax: 858-534-7390; E-mail: jingyang@ucsd.edu

doi: 10.1158/1078-0432.CCR-13-3173

©2014 American Association for Cancer Research.




Figure 1. Regulation of EMT transcription factors by tumor microenvironmental signals. TGFβ regulates upregulation of TWIST1, SNAIL1, and SNAIL2 via the SMAD signaling pathway. Drugs that inhibit TGFβ are AP 12009, GC1008, and LY573636, which are in clinical trials for advanced solid tumors. TNFα activates NF-κB to induce TWIST1, SNAIL2, and ZEB1/2 expression and TNFα/NF-κB activation also increases SNAIL1 protein stability. Therapeutic approaches to inhibit TGFβ signaling include TNFα antagonist (infliximab and etanercept) and NF-κB inhibitor (bortezomib), all of which have been assessed in phase II clinical trials for several cancer types. IL6 induces TWIST1 and SNAIL1 expression via JAK/STAT3 signaling and increases TWIST1 stability through CK2-dependent phosphorylation. An IL6 ligand-blocking antibody, CNTO 328, has been tested in phase I/II clinical trials with metastatic renal cell carcinoma. HIF1α induces TWIST1 and SNAIL1 expression and HIF1α either alone or in cooperation with TGFβ promotes SNAIL1 nuclear localization to stabilize SNAIL. Agents to inhibit HIF1α include EZN-2698, PX-478, and topotecan. Topotecan has been tested in phase I/II clinical trials in combination with conventional chemotherapy, and EZN-2698 and PX-478 are currently being tested in phase I clinical trials. Collagen I can promote SNAIL1 stability through binding to its receptor DDR2 and activating SRC/ERK2 pathway. HA binding to CD44 induces nuclear translocation of CD44 to directly induce lysyl-oxidase (LOX) expression, which in turn increases TWIST1 expression.

Downloaded from <http://aacrjournals.org/clinccancerres/article-pdf/21/5/963/20288961962> pdf by guest on 03 November 2024

shown to induce TWIST1 and SNAIL2 expression in prostate and non-small cell lung cancer (12, 13). TGF β can also induce SNAIL1 and SNAIL2 via IKK α and SMAD signaling in pancreatic cancer cells (14). Furthermore, Vincent and colleagues (15) showed that SNAIL-SMAD3/4 transcriptional repressor complex could promote TGF β -mediated EMT in breast cancer. Tumor necrosis factor- α (TNF α) is a crucial activator of the NF- κ B signaling pathway, and activated NF- κ B has been shown to induce multiple EMT transcription factors expression, including TWIST1, SNAIL2, and ZEB1/2 (16–18). Furthermore, Wu and colleagues (19) found that NF- κ B activation could stabilize SNAIL1 to further promote cell migration and invasion. The release of interleukins by immune cells, endothelial cells, and fibroblasts can also contribute to EMT. IL6 promotes EMT in head and neck cancer cells and correlates with increased TWIST1 and SNAIL1 expressions (20). Sullivan and colleagues (21) showed that an IL6-TWIST1 positive feedback loop induces EMT in breast cancer cells. Taken together, various inflammatory cytokines from TME can regulate the expression and/or protein stability of EMT transcription factors to activate EMT and tumor invasion.

Hypoxia

Hypoxia condition has been shown to select tumor cells to become more invasive and metastatic. Specifically, hypoxia can promote EMT via hypoxia-inducible factor-1 α (HIF1 α ; ref. 22). HIF1 α is found to increase SNAIL1 protein stability, leading to suppression of E-cadherin in ovarian carcinoma (23). Yang and colleagues (24) found that HIF1 α could induce TWIST1 expression by binding directly to the TWIST1 promoter. In addition, HIF1 α cooperates with inflammatory cytokines to promote EMT. For example, HIF1 α , together with TGF β , promotes SNAIL1 nuclear translocation to induce EMT through the suppression of estrogen receptor β in prostate carcinoma (25). Also, HIF1 α could enhance the expression of TWIST1 by upregulating TNF α , IL6, and TGF β in prostate cancer (26). Hypoxia, together with the Wnt/ β -catenin signaling, can also promote SNAIL1 stability by inhibiting GSK3 β (27). Taken together, HIF1 α , often in cooperation with additional TME factors, can induce EMT, suggesting a promising strategy to target hypoxic signaling for cancer therapeutics.

ECM components

ECM includes structural and nonstructural components that can activate cellular signaling through membrane-bound receptors such as integrins. The critical role of ECM in promoting EMT was already evident in the original experiments conducted by Greenburg and Hay (28). They showed that epithelial cells from embryonic and adult anterior lens cultured in three-dimensional collagen gels can elongate and migrate as individual cells. Indeed, Greenburg and Hay (28) concluded that "interactions with ECM may be a major factor in the ability of a cell to become mesenchymal."

Recently, Zhang and colleagues (29) unraveled a direct connection between ECM structural protein collagen-I and SNAIL1. They found that collagen-I binds to its receptor DDR2 and activates downstream SRC/ERK2 to stabilize SNAIL1 in breast tumors cells. SNAIL1 further upregulates MT1-MMP and collagen-I to promote tumor cell invasion. Another ECM structural component, fibronectin, partly through binding to integrin receptors, induces SNAIL1 expression in tumor cells. This study demonstrated that cooperation of fibronectin and TGF β was required to activate the downstream SRC and ERK/MAPK kinases and induce

EMT (30). Hyaluronan (HA) is a major component of ECM and signals through its membrane receptor CD44, which is overexpressed in many human cancers. HA binding to tumor cells was found to induce CD44 nuclear translocation and activate LOX expression, which in turn upregulates TWIST1 expression to promote breast cancer metastasis (31). Periostin, a nonstructural ECM component highly expressed in human tumors, could signal through integrins to increase cell survival and promote metastatic progression of colon cancer *in vivo* (32). Kim and colleagues (33) identified differential roles of periostin in EMT: it induces SNAIL1 expression in prostate cancer cells, whereas it inhibits TWIST1 expression in bladder cancer cells. These studies show that many ECM components are key regulators of EMT and tumor invasion.

ECM mechanical properties

During tumor progression, ECM is constantly remodeled by various cell types in the TME. Specifically, increasing matrix stiffness through LOX-mediated collagen cross-linking plays a critical role in tumor invasion and metastasis. Pioneer study by Paszek and colleagues (34) showed that increasing ECM stiffness induced a malignant phenotype, associated with activated FAK and ERK signaling. LOX-mediated ECM stiffening promoted tumor progression *in vivo* partially via an activated FAK signaling (35). Conversely, treatment with a LOX inhibitor reduced focal adhesions and PI3K signaling, demonstrating that LOX modulates tumor progression through ECM stiffening to drive focal adhesion assembly. Furthermore, ECM stiffening was required to cooperate with TGF β to induce EMT in human breast tumor cells (36), further strengthening the notion that mechanical properties of the tumor microenvironment are key factors regulating EMT and promoting tumor progression.

Clinical-Translational Advances

Accumulating evidence supports a critical role of EMT in many aspects of tumor development, including resistance to apoptosis and senescence, CSCs, and invasion and metastasis, thus suggesting that targeting this process could be a promising therapeutic approach. However, the core EMT transcription factors remain technically challenging to target. Instead, a number of preclinical studies suggest that inhibiting EMT-inducing TME signals could serve as alternative approaches to impinge upon the EMT program. Here, we summarize therapeutics in preclinical and clinical studies that target TME to prevent tumor progression (Table 1).

Inflammatory cytokines

Preclinical studies support the importance of inflammatory cytokines, including TNF α and IL6, in promoting EMT and tumor invasion. Several TNF α inhibitors have been tested in clinical trial in different types of cancers. For example, infliximab, a TNF α monoclonal blocking antibody, has been tested in phase II clinical trials in renal cell carcinoma and advanced cancers (37, 38). These studies suggested that TNF α inhibitor was effective to suppress the levels of IL6 and CCL2 in patients and improved progress-free survival. Two clinical studies examined the therapeutic effects of etanercept, a TNF α antagonist, in recurrent ovarian cancer and metastatic breast cancer. Etanercept is well tolerated in patients and significantly improved progress-free survival with consistent decrease in CCL and IL levels (39, 40). Because NF- κ B is the essential downstream activator of the TNF α signaling, several clinical trials tested whether inhibition of NF- κ B signaling could

Table 1. Clinical studies of drugs that target tumor microenvironment

Target	Drug	Types of drug	Cancer types	Clinical studies	Response Median/mo ^a	PFS Median/mo ^a	OS Median/mo ^a	Reference
TNF α	Infliximab	Monoclonal antibody	Renal cell carcinoma	Phase II	Study 1: 32% PR+SD (7.7) Study 2: 61% SD (6.2)	Study 1: 3.1 Study 2: 4.1	Study 1: 10 Study 2: 13.1	37 38
			Ovarian, renal, cervical cancer, endometrial stromal cell sarcoma, metastatic melanoma, and metastatic colon cancer	Phase I	Study 2: 61% SD (6.2) 6.2% SD (3.9)			
	Etanercept (Enbrel)	Monoclonal antibody	Metastatic breast cancer	Phase II	6.25% SD (4.1)			39
NF- κ B	Bortezomib	Proteasome inhibitor	Recurrent ovarian cancer	Phase I	33.3% SD (6.25)			40
			Unresectable/metastatic gastric and gastroesophageal junction adenocarcinoma	Phase II	6.25% SD (3.3)	1.28	5.08	41
			Recurrent and/or metastatic head and neck squamous cell carcinoma	Phase II	53% PR+SD (3.0)	3.0	9.4	42
IL6	Siltuximab (CNTO-328)	Monoclonal antibody	Metastatic renal cell cancer	Phase I/II	Part 2: 54% PR+SD (7.6)	3.4	5	43
TGF β	AP-12009	Antisense oligonucleotide	Recurrent malignant glioma	Phase I/II	29.1% SD (6)		11	44
	GC-1008	Monoclonal antibody	Metastatic melanoma and renal cell carcinoma	Phase I/II (ongoing)				45
	LY-573636 (tasisulam sodium)	Small-molecule inhibitor	Unresectable/metastatic non-small cell lung cancer	Phase II	43.8% SD (4.21)	2.69	8.48	46
			Unresectable/metastatic soft tissue sarcoma	Phase II	46.0% PR+SD (4.44)	2.64	8.71	47
			Unresectable/metastatic melanoma	Phase II	47.1% CR+PR+SD (6.6)	2.6	9.6	48
HIF1 α	EZN-2698	Antisense oligonucleotide	Advanced solid tumors and metastatic renal cell carcinoma	Phase I (ongoing)				49
	PX-478	Small-molecule inhibitor	Advanced metastatic cancer	Phase I (ongoing)				49
	Topotecan	Small-molecule inhibitor	Advanced, refractory non-small cell lung cancer	Phase I/II	69.1% PR+SD (5.1)	5.2	11.5	50
Tenascin C	²¹¹ At-ch81C6	Radioactive particles	GBM, anaplastic astrocytoma (AA), anaplastic oligodendroglioma (AO)	Pilot	GBM: SD (13.5)		GBM: 37.5 Non-GBM: 60	53
FAK	PF-00562271	Small-molecule inhibitor	Advanced solid tumors	Phase I	AA: SD (13) AO: SD (29) 34% SD (1.5) 17% SD (9)			59

Abbreviations: CR, complete response; OS, overall survival; PFS, progress-free survival; PR, partial response; SD, stable disease.

^aMedian/mo: median duration/months.

suppress tumor progression and metastasis. Bortezomib, a proteasome inhibitor that suppresses NF- κ B activation, was tested in phase II clinical studies with metastatic gastric adenocarcinoma, and recurrent and metastatic head and neck squamous cell carcinoma (41, 42). Although bortezomib alone showed poor response in patients, combination therapy with docetaxel or targeted inhibition of other oncogenic pathways are currently under way in solid tumors. Finally, various blocking antibodies against cytokines have been used in various clinical studies. CNTO-328, an IL6 ligand-blocking antibody, was tested in phase I/II clinical trials for the treatment of metastatic renal cell carcinoma. This study showed that CNTO-328 could increase patient survival and more than 50% of progressive metastatic renal cell carcinoma patients presented stable diseases upon treatment (43). Together, these clinical trials in progress could bring a number of promising anti-inflammatory cytokine agents to the forefront of antimetastasis therapeutics.

The TGF β signaling is extensively targeted to block tumor progression and metastasis, and various approaches have been taken to inhibit the TGF β signaling. AP-12009, an antisense oligonucleotide against TGF β II, was tested in patients with high-grade glioma and significantly improved survival compared with standard chemotherapy treatment (44). Furthermore, TGF β -neutralizing antibody GC-1008 showed promises in phase I trial for metastatic melanoma and renal cell carcinoma (45). Small-molecule inhibitor, LY-573636, used in phase II clinical studies in patients with metastatic NSCLC, soft tissue sarcoma, and melanoma, has also shown modest activity as a second/third-line therapy (46–48). These studies showed that inhibiting TGF β signaling pathway is safe, well tolerated in patients and could provide promising new therapeutics against tumor invasion.

Hypoxia

Several HIF1 α inhibitors have also shown remarkable antitumor activities in a variety of preclinical and clinical trials. EZN-2698, an antisense oligonucleotide against of HIF1 α , is being tested in a phase I clinical trial with advanced solid tumors (49). Another HIF1 α inhibitor, PX-478, which inhibit HIF1 α expression, is currently tested in phase I clinical trials in patients with advanced metastatic cancer (49). Several novel compounds have also been identified in a high-throughput screen using a cell-based reporter of HIF1 α transcriptional activity. One such compound topotecan has been tested in phase I/II clinical trials with conventional chemotherapies such as cisplatin or bevacizumab in patients with advanced lung cancer. Clinical results indicate that combination treatment is well tolerated and worthy of further clinical investigation (50), thus making them promising agents against tumor metastasis.

ECM components

Disruption of tumor ECM integrity has shown promising results in halting tumor metastasis in preclinical studies. Methylumbelliferone, a HA synthesis inhibitor, was effective in preventing bone metastasis of lung cancer *in vivo* (51). Neutralizing antibody directed against periostin resulted in 40% inhibition of tumor growth ($P < 0.001$), 80% inhibition of lung metastasis ($P < 0.001$), and significant increase in survival ($P < 0.05$) using mouse breast tumor xenografts (52).

Because cells that have undergone EMT secrete many unique ECM components, these ECM molecules have also been used for targeting drug delivery to tumors. For example, a promising

approach has been used in clinical trials for patients with glioblastoma multiforme (GBM), linking anti-Tenascin C antibody to radioactive particles to specifically target tumor cells. Result showed minimal toxicity associated with a promising antitumor benefit and encouraging overall outcomes (53). Recently, engineered HA-based conjugates have emerged as a promising strategy to efficiently target tumors with drugs exerting poor solubility and strong side effects, such as paclitaxel (54). These strategies take advantage of unique EMT-associated TME components to achieve targeted delivery of traditional chemotherapeutics, thus presenting a new anticancer therapeutic strategy.

ECM mechanical properties

In patients, the presence of fibrotic foci in breast tumors is a prognostic marker of distant metastasis and correlates with poor survival (55). In addition, LOX is essential for hypoxia-induced breast cancer metastasis and its expression in patients is correlated to a poor outcome (56). Finally, a recent study shows that LOX is critical to establish a permissive microenvironment within fibrotic tissues, characterized by increased EMT, to favor the colonization of metastasizing tumor cells (57). Thus, anti-LOX strategies could suppress metastatic progression of the disease, not only by targeting the TME of the premetastatic niche, but also by targeting tumor cells themselves, as shown by the direct effect of LOX inhibition in attenuating FAK-dependent breast cancer cell invasion in a preclinical study (58). Therapeutic inhibition of FAK, recently validated in a phase I study, may also be a promising approach to prevent the effect of TME stiffness on metastatic progression of several types of cancer. Indeed the use of pharmacologic inhibitor PF-00562271 in patients with advanced solid tumors unresponsive to existing therapies showed a significant stabilization of the disease, thus supporting FAK as a potential therapeutic target (59).

Conclusion and Discussion

As discussed, a number of inhibitors targeting TME are being tested in preclinical and clinical trials and well-tolerated in patients and several showed promising results. Because these TME signals regulate various signaling pathways, the impacts of these inhibitors on tumor progression are likely beyond the EMT program. Given the critical role of EMT in multiple steps of tumor progression, targeting the EMT-inducing TME signals is indeed worth pursuing to combat metastatic cancers.

However, there are also a number of issues to be resolved to better decide how to effectively affect tumor progression by targeting the EMT program. First, current clinical trials largely aim to shrink established metastases, in which the EMT program may not be involved. Instead, metastasis prevention trials in patients with cancer with high metastasis risk would be the appropriate setting to test the effect of EMT inhibition on metastasis occurrence. Second, recent studies demonstrated the dynamic involvement of EMT in tumor metastasis: activation of EMT promotes tumor dissemination and reversion of EMT is essential for outgrowth of macrometastases. Therefore, EMT inhibitor alone could be counter-productive in preventing distant metastases if patients already have disseminated tumor cells in distant organs. In these cases, combining therapies targeting TME signals with traditional chemotherapy and targeted therapies to simultaneously inhibit EMT and cell proliferation could be a more powerful approach to eradicate both migrating as well as proliferating tumor cells, thus halting tumor progression.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: H.-Y. Jung, L. Fattet, J. Yang

Writing, review, and/or revision of the manuscript: H.-Y. Jung, L. Fattet, J. Yang

Acknowledgments

The authors apologize to the many researchers in this field whose work they were unable to cite due to space restrictions.

Grant Support

L. Fattet is supported by a postdoctoral fellowship from the Fondation pour la Recherche Médicale (SPE20130326547). J. Yang is supported by the NCI of the NIH under award number 1R01CA168689, American Cancer Society grant RSG-09-282-01-CSM, the Hartwell Foundation, and the U.S. Department of Defense Breast Cancer Program under award number W81XWH-13-1-0132.

Received May 15, 2014; revised July 10, 2014; accepted July 18, 2014; published OnlineFirst August 8, 2014.

References

- De Craene B, Bex G. Regulatory networks defining EMT during cancer initiation and progression. *Nat Rev Cancer* 2013;13:97–110.
- Sánchez-Tilló E, Liu Y, de Barrios O, Siles L, Fanlo L, Cuatrecasas M, et al. EMT-activating transcription factors in cancer: beyond EMT and tumor invasiveness. *Cell Mol Life Sci* 2012;69:3429–56.
- Thiery JP, Acloque H, Huang RYJ, Nieto MA. Epithelial-mesenchymal transitions in development and disease. *Cell* 2009;139:871–90.
- Valdés F, Alvarez AM, Locascio A, Vega S, Herrera B, Fernández M, et al. The epithelial mesenchymal transition confers resistance to the apoptotic effects of transforming growth factor Beta in fetal rat hepatocytes. *Mol Cancer Res* 2002;1:68–78.
- Derksen PWB, Liu X, Saridin F, van der Gulden H, Zevenhoven J, Evers B, et al. Somatic inactivation of E-cadherin and p53 in mice leads to metastatic lobular mammary carcinoma through induction of anoikis resistance and angiogenesis. *Cancer Cell* 2006;10:437–49.
- Ansieau S, Bastid J, Doreau A, Morel A-P, Bouchet BP, Thomas C, et al. Induction of EMT by TWIST proteins as a collateral effect of tumor-promoting inactivation of premature senescence. *Cancer Cell* 2008;14:79–89.
- Browne G, Sayan AE, Tulchinsky E. ZEB proteins link cell motility with cell cycle control and cell survival in cancer. *Cell Cycle* 2010;9:886–91.
- Mani SA, Guo W, Liao M-J, Eaton EN, Ayyanan A, Zhou AY, et al. The epithelial-mesenchymal transition generates cells with properties of stem cells. *Cell* 2008;133:704–15.
- Morel A-P, Lièvre M, Thomas C, Hinkal G, Ansieau S, Puisieux A. Generation of breast cancer stem cells through epithelial-mesenchymal transition. *PLoS ONE* 2008;3:e2888.
- Tsai JH, Donaher JL, Murphy DA, Chau S, Yang J. Spatiotemporal regulation of epithelial-mesenchymal transition is essential for squamous cell carcinoma metastasis. *Cancer Cell* 2012;22:725–36.
- Rakoff-Nahoum S. why cancer and inflammation? *Yale J Biol Med* 2006;79:123–30.
- Shiota M, Zardan A, Takeuchi A, Kumano M, Beraldi E, Naito S, et al. Clusterin mediates TGF β -induced epithelial-mesenchymal transition and metastasis via TWIST1 in prostate cancer cells. *Cancer Res* 2012;72:5261–72.
- Slabáková E, Pernicová Z, Slavíčková E, Stars'ířhová A, Kozubík A, Souček K. TGF β 1-induced EMT of non-transformed prostate hyperplasia cells is characterized by early induction of SNAIL/Slug. *Prostate* 2011;71:1332–43.
- Brandl M, Seidler B, Haller F, Adamski J, Schmid RM, Saur D, et al. IKK α controls canonical TGF β -SMAD signaling to regulate genes expressing SNAIL and SLUG during EMT in panc1 cells. *J Cell Sci* 2010;123:4231–9.
- Vincent T, Neve EPA, Johnson JR, Kukalev A, Rojo F, Albanell J, et al. A SNAIL1-SMAD3/4 transcriptional repressor complex promotes TGF β mediated epithelial-mesenchymal transition. *Nat Cell Biol* 2009;11:943–50.
- Li CW, Xia W, Huo L, Lim SO, Wu Y, Hsu JL, et al. Epithelial-mesenchymal transition induced by TNF α requires NF κ B-mediated transcriptional upregulation of TWIST1. *Cancer Res* 2012;72:1290–300.
- Storci G, Sansone P, Mari S, D'Uva G, Tavolari S, Guarnieri T, et al. TNF α up-regulates SLUG via the NF κ B/HIF1 α axis, which imparts breast cancer cells with a stem cell-like phenotype. *J Cell Physiol* 2010;225:682–91.
- Chua HL, Bhat-Nakshatri P, Clare SE, Morimiya A, Badve S, Nakshatri H. NF κ B represses E-cadherin expression and enhances epithelial to mesenchymal transition of mammary epithelial cells: potential involvement of ZEB-1 and ZEB-2. *Oncogene* 2006;26:711–24.
- Wu Y, Deng J, Rychahou PG, Qiu S, Evers BM, Zhou BP. Stabilization of SNAIL by NF κ B is required for inflammation-induced cell migration and invasion. *Cancer Cell* 2009;15:416–28.
- Su Y-W, Xie T-X, Sano D, Myers JN. IL6 stabilizes TWIST and enhances tumor cell motility in head and neck cancer cells through activation of Casein kinase 2. *PLoS ONE* 2011;6:e19412.
- Sullivan NJ, Sasser AK, Axel AE, Vesuna F, Raman V, Ramirez N, et al. Interleukin-6 induces an epithelial-mesenchymal transition phenotype in human breast cancer cells. *Oncogene* 2009;28:2940–7.
- Vooijs M, Gort E, Groot A, Wall der E, van Diest P. Hypoxic regulation of metastasis via hypoxia-inducible factors. *Curr Mol Med* 2008;8:60–7.
- Imai T, Horiuchi A, Wang C, Oka K, Ohira S, Nikaido T, et al. Hypoxia attenuates the expression of E-Cadherin via up-regulation of SNAIL in ovarian carcinoma cells. *Am J Pathol* 2003;163:1437–47.
- Yang M-H, Wu M-Z, Chiou S-H, Chen P-M, Chang S-Y, Liu C-J, et al. Direct regulation of TWIST by HIF1 α promotes metastasis. *Nat Cell Biol* 2008;10:295–305.
- Mak P, Leav I, Pursell B, Bae D, Yang X, Taglienti CA, et al. ER β impedes prostate cancer EMT by destabilizing HIF1 α and inhibiting VEGF-mediated SNAIL nuclear localization: implications for Gleason grading. *Cancer Cell* 2010;17:319–32.
- Kim H-J, Park J-W, Cho Y-S, Cho C-H, Kim J-S, Shin H-W, et al. Pathogenic role of HIF1 α in prostate hyperplasia in the presence of chronic inflammation. *Biochim Biophys Acta* 2013;1832:183–94.
- Cannito S, Novo E, Compagnone A, Valfrè di Bonzo L, Busletta C, Zamara E, et al. Redox mechanisms switch on hypoxia-dependent epithelial-mesenchymal transition in cancer cells. *Carcinogenesis* 2008;29:2267–78.
- Greenburg G, Hay ED. Epithelia suspended in collagen gels can lose polarity and express characteristics of migrating mesenchymal cells. *J Cell Biol* 1982;95:333–9.
- Zhang K, Corsa CA, Ponik SM, Prior JL, Piwnicka-Worms D, Eliceiri KW, et al. The collagen receptor discoidin domain receptor 2 stabilizes SNAIL1 to facilitate breast cancer metastasis. *Nat Cell Biol* 2013;15:677–87.
- Park JH, Schwarzbauer JE. Mammary epithelial cell interactions with fibronectin stimulate epithelial-mesenchymal transition. *Oncogene* 2013;33:1649–57.
- El-Haibi CP, Bell GW, Zhang J, Collmann AY, Wood D, Scherber CM, et al. Critical role for lysyl oxidase in mesenchymal stem cell-driven breast cancer malignancy. *Proc Natl Acad Sci U S A* 2012;109:17460–5.
- Bao S, Ouyang G, Bai X, Huang Z, Ma C, Liu M, et al. Periostin potently promotes metastatic growth of colon cancer by augmenting cell survival via the Akt/PKB pathway. *Cancer Cell* 2004;5:329–39.
- Kim CJ, Sakamoto K, Tambe Y, Inoue H. Opposite regulation of epithelial-to-mesenchymal transition and cell invasiveness by periostin between prostate and bladder cancer cells. *Int J Oncol* 2011;38:1759–66.
- Paszek MJ, Zahir N, Johnson KR, Lakins JN, Rozenberg GI, Gefen A, et al. Tensional homeostasis and the malignant phenotype. *Cancer Cell* 2005;8:241–54.
- Levental KR, Yu H, Kass L, Lakins JN, Egeblad M, Erler JT, et al. Matrix crosslinking forces tumor progression by enhancing integrin signaling. *Cell* 2009;139:891–906.
- Leight JL, Wozniak MA, Chen S, Lynch ML, Chen CS. Matrix rigidity regulates a switch between TGF β 1-induced apoptosis and epithelial-mesenchymal transition. *Mol Biol Cell* 2012;23:781–91.

37. Harrison ML, Obermueller E, Maisey NR, Hoare S, Edmonds K, Li NF, et al. Tumor Necrosis Factor α as a new target for renal cell carcinoma: two sequential phase II trials of Infliximab at standard and high dose. *J Clin Oncol* 2007;25:4542–9.
38. Brown ER, Charles KA, Hoare SA, Rye RL, Jodrell DI, Aird RE, et al. A clinical study assessing the tolerability and biological effects of infliximab, a TNF α inhibitor, in patients with advanced cancer. *Ann Oncol* 2008;19:1340–6.
39. Madhusudan S. A phase II study of Etanercept (Enbrel), a tumor necrosis factor α inhibitor in patients with metastatic breast cancer. *Clin Cancer Res* 2004;10:6528–34.
40. Madhusudan S. Study of etanercept, a tumor necrosis factor-alpha inhibitor, in recurrent ovarian cancer. *J Clin Oncol* 2005;23:5950–9.
41. Shah MA, Power DG, Kindler HL, Holen KD, Kemeny MM, Ilson DH, et al. A multicenter, phase II study of Bortezomib (PS-341) in patients with unresectable or metastatic gastric and gastroesophageal junction adenocarcinoma. *Invest New Drugs* 2010;29:1475–81.
42. Chung CH, Aulino J, Muldowney NJ, Hatakeyama H, Baumann J, Burkey B, et al. Nuclear factor-kappa B pathway and response in a phase II trial of bortezomib and docetaxel in patients with recurrent and/or metastatic head and neck squamous cell carcinoma. *Ann Oncol* 2010;21:864–70.
43. Rossi J-F, Négrier S, James ND, Kocak I, Hawkins R, Davis H, et al. A phase I/II study of siltuximab (CNTO 328), an anti-interleukin-6 monoclonal antibody, in metastatic renal cell cancer. *Br J Cancer* 2010;103:1154–62.
44. Hau P, Jachimczak P, Schlingensiepen R, Schulmeyer F, Jauch T, Steinbrecher A, et al. Inhibition of TGF β 2 with AP 12009 in recurrent malignant gliomas: from preclinical to phase I/II studies. *Oligonucleotides* 2007;17:201–12.
45. Nagaraj NS, Datta PK. Targeting the transforming growth factor- β signaling pathway in human cancer. *Expert Opin Investig Drugs* 2010;19:77–91.
46. Scagliotti CV, Ilaria R Jr, Novello S, Pawel von J, Fischer JR, Ermisch S, et al. Tasisulam sodium (LY573636 Sodium) as third-line treatment in patients with unresectable, metastatic non-small-cell lung cancer. *J Thorac Oncol* 2012;7:1053–7.
47. Ryan CW, Matias C, Agulnik M, Lopez-Pousa A, Williams C, Alwis DP, et al. A phase II study of tasisulam sodium (LY573636 sodium) as second-line or third-line treatment for patients with unresectable or metastatic soft tissue sarcoma. *Invest New Drugs* 2012;31:145–51.
48. Kirkwood JM, Gonzalez R, Reintgen D, Clingan PR, McWilliams RR, de Alwis DP, et al. A phase 2 study of tasisulam sodium (LY573636 sodium) as second-line treatment for patients with unresectable or metastatic melanoma. *Cancer* 2011;117:4732–9.
49. Onnis B, Rapisarda A, Melillo G. Development of HIF1 inhibitors for cancer therapy. *J Cell Mol Med* 2009;13:2780–6.
50. Powell SF, Beitinjaneh A, Tessema M, Bliss RL, Kratzke RA, Leach J, et al. Phase II study of topotecan and bevacizumab in advanced, refractory non-small-cell lung cancer. *Clin Lung Cancer* 2013;14:495–501.
51. Futamura N, Urakawa H, Arai E, Kozawa E, Ishiguro N, Nishida Y. Hyaluronan synthesis inhibitor supplements the inhibitory effects of zoledronic acid on bone metastasis of lung cancer. *Clin Exp Metastasis* 2013;30:595–606.
52. Morishita R. Role of periostin in cancer progression and metastasis: inhibition of breast cancer progression and metastasis by anti-periostin antibody in a murine model. *Int J Mol Med* 2011;28:181–6.
53. Zalutsky MR, Reardon DA, Akabani G, Coleman RE, Friedman AH, Friedman HS, et al. Clinical experience with α -particle emitting 211At: treatment of recurrent brain tumor patients with 211At-labeled chimeric antitenascin monoclonal antibody 81C6. *J Nucl Med* 2007;49:30–8.
54. Arpicco S, Milla P, Stella B, Dosio F. Hyaluronic acid conjugates as vectors for the active targeting of drugs, genes and nanocomposites in cancer treatment. *Molecules* 2014;19:3193–230.
55. Colpaert C, Vermeulen P, VanMarck E, Dirix L. The presence of a fibrotic focus is an independent predictor of early metastasis in lymph node-negative breast cancer patients. *Am J Surg Pathol* 2001;25:1557.
56. Erler JT, Bennewith KL, Nicolau M, Dornhöfer N, Kong C, Le Q-T, et al. Lysyl oxidase is essential for hypoxia-induced metastasis. *Nature* 2006;440:1222–6.
57. Cox TR, Bird D, Baker A-M, Barker HE, Ho MWY, Lang G, et al. LOX-mediated collagen crosslinking is responsible for fibrosis-enhanced metastasis. *Cancer Res* 2013;73:1721–32.
58. Chen L-C, Tu S-H, Huang C-S, Chen C-S, Ho C-T, Lin H-W, et al. Human breast cancer cell metastasis is attenuated by lysyl oxidase inhibitors through down-regulation of focal adhesion kinase and the paxillin-signaling pathway. *Breast Cancer Res Treat* 2012;134:989–1004.
59. Infante JR, Camidge DR, Mileskin LR, Chen EX, Hicks RJ, Rischin D, et al. Safety, pharmacokinetic, and pharmacodynamic phase I dose-escalation trial of PF-00562271, an inhibitor of focal adhesion kinase, in advanced solid tumors. *J Clin Oncol* 2012;30:1527–33.