

---

# 45 Field Studies of Social Cognition in Spotted Hyenas

Kay E. Holekamp and Anne L. Engh

Considered in relation to body size, the brains of primates are relatively large and complex compared with those of other animals, including most nonprimate mammals (Macphail 1982; Harvey and Krebs 1990). In particular, the primate neocortex is very large in relation to the rest of the brain (Barton and Dunbar 1997). Furthermore, primates appear to be endowed with cognitive abilities that are superior to, and qualitatively different from, those observed in most other mammals (reviewed in Byrne and Whiten 1988; Harcourt and de Waal 1992; Tomasello and Call 1997).

Two different types of selection pressures have been hypothesized to favor the evolution of large brains and great intelligence (as defined by Kamil 1987) in primates. The first hypothesis suggests that intelligence has been favored in primates by selection pressures associated with complexity in the physical environment, particularly that confronted when navigating through a three-dimensional arboreal world (e.g., Povinelli and Preuss 1995) or when finding and obtaining food (e.g., Milton 1981). The second hypothesis suggests instead that the key selection pressures have been imposed by complexity associated with the labile social behavior of conspecific group members (Byrne and Whiten 1988). Predictions of this social complexity hypothesis have now been confirmed in a number of Old World primate species, suggesting that the evolution of intelligence has been more strongly influenced by social pressures than by nonsocial aspects of the environment (reviewed in Byrne 1994; Tomasello and Call 1997). Unfortunately the generality of this hypothesis is severely limited by the current dearth of information about social cognition in animals other than primates (Harcourt and de Waal 1992). In fact, most work in this area has focused exclusively on cercopithecine primates and great apes. The social complexity hypothesis, however, predicts that nonprimate animals

that share with primates most salient features of their social life and resource distribution should possess many of the same features of social intelligence found in monkeys and apes.

Like most primates, many mammalian carnivores live in permanent, complex social groups that contain both males and females from multiple, overlapping generations. Gregarious carnivores also engage in a variety of behaviors, such as cooperative hunts of large vertebrate prey, which have prompted many observers to claim that these predators must possess extraordinary intellectual powers (e.g., Guggisberg 1962). The size of the carnivore neocortex is positively correlated with group size (Dunbar and Bever 1998). However, the cognitive abilities of carnivores have seldom been the subject of systematic study, and they are currently very poorly understood (e.g., Byrne 1994).

In our own research we are examining the cognitive mechanisms underlying social behavior and communication in one gregarious carnivore, the spotted hyena (*Crocuta crocuta*). Our ultimate objective is to determine whether hyenas exhibit some of the same cognitive abilities that are observed in primates. Evidence for the existence of shared cognitive abilities would suggest convergent evolution in these two distantly related taxa and would strongly support the social complexity hypothesis. In contrast, the failure to obtain such evidence would suggest that the social complexity hypothesis should be either rejected or revised. We use three different methods to investigate the cognitive abilities of gregarious carnivores: (1) comparative analysis based on literature review, (2) field experiments in the natural habitat, and (3) controlled observations of free-living hyenas. Here we summarize our work to date using each of these methods, although we hope readers will bear in mind the fact that much of it is still in progress.

---

### Comparative Analysis of Group Travel

Here we sought to understand whether coordinated movements by gregarious carnivores reveal the operation of complex mental abilities. Like many primates (e.g., Boinski and Garber 2000), gregarious carnivores frequently travel with other members of their social groups. Group hunts by lions, spotted hyenas, and other carnivores often appear to involve intelligent coordination and division of labor among hunters (e.g., Guggisberg 1962; Stander 1992a,b). Studies of carnivore predatory behavior reveal that their group hunts represent more complexly organized phenomena than mere opportunistic grabs at prey. However, although myriad observers have claimed that the group hunting activity of large carnivores requires the operation of humanlike mental processes, coordinated hunting behavior by lions, hyenas, and social canids can in fact be most parsimoniously explained by the operation of a few simple mental rules of thumb, such as “Move wherever you need to in order to keep the selected prey animal between you and another hunter” (Holekamp et al. 2000). Currently there is no evidence that gregarious carnivores use mental algorithms more complex than simple rules of thumb to surround and capture prey. Disproving the rules of thumb hypothesis will require experimental evidence, not only that individual carnivores monitor both their prey and their fellow hunters (e.g., Stander 1992b), but also that they accurately anticipate the behavior of the latter based on knowledge of their goals.

Another form of group travel that reveals the mental operations of gregarious carnivores takes place when young cubs or pups are moved from one den site to another (Holekamp et al. 2000). Whereas group hunts make carnivores look very smart, den moves make them appear very stupid. When gregarious carnivores move infants ( $N$ ) to a new den, they always make at least one extra trip ( $N + 1$ ) to the old den, suggesting that instead of being able to count, during den moves

carnivores operate according to a rule of thumb that says “Revisit the old den until you find no more of your infants there.” Thus the mental processes required to initiate and maintain seemingly complex movements need not be particularly sophisticated in carnivores. Interestingly, it appears that much coordinated group travel in primates, including cooperative hunting behavior by chimpanzees (e.g., Boesch and Boesch 1989), might also be explained effectively by these same simple rules of thumb.

---

### Field Experiments and Controlled Observations of Spotted Hyenas

Many primates demonstrate the ability to form mental representations of tertiary, or third-party, relationships among conspecific group members (Tomasello and Call 1997). These involve interactions and relationships in which the observer is not directly involved. For example, adult male hamadryas baboons make adaptive decisions about whether to challenge a rival male based on their assessment of the strength of the social bond between the rival and his female associates (Bacchman and Kummer 1980). Tomasello and Call (1997) hypothesize that an ability to understand third-party relationships is unique to primates, and furthermore, that this distinguishes their mental abilities from those of all other animals. Indeed, the ability to recognize third-party relationships has not been documented in non-primate mammals, even in species living in complex, stable societies such as elephants (Moss 1988), dolphins (Connor et al. 1992), or lions (Packer 1994). However, few published studies of these other gregarious mammals have focused on the cognitive abilities of these animals, so it remains unclear whether the absence of evidence that any of these nonprimate mammals can recognize tertiary relationships actually means that this ability is also absent. Our own field work has attempted to determine whether a primate-

like ability to recognize tertiary relationships is exhibited by the spotted hyena, a carnivore whose social life is remarkably similar in most respects to that of many Old World primates.

### **The Social Lives of Hyenas and Old World Monkeys**

Both spotted hyenas and Old World monkeys usually live in permanent social groups that commonly contain multiple adult males and multiple matrilineal lines of adult female kin with offspring. Males leave their natal groups, whereas females are usually philopatric (Cheney and Seyfarth 1983; Smale et al. 1997). Adults of both taxa can be ranked in a linear dominance hierarchy based on outcomes of agonistic interactions, and priority of access to resources varies with social rank (Tilson and Hamilton 1984; Andelman 1985). In both taxa, members of the same matriline occupy adjacent rank positions in the group's hierarchy, and female dominance relations are extremely stable across a variety of contexts and over extended periods. Juvenile hyenas of both sexes acquire ranks immediately below those of their mothers (Holekamp and Smale 1991), and they do this via the same associative learning mechanisms as those documented in cercopithecine primates (Horrocks and Hunte 1983; Engh et al. 2000). Thus social status is often not determined by size, strength, or fighting ability in either of these taxa, as it is in many mammals (e.g., Clutton-Brock et al. 1982). Instead, juvenile monkeys and hyenas both learn during early life that they can dominate individuals ranked lower than their mothers. In both monkeys and hyenas, kin associate more closely than do non-kin, and individuals direct affiliative behavior more frequently toward kin than non-kin (Seyfarth 1980; East et al. 1993; Holekamp et al. 1997).

In both taxa, high-ranking animals are preferred over lower-ranking individuals as social companions (Seyfarth, 1980; Holekamp et al. 1997). Furthermore, the patterns of greeting behavior in *Crocuta* follow primate patterns of

social grooming (East et al. 1993), in which individuals prefer to direct affiliative behavior toward high-ranking non-kin (Seyfarth and Cheney 1984). This suggests that hyenas, like monkeys, may recognize that some group members are more valuable social partners than others. Finally, triadic and more complex interactions (e.g., coalitions) appear to play important roles in both maintenance and reversals of social rank in free-living *Crocuta* (Zabel et al. 1992; Smale et al. 1993), as they do in many cercopithecine primates (Walters 1980; Datta 1986; Chapais 1992).

### **Playback Experiments on Vocal Recognition**

The similarities between the social lives of spotted hyenas and Old World primates present an ideal opportunity for comparative analysis of their social knowledge. In our first attack on this problem (Holekamp et al. 1999), we conducted a series of playback experiments with hyenas in the field and compared the hyenas' behavioral responses during these experiments with those observed earlier in vervet monkeys (Cheney and Seyfarth 1990) and in other cercopithecine primates (e.g., Gouzoules et al. 1995).

In the vervet experiments (Cheney and Seyfarth 1980), the distress scream of a juvenile was played through a hidden loudspeaker to groups of females that contained the juvenile's mother as well as other adult breeding females (controls). The mothers responded more strongly to screams than the control females, and the controls often responded by looking at the mother without any apparent cues from the mother herself. Thus vervets behaved as though they recognized the close associative relationships existing among group members unrelated to themselves (Cheney and Seyfarth 1980, 1990).

We used playback experiments to determine whether hyenas are capable of identifying individual conspecifics on the basis of their long-distance "whoop" vocalizations and whether they recognize tertiary relationships among non-

kin members of their clan, as occurs in vervets. Although the results obtained during our playback experiments showed clearly that hyenas can recognize the whoops of specific individuals, they yielded no evidence that hyenas can recognize third-party relationships (Holekamp et al. 1999). In contrast to vervets, after the playback, the control females in our experiments were no more likely to look at the mother of the whooping cub than at other control females. Although this result suggests that spotted hyenas do not recognize third-party relationships, it remains possible that they do recognize them, but that their orientation responses during playback experiments are poor indicators of this ability. If this alternative hypothesis is correct, then other social situations, in which actions based on social knowledge will unambiguously improve fitness outcomes for the actor, should reveal that hyenas can indeed recognize third-party relationships. Therefore we recently turned our attention to these other types of social situations.

### *Coalition Formation*

Coalitions play an important role in the acquisition and maintenance of social rank in spotted hyenas (Zabel et al. 1992; Smale et al. 1993; Engh et al. 2000). When aggression between two hyenas escalates, one or more others may join the skirmish by forming a coalition with the attacker against the target individual. Typically, animals joining to form coalitions are all dominant to the victim. Thus when attempting to displace a larger subordinate animal from food, a hyena might benefit, for example, by delaying its attack until the arrival of a potential coalitionary ally who is higher ranking than the target animal.

We are currently analyzing changes in rates of aggression directed against subordinates after the arrival of hyenas of high and intermediate rank in order to learn more about hyenas' abilities to recognize tertiary relationships (Engh et al. in preparation). If hyenas increase their rates of aggression only after higher-ranking hyenas arrive on the scene, then they may be following a

simple rule of thumb, such as "Only attack a larger subordinate when another individual is present who is higher ranking than yourself." Alternatively, if the attack rate also increases following the arrival of an individual who is dominant to the victim but subordinate to the attacker, then the attacking hyena must recognize the relative ranks of the other two individuals. In the latter case, the hyenas would be indicating to us that they can indeed recognize tertiary relationships. Although we are still in the early stages of this study, the preliminary results suggest that hyenas increase their rates of aggression only when animals higher ranking than themselves arrive, indicating once again that they are probably using simple rules of thumb to make decisions, rather than knowledge of complex relationships.

### *Reconciliation and Redirected Aggression*

Affiliative gestures functioning to repair social relationships damaged during a fight are called "reconciliation behaviors" (de Waal 1993). Reconciliation occurs in many primates during friendly reunions between former opponents shortly after aggressive conflicts (reviewed by Aurie and de Waal 2000). Similarly, spotted hyenas appear to reconcile after 10 to 15 percent of fights, often by the loser lifting its leg in a friendly "greeting" gesture directed at its former opponent (Hofer and East 2000; Wahaj et al. 2001). After fights, Old World primates are known to reconcile, not only with their former opponents, but also with the kin of former opponents (e.g., Cheney and Seyfarth 1989), indicating that the conciliatory monkeys recognize those tertiary relationships. In contrast, we rarely observe hyenas reconciling with any animals but their former opponents. This suggests that even though the ability to recognize tertiary relationships under these circumstances might well enhance their fitness, hyenas do not behave as though they possess this ability.

In many primates, an animal that has been involved in a fight will redirect aggression and

threaten a third, previously uninvolved individual (e.g., Cheney and Seyfarth 1989). Often such redirected aggression is directed toward a close relative of the prior opponent. Thus patterns of redirected aggression in primates resemble those seen in kin-based reconciliation, and both indicate that monkeys can recognize third-party relationships. Although we have not yet begun work on redirected aggression in spotted hyenas, it is very common in this species, and we plan in the near future to use these interactions to ask once again whether hyenas can recognize third-party relationships.

---

### Conclusions

Much of our work is still in progress, but the initial results from our studies of social cognition in hyenas suggest that even though these predators live in the same types of complex, stable groups as many primates do, they lack some of the mental abilities common in monkeys and apes. If spotted hyenas cannot recognize third-party social relationships, then features of the social environment common to both primates and spotted hyenas may be necessary, but cannot be sufficient, to account for the evolution of the primates' ability to recognize relationships among unrelated group members. Our data to date suggest that hyenas may accomplish many of the same social feats as monkeys do by following simple rules. If it is confirmed in our ongoing studies, the finding that hyenas cannot recognize third-party relationships will suggest that the social complexity hypothesis should be modified to focus only on those aspects of primate social life that are not shared with gregarious carnivores.

---

### Acknowledgments

This work was supported by National Science Foundation grant IBN9906445. We thank L. Smale for insightful comments on an earlier draft of this paper.

---

### References

- Andelman, S. J. (1985). Ecology and Reproductive Strategies of Vervet Monkeys (*Cercopithecus aethiops*) in Amboseli National Park, Kenya. Ph.D. thesis, University of Washington, Seattle.
- Aurlei, F. and de Waal, F. B. M. (2000). *Natural Conflict Resolution*. Berkeley: University of California Press.
- Bacchman, C. and Kummer, H. (1980). Male assessment of female choice in hamadryas baboons. *Behavioral Ecology and Sociobiology* 6: 315–321.
- Barton, R. A. and Dunbar, R. I. M. (1997). Evolution of the social brain. In *Machiavellian Intelligence II: Extensions and Evaluations*, A. Whiten and R. W. Byrne, eds., pp. 240–263. Cambridge: Cambridge University Press.
- Boesch, C. and Boesch, H. (1989). Hunting behaviour of wild chimpanzees in the Taï National Park. *American Journal of Physical Anthropology* 78: 547–573.
- Boinski, S. and Garber, P. (eds.) (2000). *On the Move: How and Why Animals Travel in Groups*. Chicago: University of Chicago Press.
- Byrne, R. W. (1994). The evolution of intelligence. In *Behaviour and Evolution*, P. J. B. Slater and T. R. Halliday, eds., pp. 223–265. Cambridge: Cambridge University Press.
- Byrne, R. W. and Whiten, A. (eds.) (1988). *Machiavellian Intelligence*. Oxford: Clarendon Press.
- Chapais, B. (1992). The role of alliances in the social inheritance of rank among female primates. In *Coalitions and Alliances in Humans and Other Animals*, A. H. Harcourt and F. B. M. de Waal, eds., pp. 29–60. Oxford: Oxford University Press.
- Cheney, D. L. and Seyfarth, R. M. (1980). Vocal recognition in free-ranging vervet monkeys. *Animal Behaviour* 28: 362–367.
- Cheney, D. L. and Seyfarth, R. M. (1983). Non-random dispersal in free-ranging vervet monkeys: Social and genetic consequences. *American Naturalist* 122: 392–412.
- Cheney, D. L. and Seyfarth, R. M. (1989). Redirected aggression and reconciliation among vervet monkeys, *Cercopithecus aethiops*. *Behaviour* 110: 258–275.
- Cheney, D. L. and Seyfarth, R. M. (1990). *How Monkeys See the World*. Chicago: University of Chicago Press.

- Clutton-Brock, T. H., Guinness, F. E., and Albon, S. D. (1982). *Red Deer: Behavior and Ecology of Two Sexes*. Chicago: University of Chicago Press.
- Connor, R. C., Smolker, R. A., and Richards, A. F. (1992). Dolphin alliances and coalitions. In *Coalitions and Alliances in Humans and Other Animals*, A. H. Harcourt and F. B. M. de Waal, eds., pp. 415–443. Oxford: Oxford University Press.
- Datta, S. B. (1986). The role of alliances in the acquisition of rank. In *Primate Ontogeny, Cognition and Social Behavior*, G. J. Else and P. C. Lee, eds., pp. 219–225. Cambridge: Cambridge University Press.
- de Waal, F. B. M. (1993). Reconciliation among primates: A review of empirical evidence and unresolved issues. In *Primate Social Relationships*, W. A. Mason and S. P. Mendoza, eds., pp. 111–114. New York: State University of New York Press.
- Dunbar, R. I. M. and Bever, J. (1998). Neocortex size predicts group size in carnivores and some insectivores. *Ethology* 104: 695–708.
- East, M. L., Hofer, H., and Wickler, W. (1993). The erect “penis” as a flag of submission in a female-dominated society: Greetings in Serengeti spotted hyenas. *Behavioral Ecology and Sociobiology* 33: 355–370.
- Eng, A. L., Esch, K., Smale, L., and Holekamp, K. E. (2000). Mechanisms of maternal rank “inheritance” in the spotted hyaena, *Crocuta crocuta*. *Animal Behaviour* 60: 323–332.
- Eng, A. L., Siebert, E., Jensen, A. D., and Holekamp, K. E. (in preparation). Do spotted hyenas use allies strategically?
- Gouzoules, S., Gouzoules, H., and Marler, P. (1995). Representational signaling in non-human primate vocal communication. In *Current Topics in Primate Vocal Communication*, E. Zimmerman, J. D. Newman, and U. Jürgens, eds., pp. 235–252. New York: Plenum.
- Guggisberg, C. A. W. (1962). *Simba*. London: Bailey Brothers and Swinfen.
- Harcourt, A. H. and de Waal, F. B. M. (eds.) (1992). *Coalitions and Alliances in Humans and Other Animals*. Oxford: Oxford University Press.
- Harvey, P. H. and Krebs, J. R. (1990). Comparing brains. *Science* 249: 140–146.
- Hofer, H. and East, M. L. (2000). Conflict management in female-dominated spotted hyenas. In *Natural Conflict Resolution*, F. Aureli and F. B. M. de Waal, eds., pp. 232–234. Berkeley: University of California Press.
- Holekamp, K. E. and Smale, L. (1991). Rank acquisition during mammalian social development: The “inheritance” of maternal rank. *American Zoologist* 31: 306–317.
- Holekamp, K. E., Cooper, S. M., Katona, C. I., Berry, N. A., Frank, L. G., and Smale, L. (1997). Patterns of association among female spotted hyenas (*Crocuta crocuta*). *Journal of Mammalogy* 78: 55–64.
- Holekamp, K. E., Boydston, E. E., Szykman, M., Graham, I. Nutt, K. J., Birch, S., Piskiel, A., and Singh, M. (1999). Vocal recognition in the spotted hyaena and its possible implications regarding the evolution of intelligence. *Animal Behaviour* 58: 383–395.
- Holekamp, K. E., Boydston, E. E., and Smale, L. (2000). Group travel in social carnivores. In *On the Move: How and Why Animals Travel in Groups*, S. Boinski and P. Garber, eds., pp. 587–627. Chicago: University of Chicago Press.
- Horrocks, J. and Hunte, W. (1983). Maternal rank and offspring rank in vervet monkeys: An appraisal of the mechanisms of rank acquisition. *Animal Behaviour* 31: 772–782.
- Kamil, A. C. (1987). A synthetic approach to the study of animal intelligence. *Nebraska Symposium on Motivation* 7: 257–308.
- Macphail, E. M. (1982). *Brain and Intelligence in Vertebrates*. Oxford: Clarendon Press.
- Milton, K. (1981). Distribution patterns of tropical plant foods as an evolutionary stimulus to primate mental development. *American Anthropologist* 83: 534–548.
- Moss, C. J. (1988). *Elephant Memories*. Boston: Houghton Mifflin.
- Packer, C. (1994). *Into Africa*. Chicago: University of Chicago Press.
- Povinelli, D. J. and Preuss, T. M. (1995). Theory of mind: Evolutionary history of a cognitive specialization. *Trends in Neuroscience* 18: 418–424.
- Seyfarth, R. M. (1980). The distribution of grooming and related behaviors among adult female vervet monkeys. *Animal Behaviour* 28: 798–813.



- Seyfarth, R. M. and Cheney, D. L. (1984). Grooming, alliances, and reciprocal altruism in vervet monkeys. *Nature* 308: 541–543.
- Smale, L., Frank, L. G., and Holekamp, K. E. (1993). Ontogeny of dominance in free-living spotted hyenas: Juvenile rank relations with adults. *Animal Behaviour* 46: 467–477.
- Smale, L., Nunes, S., and Holekamp, K. E. (1997). Sexually dimorphic dispersal in mammals: Patterns, causes and consequences. *Advances in the Study of Behavior* 26: 181–250.
- Stander, P. E. (1992a). Foraging dynamics of lions in a semi-arid environment. *Canadian Journal of Zoology* 70: 8–21.
- Stander, P. E. (1992b). Cooperative hunting in lions: The role of the individual. *Behavioral Ecology and Sociobiology* 29: 445–454.
- Tilson, R. T. and Hamilton, W. J. (1984). Social dominance and feeding patterns of spotted hyaenas. *Animal Behaviour* 32: 715–724.
- Tomasello, M. and Call, J. (1997). *Primate Cognition*. Oxford: Oxford University Press.
- Wahaj, S., Guze, K., and Holekamp, K. E. (2001). Reconciliation in the spotted hyena (*Crocuta crocuta*). *Ethology* 107: 1057–1074.
- Walters, J. (1980). Interventions and the development of dominance relationships in female baboons. *Folia Primatologica* 34: 61–89.
- Zabel, C. J., Glickman, S. E., Frank, L. G., Woodmansee, K. B., and Keppel, G. (1992). Coalition formation in a colony of prepubertal spotted hyaenas. In *Coalitions and Alliances in Humans and Other Animals*, A. H. Harcourt and F. B. M. de Waal, eds., pp. 113–135. Oxford: Oxford University Press.





This is a section of [doi:10.7551/mitpress/1885.001.0001](https://doi.org/10.7551/mitpress/1885.001.0001)

# The Cognitive Animal

## Empirical and Theoretical Perspectives on Animal Cognition

**Edited by: Marc Bekoff, Colin Allen, Gordon M. Burghardt**

### Citation:

*The Cognitive Animal: Empirical and Theoretical Perspectives on Animal Cognition*

Edited by: Marc Bekoff, Colin Allen, Gordon M. Burghardt


DOI: 10.7551/mitpress/1885.001.0001

ISBN (electronic): 9780262268028

Publisher: The MIT Press

Published: 2002

The open access edition of this book was made possible by generous funding and support from Arcadia – a charitable fund of Lisbet Rausing and Peter Baldwin



The MIT Press

©2002 Massachusetts Institute of Technology

All rights reserved. No part of this book may be reproduced in any form by any electronic or mechanical means (including photocopying, recording, or information storage and retrieval) without permission in writing from the publisher.

This book was set in Times New Roman on 3B2 by Asco Typesetters, Hong Kong. Printed and bound in the United States of America.

Library of Congress Cataloging-in-Publication Data

The cognitive animal: empirical and theoretical perspectives on animal cognition /  
edited by Marc Bekoff, Colin Allen, and Gordon M. Burghardt.

p. cm.

“A Bradford book.”

Includes bibliographical references.

ISBN 0-262-02514-0 (hc. : alk. paper)—ISBN 0-262-52322-1 (pbk. : alk. paper)

1. Cognition in animals. I. Bekoff, Marc. II. Allen, Colin. III. Burghardt, Gordon M.,  
1941—

QL785 .C485 2002

591.5'13—dc21

2001057965