

# The Effects of Learning on the Evolution of Saussurean Communication

Edgar E. Vallejo<sup>1,2</sup> and Charles E. Taylor<sup>2</sup>

<sup>1</sup>Tecnológico de Monterrey, Campus Estado de México, Atizapán de Zaragoza 52926, México

<sup>2</sup>University of California Los Angeles, Los Angeles, CA 90025, USA

## Abstract

This paper presents a computational framework for studying the influence of learning on the evolution of communication. In our model, an evolving population of learning agents is engaged in pairwise communicative interactions. Simulation results show the genetic assimilation of transmission behaviors as a consequence of saussurean learning.

## Introduction

The evolution of communication is an excellent domain for studying fundamental questions of artificial life research. Previous work by Ackley and Littman (2), Arita and Taylor (3), Di Paolo (5), Hashimoto and Ikegami (8), and Steels (18), among others, have shown that we are able to explore important issues such as emergence, self-organization and cultural evolution within this framework.

The aim of this work is to study the influence of learning on the evolution of communication. We believe this is an instrumental aspect for understanding the origin and evolution of communication systems with the complexity of human languages.

In nature, learning seem to influence both the association of signaling behaviors to appropriate external referents and the development of responses to different signal types (10). However, there are several examples of innate underpinnings in transmission behaviors. For example, young vervet monkeys seem predisposed to alarm calling, dividing up the universe of predators into different ill-defined categories, leaving social experience to sharpen the boundaries of exclusive predator categories and its corresponding association to particular call types (4). Moreover, there is indication of innate predispositions in human languages (16). On the other hand, there is a growing conviction that learning plays a substantial role in the development of how a receiver responds on hearing a signal (10).

A fundamental question that arises in the context of animal communication is whether a communication system is likely to be symbolic. According to Marler, for an animal communication system to qualify as symbolic, information about external referents has to be both encoded by signalers and decoded by receivers (10). This definition is consistent with the nature of the saussurean linguistic sign (17).

In his seminal paper, Hurford proposed a model for studying the evolution of saussurean communication (9). In this model, an agent consists of two probabilistic matrices that provide a framework for semiotic interactions. Computational simulations were conducted to investigate the evolutionary potentials of different learning strategies. Further studies by Oliphant (14), Oliphant and Batali (15) and Nowak et al (13) have contributed to elucidate the fundamental properties of this model.

These studies have been largely conducted within the framework of Lamarkian evolution. In most cases, learned communicative behaviors are written back to the genetic description of agents and thus transmitted to offspring during reproduction. We believe that Darwinian evolution provides a more convenient framework for studying the evolution of communication. In their influential work, Hinton and Nowlan proposed a computational framework for conducting studies on the effects of learning on evolution in Darwinian evolution (7) (1). This holds much promise. For example, Turkel has shown that this computational framework can be used for exploring how the capacity of human language could have evolved via natural selection (19).

In this work, we conducted computational studies about the influence of learning on the evolution of communication. Experimental results show the genetic assimilation of transmission behaviors. This results are consistent with dominant theories on the evolution of communication (6).

Copyrighted Material

## The model

### Agent architecture

The formal definition of the agent architecture presented below is based on considerations of the models proposed by Hurford (9) and Oliphant (14).

**Agent** Let  $O = \{o_1, \dots, o_n\}$  be a finite set of  $n$  objects and  $S = \{s_1, \dots, s_m\}$  be a finite set of  $m$  signals. A *learning communicative agent* is a triple  $(\delta, \phi, \sigma)$ , where

1.  $\delta : O \rightarrow S \cup \{s_\#\}$  is the transmission function, where  $s_\#$  is the undetermined signal,
2.  $\phi : S \rightarrow O \cup \{o_\#\}$  is the reception function, where  $o_\#$  is the undetermined object, and
3.  $\sigma$  is the learning strategy.

**Communication** An agent  $A_1 = (\delta_1, \phi_1, \sigma_1)$  communicates to an agent  $A_2 = (\delta_2, \phi_2, \sigma_2)$  as follows. Initially,  $A_1$  perceives an object  $o_i$  and produces a signal  $s_j$  according to the mapping described by the transmission function  $\delta_1$ , such that  $\delta_1(o_i) = s_j$ . Once  $A_1$  produces the signal  $s_j$ , the agent  $A_2$  interprets the signal  $s_j$  as the object  $o_k$  according to the mapping described by the reception function  $\phi_2$ , such that  $\phi_2(s_j) = o_k$ . A communication event from  $A_1$  to  $A_2$  is successful if the following conditions are satisfied

1.  $\delta_1(o_i) = s_j$ ,
2.  $\phi_2(s_j) = o_k$ , and
3.  $o_i = o_k$

**Homonymy** An agent  $A = (\delta, \phi, \sigma)$  is said to possess homonymy if there exist a signal  $s_j \neq s_\#$  and a pair of objects  $o_i$  and  $o_k$  that satisfy the following conditions

1.  $\delta(o_i) = s_j$ ,
2.  $\delta(o_k) = s_j$ , and
3.  $o_i \neq o_k$

**Synonymy** An agent  $A = (\delta, \phi, \sigma)$  is said to possess synonymy if there exist an object  $o_i \neq o_\#$  and a pair of signals  $s_j$  and  $s_k$  that satisfy the following conditions

1.  $\phi(s_j) = o_i$ ,
2.  $\phi(s_k) = o_i$ , and
3.  $s_j \neq s_k$

**Innate transmission** Let  $A = (\delta, \phi, \sigma)$  be an agent. A transmission from  $A$  for a given object  $o_i$  is said to be innate if  $\delta(o_i) \neq s_\#$  and is said to be subject to learning if  $\delta(o_i) = s_\#$ .

**Innate reception** Let  $A = (\delta, \phi, \sigma)$  be an agent. A reception of  $A$  for a given signal  $s_j$  is said to be innate if  $\phi(s_j) \neq o_\#$  and is said to be subject to learning if  $\phi(s_j) = o_\#$ .

**Learning** In our model, both transmission and reception behaviors are partially learned. Before a communication event from  $A_1$  to  $A_2$  takes place,  $A_1$  replaces the undetermined signals in  $\delta_1$  with signals in  $S$  using the learning strategy  $\sigma_1$ . Similarly,  $A_1$  replaces the undetermined objects in  $\phi_1$  using the learning strategy  $\sigma_1$ . Agent  $A_2$  proceeds similarly.

A fundamental aspect of our model is that learning is performed for communication purposes and does not permanently modify the actual description of an agent.

### Learning strategies

We consider three different learning strategies: imitator, calculator and saussurean, after (9). In addition, we introduce a fourth learning strategy: random learner.

**Imitator** An imitator agent replaces the undetermined signals in his transmission function by the corresponding signals in the transmission function of another agent. Similarly, he replaces the undetermined objects in his reception function by the corresponding objects in the reception function of another agent.

Formally, an agent  $A_1 = (\delta_1, \phi_1, \sigma_1)$  imitates an agent  $A_2 = (\delta_2, \phi_2, \sigma_2)$  as follows.

1.  $\delta_1(o_i)$  is set to  $\delta_2(o_i)$  if  $\delta_1(o_i) = s_\#$ ,  $\delta_2(o_i) \neq s_\#$ , for  $i = 1, \dots, n$ , and
2.  $\phi_1(s_j)$  is set to  $\phi_2(s_j)$  if  $\phi_1(s_j) = o_\#$ ,  $\phi_2(s_j) \neq o_\#$ , for  $j = 1, \dots, m$ .

**Calculator** A calculator agent replaces the undetermined signals in his transmission function in such a way that a communication event to another agent would be successful. Similarly, he replaces the undetermined objects in his reception function in such a way that the communication event from another agent would be successful.

Formally, an agent  $A_1 = (\delta_1, \phi_1, \sigma_1)$  calculates an agent  $A_2 = (\delta_2, \phi_2, \sigma_2)$  as follows.

1.  $\delta_1(o_i)$  is set to  $s_k$  if  $\phi_2(s_k) = o_i$ ,  $\delta_1(o_i) = s_\#$ , for  $i = 1, \dots, n$ , and

strategy	code
imitator	0
calculator	1
saussurean	2
random	3

Table 1: Codification of learning strategies

- $\phi_1(s_j)$  is set to  $o_k$  if  $\delta_2(o_k) = s_j$ ,  $\phi_1(s_j) = o_{\#}$ , for  $j = 1, \dots, m$ .

**Saussurean** A saussurean agent replaces the undetermined signals in his transmission function by the corresponding signals in the transmission function of another agent. Similarly, he replaces the undetermined objects in his reception function in such a way that a communication event to himself would be successful.

Formally, an agent  $A_1 = (\delta_1, \phi_1, \sigma_1)$  is saussurean with respect to and agent  $A_2 = (\delta_2, \phi_2, \sigma_2)$  as follows.

- $\delta_1(o_i)$  is set to  $\delta_2(o_i)$  if  $\delta_1(o_i) = s_{\#}$ ,  $\delta_2(o_i) \neq s_{\#}$ , for  $i = 1, \dots, n$ , and
- $\phi_1(s_j)$  is set to  $o_k$  if  $\delta_1(o_k) = s_j$ ,  $\phi_1(s_j) = o_{\#}$ , for  $j = 1, \dots, m$ .

**Random** A random agent can assume any of the strategies described above. Before a communication event takes place, a random agent randomly selects one of the following strategies: imitator, calculator or saussurean.

For convenience, we consider a codification of learning strategies as shown in table 1.

## Evolution of communication

In our model, a population of learning communicative agents are intended to evolve successful communication at the population level. We use genetic algorithms for this purpose. The design decisions presented below are based on considerations of the performance of genetic algorithms in practical applications (12).

**Genome representation** A learning communicative agent  $A = (\delta, \phi, \sigma)$  is represented linearly as follows

$$A = (\delta(o_1), \dots, \delta(o_n), \phi(s_1), \dots, \phi(s_m), \sigma)$$

For example, consider a set of objects  $O = \{1, 2, 3\}$ , a set of signals  $S = \{a, b, c\}$ . Consider imitator agent  $A_1 = (\delta_1, \phi_1, \sigma_1)$ . In addition, consider the functions  $\delta_1$  and  $\phi_1$  described in table 2.

$o_i$	$\delta_1(o_i)$	$s_j$	$\phi_1(s_j)$
1	$b$	$a$	$o_{\#}$
2	$s_{\#}$	$b$	3
3	$a$	$c$	2

Table 2: Transmission and reception functions

The linear representation of agent  $A_1$  is

$$(b, s_{\#}, a, o_{\#}, 3, 2, 1)$$

**Genetic operators** Agents produce a new offspring by means of genetic operators. Fitness proportional selection, one-point recombination and point mutation operate on the linear representation of agents described above.

**Fitness function** Fitness was defined as the communicative accuracy of learning communicative agents. The communicative accuracy is the ability of an agent to successfully communicate with a collection of other agents.

Let  $P$  be a finite population of agents,  $A$  be an agent in  $P$ , and  $Q \subseteq P$  be a non empty collection of agents. The communicative accuracy of  $A$  with respect to  $Q$  given the set of objects  $O = \{o_1, \dots, o_n\}$  and the set of signals  $S = \{s_1, \dots, s_m\}$ ,  $C(A, Q, O)$ , is defined as

$$C(A, Q, O) = \frac{\sum_{o_i \in O} \sum_{A_k \in Q} c(A, A_k, o_i) + c(A_k, A, o_i)}{|Q|}$$

where  $c(A, A_k, o_i) = 1$  if the communication event from  $A$  to  $A_k$  is successful given the object  $o_i$ , and 0 otherwise;  $|Q|$  is the cardinality of  $Q$ .  $c(A_k, A, o_i)$  is defined similarly.

## Experiments and results

Experiments were conducted to investigate whether a population of learning communicative agents is likely to arrive to successful communication at the population level. In addition, we validated the evolutionary performance of competing learning strategies. Most importantly, we were interested in exploring the effects of learning on the genetic description of an evolving population of learning communicative agents. The simulation procedure is described in table 3.

Several simulations were conducted using different combinations of parameter values as shown in table 4. The following were the major results:

1. Create an initial random population  $P$  of agents
2. Do **until** number generations is met
  - (a) **For each** individual  $A_i = (\delta_i, \phi_i, \sigma_i) \in P$  **do**
    - i. Perform the learning process of  $A_i$  according to the learning strategy  $\sigma_i$  with respect to a random agent  $A_h \in P$
    - ii. Select a random subpopulation of agents  $Q \subseteq P$
    - iii. Perform the learning process for all  $A_j \in Q$  according to the learning strategy of  $A_j$  with respect to a random agent  $A_k \in P$
    - iv. Measure the communicative accuracy of  $A_i$  with respect to  $Q$ ,  $C(A_i, Q, O)$ , given the set of objects  $O$
  - End for**
  - (b) Select two individuals  $A_{mother} \in P$  and  $A_{father} \in P$  for reproduction using fitness proportional selection
  - (c) Produce an offspring  $A_{new}$  from  $A_{mother}$  and  $A_{father}$  using one-point recombination and point mutation
  - (d) Select a random individual  $A_{old} \in P$
  - (e) Replace  $A_{old}$  by  $A_{new}$
- End do**

Table 3: Simulation procedure

Parameter	Value
generations	2000–3000
population $P$	128–512
subpopulation $Q$	4–16
signals	4–8
meanings	4–8
crossover probability	0.6–0.7
mutation probability	0.001–0.01

Table 4: Parameters for simulations

1. Agents arrived at highly successful communication at the population level. However, simulations showed that there exists a threshold condition on the number of interactions (size of set  $Q$ ) required to achieve accuracy in communication. Figure 1 shows the results of the simulations for different number of interactions. Agents reached local minima in communication accuracy as the number of interactions is reduced.
2. Local minima were produced mostly by the presence of homonymy – i.e. when the same signal is used to describe two or more different objects. Synonymy – when two or more signals are interpreted as the same object, did not appear consistently.
3. In pairwise strategy contests, populations of calculators dominated a populations of imitators. However, populations of saussureans outcompeted both calculators and imitators.
4. In all strategy contests: imitators, calculators, saussureans and randoms, experimental results showed the superiority of the saussurean learning strategy. In most cases, saussureans took over the entire population. Only rarely, did imitators or calculators dominate such populations. Random learners never became dominant. Figure 2 shows the frequency of strategies in the population for a typical simulation where all strategies were initially present.
5. The undetermined signal trait disappeared in the population. All the signals produced for every object were genetically assimilated in the transmission gene segment of the genome for all agents. The frequency of undetermined signals in the population as evolution proceeded is shown in figure 3.
6. The undetermined objects trait prevailed in the population. Objects interpreted for every signal were not genetically assimilated in the reception gene segment of the genome. The frequency of undetermined objects in the population as evolution proceeded is shown in figure 3.
7. Most simulations considered a set of 4 objects and a set of 4 signals. Both homonymy and synonymy began to appear more consistently when both the number of objects and signals was increased. However, homonymy continued to be more common.

## Discussion

The overall experimental results indicate that given a sufficient number of interactions, a population of learning communicative agents is capable of arriving at highly successful communication. Surprisingly, transmission behaviors became innate as a consequence of the evolution saussurean learning. On the other hand, reception behaviors prevailed as characteristics that are subject to learning. Our results are consistent with several hypotheses formulated for the evolution of communication (10).

### Why transmission was genetically assimilated

In general, experimental results showed that there is an evolutionary preference for saussurean agents. Saussurean learning implements a tight coupling between transmission and reception behaviors.

Why did signaling behaviors become innate? First, imitation of transmission in a static environment provides the opportunity for genetic assimilation of signaling behaviors. Second, a tight coupling between reception and innate transmission provide the opportunity for successful learning in reception behaviors.

### Why saussureans prevailed

From the evolutionary perspective, how good is saussurean learning for evolving communication?. Could an alternative communicative strategy invade a population of saussurean learners?.

Maynard-Smith(11) has demonstrated that game theory can be used as a framework to explain the evolution of most phenotypic traits in situation in which fitness of a trait depends on what others are doing. He has also provided the notion of evolutionary stable strategy (ESS). An ESS is a phenotype such that, if almost all individuals have that phenotype, no alternative phenotype can invade the population.

In our model, experimental results showed that saussurean learning is likely to be an ESS. However, further studies are required to formally demonstrate what types of learning are evolutionary stable strategies.

In this study, we did not consider the cost of producing a signal. Previous studies suggest that honesty of communication become an issue when cost is considered (2).




Figure 1: Threshold condition on interactions




Figure 2: Evolved strategies




Figure 3: Genetically assimilated traits

Copyrighted Material

## Acknowledgements

This work was supported by the UC MEXUS-CONACYT, the UCLA Center for Embedded Sensor Networks, the Defense Advance Research Projects (DARPA), administered by the Army Research Office under Emergent Surveillance Plexus MURI Award No. DAAD19-01-1-0361, and DARPA MURI administered by the US Air Force No. F49620-01-1-0361. Any opinions, findings and conclusions or recommendations expressed in this publication are those of the authors and do not necessarily reflect the views of the sponsoring agencies.

## References

- [1] Ackley, D.H. and Littman, M.L. 1992. Interactions between learning and evolution. In C.G. Langton, C.E. Taylor, J.D. Farmer and S. Rasmussen. 1992. *Artificial Life II*. Addison Wesley.
- [2] Ackley D,H. and Littman, M.L. 1994. Altruism in the Evolution of Communication. In R.A. Brooks, P. Maes, (eds), *Artificial Life IV. Proceedings of the Fourth International Conference on Artificial Life*, The MIT Press, pp.40–48.
- [3] Arita, T. and Taylor, C.E. 1996. A simple model for the evolution of communication. In: L.J. Fogel, P.J. Angeline, T. Back, (eds), *Evolutionary Programming V*. The MIT Press, pp.405–410.
- [4] Cheney, D. and Seyfarth, R.M. 1990. *How Monkeys See the World*. University of Chicago Press.
- [5] Di Paolo, E. A. 1998. An investigation into the evolution of communication. *Adaptive Behavior*, 6(2):285–324.
- [6] Hauser, M. 1996. *The Evolution of Communication*. The MIT Press.
- [7] Hinton, G.E. and Nowlan, S.J. 1987. How learning can guide evolution. *Complex Systems*1, pp.495–502.
- [8] Hashimoto, T. and Ikegami, T. 1996. Emergence of net-grammar in communicating agents, *BioSystems* 38:1–14.
- [9] Hurford, J. R. 1989. Biological evolution of the saussurean sign as a component of the language acquisition device. *Lingua* 77, pp.187–222.
- [10] Marler, P. 1998. Animal communication and human language. In N.G. Jablonski and L.C. Aiello (eds.) *The Origin and Diversification of Language*. University of California Press.
- [11] Maynard-Smith J (1982), *Evolution and the Theory of Games*. Cambridge University Press.
- [12] Mitchell, M. 1996. *An introduction to genetic algorithms*. The MIT Press.
- [13] Nowak M.A., Plotkin, J.B. and Krakauer, D. 1999. The evolutionary language game. *Journal of Theoretical Biology*, 200(2):147–162.
- [14] Oliphant, M. 1996. The dilemma of saussurean communication. *BioSystems*, 37(1-2), pp.31–38.
- [15] Oliphant, M. and Batali, J. 1997. Learning and the emergence of coordinated communication. *Center for Research on Language Newsletter*, 11(1).
- [16] Pinker, S. 1994. *The Language Instinct*. William Morrow and Company.
- [17] de Saussure, F. 1986. *Course in general linguistics*. Open Court.
- [18] Steels, L. 1996. Self-organizing vocabularies. In: C. Langton and T. Shimohara (eds), *Artificial Life V. Proceedings of the Fifth International Conference on Artificial Life*, The MIT Press, pp.179–184.
- [19] Turkel, W.J. 2002. The learning guided evolution of natural language. In T. Briscoe (ed.) *Linguistic Evolution through Language Acquisition*. Cambridge University Press, pp.235–254.