

Index

- agricultural water use 313–314, 318
- Amurga Massif, Gran Canaria 130–135
- aquifer geometry models 214–215
- aquifers as archives 2, 12–13, 155
- Aquitaine Basin 300–302
- Astian aquifer 101–104, 203, 308
 - groundwater management 322
- Aveiro Cretaceous aquifer 139–153
 - geology 140–142
 - hydrogeochemistry 144–150
 - inverse modelling 246–247
 - isotope analysis 150–152, 203
 - modern groundwater 282
 - noble gas recharge temperature 151
 - palaeohydrology 302–306
 - radiocarbon analysis 150–151
 - water quality 316
- BALANCE 232
- Baltic Ice Lake 25, 179–181, 292–293
- Baltic Sea 181, *see also* Baltic Ice Lake
- Belgium, palaeoclimate 50
- boreholes 252
 - Brighton 260–265
 - contamination 36
 - North Kent 78
- British Isles, climate change 71–73
- Cambrian–Vendian aquifer 7–8, 17–27, 202
 - age structure 25
 - groundwater flow model 20–21
 - hydrogeology 17–21
 - isotopic analysis 21–23
 - modern water 23
 - noble gas recharge temperature 23–25
 - palaeogroundwater formation 25–26
 - recharge 22, 26
- cation exchange 51–59, 112, 234–247
- closed aquifer systems 214–215, 221–222
- coastal aquifers 5–6
 - management 7, 314–316
 - water quality 316–318
- confined aquifers 109–112, 251, 257
- connate water 67, 111–112
- contaminants as tracers 277–278
- Cretaceous Chalk aquifer 74–77
- cryogenic metamorphism 26
- Darum 42–43
- dating tools 273–277
- Denmark, Cenozoic succession 33, 34
- density driven flow 223
- desalination 318
- Devensian glaciation 4, 71–73, 84–86, 297, 299
- DIC *see* dissolved inorganic carbon
- diffusion 41
- diffusive transport 221–222
- dissolved inorganic carbon 38, 40–41
- dissolved organic carbon 38
- DOC *see* dissolved organic carbon
- Dogger aquifer 94–101, 203
 - modern groundwater 282
- Doñana aquifer
 - contamination 281–282
 - geology 116–118
 - hydrogeochemistry 118
 - isotope analysis 119–122, 203
 - palaeohydrology 302–306
- Dorset Basin Chalk aquifer 82–83
- drilling 319
- East Midlands Sandstone aquifer 10–12, 83–86
 - impact of glaciation 225
 - inverse geochemical modelling 234–239
 - isotope analysis 84–85, 202
 - modern groundwater 280, 281
 - water quality 316
- East Street borehole 260–265
- eastern Baltic, palaeohydrology 290–293
- Ebro delta 306–308
- Elbe Urstromtal 31–32, 293–294
- England
 - palaeowater 88–90
 - sea-level changes 72–73
- English Channel
 - palaeohydrology 297–300
 - sea-level change 73, 177–179, 300
- environmental tracers 273–277
- Estonia 17–27
 - groundwater
 - evolution 292
 - exploitation 21, 26
 - isotopic analysis 21–23
 - noble gas recharge temperature 23–25
 - quality 316
 - hydrogeology 17–21, *see also* Cambrian–Vendian aquifer
 - palaeogroundwater formation 25–26
- Fennoscandian ice sheet 25–26, 170–175, 177, 224, 292–293
- fissured aquifers 307–308
- Fjand 36, 42–43
 - geophysical logging 256
- Flandrian transgression 109–110, 116
- Flemish Valley 50, 66
- flow systems 215
 - characteristic times 215–216
 - density driven flow 223
 - diffusive transport 221–222
 - impact of glaciers 223–225
 - impact of permafrost 225–226
 - interface movement 218–221

- flow systems *continued*
 steady state interface 216–218
 fluid logging 252–265
 forward modelling 232, 239–240
 fossil organic matter 40
 France, coastal aquifers 93–105
 French Atlantic coast 177, 300–302
 freshening 33, 38, 50, 65
- Gdov aquifer 19–20
 geophysical logging 36, 42, 252–268
 in coastal aquifers 253–265
 methods 252
 glaciation
 Devensian 4, 71–73, 84–86, 171
 Glatt Valley aquifer 155–156
 Weichselian 170–174
 glaciers
 impact on flow systems 223–225
 East Midlands aquifer 86, 225
 northern Belgium 224
 Ribe Formation 225
 Glatt Valley aquifer 155–161, 203
 groundwater
 age estimate 194–198
 Cambrian–Vendian aquifer 25
 Ribe Formation 38, 40–41
 emplacement 13–14
 as injection fluid 42
 isotopic composition 185–186
 management 314, 318–326
see also modern groundwater
 groundwater evolution 4–5, 289–309
 East Midlands Sandstone aquifer 10–12
 eastern Baltic 290–293
 English Channel 297–300
 French Atlantic coast 300–302
 Iberian Atlantic coast 302–306
 sea-level change 289–290, 297, 300–302, 305, 308
 southern North Sea 293–297
 western Mediterranean coast 306–308
 groundwater flow modelling
 Cambrian–Vendian aquifer 20–21
 Dorset Basin 83
 East Midlands Sandstone aquifer 86
 Ledo-Paniselian aquifer 50–51, 60–65
 Ribe Formation 41–42
see also flow systems
 groundwater interfaces 217–221, 272
 groundwater quality 231, 272, 284–285
 groundwater systems 3
 timescales of change 3–4
- Heligoland Channel 31–32, 35, 293–294
 human impact on groundwater 38, 271–272
 indicators 273
 hydraulic conductivity
 Cambrian–Vendian aquifer 20–21, 26
 Ledo-Paniselian aquifer 62
 Ribe formation 35–36
 hydrochemical evolution 203–205
 hydrochemical modelling 231–248
- hydrogeochemical transport models 234, *see also*
 forward modelling
- Iberian Atlantic coast 302–306
 Inca-Sa Pobra aquifer 123–130, 306–307
 contamination 282
 geology 123–124
 hydrogeochemistry 124–126
 isotope analysis 126–128
 interface modelling 217–221
 inverse geochemical modelling 231, 234–248
 Irish Sea 177
 isolation preservation 321
 isotope analysis
 Amurga Massif, Gran Canaria 133–135
 Astian aquifer 101–103
 Aveiro Cretaceous aquifer 150–152
 Cambrian–Vendian aquifer 21–23
 Cretaceous Chalk aquifer 75–76
 Dogger aquifer 94–100
 Doñana aquifer 119–122
 Dorset Basin 82–83
 East Midlands Sandstone aquifer 11–12, 84–85
 Glatt Valley aquifer 157–159
 Inca-Sa Pobra aquifer 126–128
 Kent aquifers 78–79
 Ledo-Paniselian aquifer 54–56
 Llobregat River Delta aquifers 112–115
 Ribe Formation aquifer 38
 South Downs aquifer 81–82
 Spilsby Sandstone 76
 isotopic composition of groundwater 185–186
 isotopic methods 193–194
 European scale 206–208
 PALAEAUX programme 200–203
 recharge condition indicators 198–200
 residence time indicators 194–198
see also dating tools
- Jylland
 geology 30–34
 hydrogeological model 34–35
 water supply 38, 42
- karstic aquifers 307–308
 Kent aquifers 78–79
 Kotlin Formation 19–20
- last glacial maximum 29–31, 66, 85, 108, 164–167, 170,
 176–177, 292, 305
 Ledo-Paniselian aquifer
 cation exchange 51–59
 effects of pumping 62–65
 forward modelling 239–240
 geology 50
 groundwater flow 51
 hydrodynamic model 60–65
 impact of permafrost 67–68, 225–226
 inverse modelling 240–246
 isotope analysis 54–56, 204

- palaeoclimate 50
 - palaeoenvironmental model 65–67
 - pore water analysis 56–58
 - radiocarbon analysis 52–56, 202
 - recharge conditions 66
- LGM *see* last glacial maximum
- Lincolnshire, Cretaceous Chalk aquifer 74–76
- Llobregat River Delta aquifers 108–116, 306
 - exploitation of groundwater 108–109, 111
 - groundwater management 322–324
 - hydrogeochemistry 111–112
 - isotope analysis 112–115
- Lontova Formation 20
- Lower Greensand aquifer 79, 82–83
- Lükati–Lontova aquitard 19

- management of coastal aquifers 314–326
- mass balance models 232–233, *see also* inverse
 - geochemical modelling
- Mediterranean
 - aquifers 306–308
 - sea-level change 182–183
- MICRO-FEM 234
- microbial contaminants 278
- mineralization in groundwater 96–97
- MOC 234
- modern groundwater 271
 - contamination 271–272, 277–278, *see also* pesticides
 - in groundwater
 - indicators 273–279, 282
 - PALAEAUX 279
- modern groundwater advance 36–38, 279–285
- modern water, infiltration 22–23, 36, 75
- MODFLOW 41, 61, 63, 86, 234

- Netherlands, palaeoclimate 50
- NETPATH 232
- noble gas recharge temperatures 200
 - Aveiro Cretaceous aquifer 151
 - Cambrian–Vendian aquifer 23–25
 - Doñana aquifer 122–123
 - East Midlands Sandstone aquifer 11–12
 - Glatt Valley aquifer 157–160
 - Ribe Formation 41
- non-aqueous phase liquids 278
- North Sea
 - climate change 34, 73
 - geology 30–35
 - palaeogeography 30–32
- North Sea Basin, palaeohydrology 293–297

- offshore freshwater 42, 44, 76
- open aquifer systems 214, 220, *see also* unconfined
 - aquifers

- PALAEAUX programme 7–10
 - hydrochemical evolution 203–205
 - modern groundwater indicators 279
 - palaeoclimate 208
 - palacowater identification 200–203
- palaeoclimatic conditions 265–268
 - Glatt Valley aquifer 157–159
 - Ledo–Paniselian aquifer 50
 - Weichselian 165–176
- palaeoenvironmental model, Ledo–Paniselian aquifer
 - 65–67
- palaeogroundwater formation 25–26
- palaeohydrology
 - eastern Baltic 290–293
 - English Channel 297–300
 - French Atlantic coast 300–302
 - Iberian Atlantic coast 302–306
 - southern North Sea 293–297
 - western Mediterranean coast 306–308
- palacowater
 - definition 2–3
 - exploitation 314
 - identification 200–203
 - investigative techniques 193–200
 - management 313, 318–326
 - legislation 325–326
 - quality 14, 316–318
- permafrost 66–67, 71, 86, 174–176
 - impact on flow systems 223, 225–226
 - Ledo–Paniselian aquifer 225–226
- pesticides in groundwater 282–285
- PHREEQC 59, 232, 234–247
- Pleistocene base level change 251–252, 265–268
- Poole Harbour 82–83, 300
- pore-water salinity 42, 112
- precipitation recharge 35, 50–51, 66
- pumping, effect on groundwater 62–65, 68–69

- radiocarbon analysis 194–196
 - Astian aquifer 103
 - Aveiro Cretaceous aquifer 150–151
 - Dogger aquifer 98
 - East Midlands Sandstone aquifer 10–11
 - Estonia 21, 22
 - Glatt Valley aquifer 157–158
 - Ledo–Paniselian aquifer 52–56
 - Ribe Formation 38, 40–41
 - southern England 88
 - see also* dating tools
- recharge
 - artificially enhanced 111, 323
 - periglacial 17
 - precipitation 35, 50–51, 66, 133
 - river 111–116
- recharge condition indicators 198–200
- Reculver borehole 78, 257–260
- residence time indicators 194–198
- Rhone delta 306–308
- Ribe Formation aquifer 29–44
 - geochemistry 35–36, 40–41
 - geophysical logging 36, 42
 - groundwater age 40–41, 202
 - groundwater flow model 41–42
 - impact of glaciation 225
 - isotope analysis 38
 - modern groundwater 280
 - offshore freshwater 42, 44
 - palaeohydrology 296

- Ribe Formation aquifer *continued*
 radiocarbon analysis 38, 40–41
 recharge 35, 38
 structure 32–33
- Sa Pobla boreholes 265
- S'Albufera wetland 123–130
 boreholes 265
- saline intrusion 17, 74–75, 80–82, 95–97, 99–101,
 111–113, 260–265
- saline water monitoring 253
- salinity, airborne 133
- salinity profiles 221–222
- salinization 33, 104–105
- Scandinavian ice sheet 29, 31
- sea-level change 4–5, 30–34, 35, 50, 65, 72–73, 251–255
 Atlantic coast 183–185
 Baltic 179–182
 English Channel 300
 Iberian Peninsula 182–183, 305–306
 Northwest Europe 177–179
 Weichselian 172–173, 176–185
 western Mediterranean 308
see also groundwater evolution
- semi-confined aquifers 32
- Solent River 177, 299
- solutes 277–278
- South Downs aquifer 79–82
- Spain, coastal aquifers 107–135
- speleothem growth 4, 72, 174–175
- Spilsby Sandstone aquifer 76–77
- St Peter's Church borehole 260–265
- steady state interface 216–222
- subglacial hydrology 173–174
- subglacial infiltration 25–26
- SUTRA 83
- Tertiary marine sediments 32–33
- TOC *see* total organic carbon
- total organic carbon 38
- tracers 273–277
- tunnel valleys 26, 173–174
- unconfined aquifers 251, *see also* open aquifer systems
- unconsolidated aquifers 306–307
- Victoria Gardens borehole 260–265
- Voosi Formation 20
- Voronka aquifer 19–20
- water demand 314–316
- water quality 316–318
- Weichselian
 glaciation 170–173, 225
 palaeoclimate 165–170
 permafrost 174–176
 sea-level change 172–173, 176–185
 subglacial hydrology 173–174
- wells 314, 319, 321, 324
- Western Lawns borehole 253–255
- western Mediterranean coastal aquifers 306–308
- Yorkshire, Cretaceous Chalk aquifer 74–75

Palaeowaters in Coastal Europe: evolution of groundwater since the late Pleistocene

edited by

W. M. Edmunds and C. J. Milne
(British Geological Survey, Wallingford, UK)

Palaeowaters in Coastal Europe contains 17 contributions from an international array of authors. They discuss the history of groundwater evolution during the late Pleistocene in the coastal areas of Europe from the Baltic region to the Iberian peninsula and the Canary Islands. Geochemical and geophysical techniques for evaluating palaeowaters are reviewed. The focus of the book is on changes in the hydrogeological regime during the Quaternary and their impacts on groundwater movement and chemistry in European coastal aquifers.

The work summarized in the papers was carried out by a partnership of European scientists under the auspices of the PALAEAUX project, an EC initiative. Researchers from the fields of hydrogeology, geochemistry, isotope hydrology and Quaternary studies attempted to reconstruct the most probable movement of groundwater in the study area over the past 100 000 years and its response to climatic events of global significance during the last glacial cycle. The results of this work, summarized in this volume, allow a better understanding of the water resources found at and near the coastlines of northern and western Europe. During times of lowered sea level, it appears that groundwaters were replenished to depths greater than occur at the present day. These pristine freshwater reserves are an irreplaceable asset. Their location at coastlines where populations and water demands are high and often seasonal means that they need careful management to avoid over-exploitation or contamination. The inevitable conflicts that this resource management creates are discussed.

Palaeowaters in Coastal Europe: evolution of groundwater since the late Pleistocene will be of interest to Quaternary scientists, hydrogeologists, marine scientists engaged in coastal research and those involved in environmental science and the management of groundwater assets.

- 344 pages
- 17 papers
- 180 illustrations, including colour
- index

Visit our on-line bookshop:
<http://bookshop.geolsoc.org.uk>

Cover illustration: Dorset coast, southern England. The present-day European coastline has existed only for 8000 years. Before this time groundwater flowed 'offshore' and is now preserved at depth in several locations. Water resources in coastal regions of Europe are subject to immense seasonal pressures and the palaeowaters at depth offer one possible alternative for supply. Photograph by D. K. Buckley.

ISBN 1-86239-086-X



9 781862 390867 >