

Increased Levels of Free Fatty Acids in Fasted Mice Stimulate In Vivo β -Cell Electrical Activity

Juana Fernandez and Miguel Valdeolmillos

The electrical activity of pancreatic β -cells in 48-h fasted mice has been recorded in vivo. Their electrical activity is exceedingly high at low levels of blood glucose when compared with control animals. For example, at a blood glucose concentration of 4.5 mmol/l, at which β -cells are permanently hyperpolarized in control animals, fasted animals show continuous spiking activity. In fasted animals, hyperpolarization only occurs at glycemia below 2.2 mmol/l. As in fed animals, the electrical activity in fasted mice can be decreased or suppressed by the injection of diazoxide, indicating the participation of K_{ATP} channels. The treatment of fasted animals with nicotinic acid, an inhibitor of lipolysis, produces a decrease in the levels of free fatty acids (FFAs) and a decrease in electrical activity, thereby restoring the dose-response curve for glucose in fasted animals to values close to those found in fed animals. Conversely, the injection of palmitic acid produces an increase in electrical activity without a change in blood glucose. These results point to FFAs as important regulators of electrical activity during fasting in vivo. They also indicate a dissociation of electrical activity and insulin release in fasted animals, since an increase in electrical activity is not associated with increased insulin secretion. *Diabetes* 47:1707–1712, 1998

The secretion of insulin by pancreatic β -cells is regulated by a variety of factors, among which the most important is glucose. It is accepted that metabolism of the sugar is a necessary step in the chain of events that leads to insulin secretion (1). Glucose metabolism generates diverse coupling factors that modulate the activity of channels involved in the regulation of electrical activity and the function of proteins important for secretion. Among the former, the ATP-dependent potassium channel (K_{ATP}) is regulated by the relative levels of ATP and ADP. The closure of K_{ATP} channels, resulting from an increase in the ATP-to-ADP ratio, leads to membrane depolarization and the appearance of oscillatory electrical activity (2–4). The activation of voltage-dependent calcium channels during depo-

larization leads to an increase in intracellular calcium, which plays a central role in insulin secretion (5). In this scenario, changes in electrical activity are a necessary intermediate step in glucose-induced insulin secretion. In fact, studies in vitro have shown a positive correlation between glucose concentration, oscillatory electrical activity, intracellular calcium oscillations, and pulsatile insulin release (6–8).

Although there is general agreement about the basic features of the fuel hypothesis, there are important aspects that are poorly understood. For instance, the specific signal(s) that links metabolism with the closure of K_{ATP} channels and insulin secretion is not known. In addition to the levels of high energy phosphates, other signals, such as adenine nucleotides (9,10) and GTP-dependent mechanisms (11), have been implicated as important regulatory factors.

The interaction between lipid and glucose metabolism is well known; however, the effects of lipid metabolism on insulin secretion are poorly understood (12). Acyl-CoA esters have been implicated in several aspects of insulin secretion. For example, they have been found to modulate K_{ATP} channels (13) and to modify intracellular calcium handling through an effect in a nonmitochondrial compartment (14). Under normal metabolic conditions, with low levels of free fatty acids (FFAs), glucose metabolism leads to a change in the intracellular lipid profile, consisting of an increase in malonyl-CoA and the suppression of fatty acid oxidation (15–18), which seems to be a key event in glucose-induced insulin secretion. During starvation, the acute phase of insulin secretion in response to glucose is impaired when measured in isolated pancreatic islets (19,20). However, in the whole animal, such a response is not greatly depressed. Recent evidence suggests that elevated levels of FFAs play an important role in the ability of islets to secrete insulin in response to glucose in starved animals (21,22). Thus, the blockade of lipolysis during starvation greatly impairs glucose-induced insulin secretion. Conversely, the exposure of islets to fatty acids results in a shift to the left of the glucose-induced insulin secretion (23). From studies in vitro and in vivo, the effect of fatty acids on β -cell function seems to be dual. Short-term exposure to fatty acids produces a stimulation of glucose-stimulated insulin secretion (16,22,24,25), whereas long-term exposure produces an impairment (26–28). This long-term inhibition of insulin secretion may be related to early forms of NIDDM, where high glycemia and high lipidemia produce overstimulation and, finally, β -cell dysfunction (29–31). The overall emergent picture is that lipids may play an important role in β -cell physiology, probably at different steps along the stimulus secretion coupling chain.

The electrical activity of pancreatic β -cells has been recently recorded in vivo (32–34). Such studies have the

From the Instituto de Neurociencias, Campus de San Juan, San Juan de Alicante, Spain.

Address correspondence and reprint requests to Miguel Valdeolmillos, Universidad Miguel Hernandez, Instituto de Neurociencias, Campus de San Juan. Apdo. correos 18, 03550 San Juan de Alicante, Spain. E-mail: miguel.valdeolmillos@umh.es.

Received for publication 8 April 1998 and accepted in revised form 31 July 1998.

EC_{50} , glycemia at which the cells are in the active phase half of the time; FFA, free fatty acid.




FIG. 1. Representative records of the electrical activity recorded *in vivo* in fed (*A*, label control) and 48-h fasted animals (*B*). The values above each record, denoted by the letter *G*, are the blood glucose concentrations measured during electrical recording. Each record is from a different animal.

advantage of providing insight into factors other than glucose that may be important in the modulation of β -cell activity. The most relevant results from these studies related to the present experiments was the finding that glucose-dependent oscillatory electrical activity is maximally sensitive to changes in blood glucose concentration within the normal range of glycemia, ~ 5 – 8 mmol/l. The half-maximal sensitivity of isolated islets of Langerhans is obtained at 12–15 mmol/l glucose concentration.

The present study was undertaken to investigate the relationship between blood glucose concentration and electrical activity recorded *in vivo* under restricted metabolic conditions. It was expected that in starved animals, which have a lower blood glucose concentration, electrical activity would be proportionally decreased. Contrary to our expectations, we have found that in starved animals there is an overstimulation of electrical activity that is not associated with an increase in insulin secretion. Such overactivity is directly related to the increased levels of FFAs secondary to the fasting state.

RESEARCH DESIGN AND METHODS

The methods used for *in vivo* recording have been described elsewhere (32). Albino mice (8–10 weeks old, 25–35 g wt) were anesthetized by intraperitoneal injection of 90 mg/kg Nembutal. Control animals had free access to food and water, while starved animals were fasted for 48 h. The degree of anesthesia was checked periodically during the experiment by testing cutaneous reflexes. The experiments were carried out according to institutional animal care guidelines. The animals were laparotomized, and the duodenal part of the pancreas was dissected free from the vena cava and the abdominal aorta were cannulated in their caudal parts for solution infusion and blood sample collection, respectively. During the experiment, the animal was laid on its back on a heated bed maintained at 37°C. For the electrical recording, the duodenal part of the pancreas was spread out on top of a platform (6 × 15 mm) covered by a 3-mm layer of Sylgard (Dow Corning, Midland, MI). The platform was attached to a micromanipulator. The pancreas was isolated from peristaltic and respiratory movements by fixing it to the Sylgard with dissection pins, taking care not to affect the blood supply. The islets were impaled with high-resistance glass microelectrodes (100–150 M Ω) filled with a solution of 3 mol/l potassium citrate plus 100 mmol/l potassium chloride, and the electrical activity was recorded with an Axoclamp-2A amplifier (Axon Instruments, Foster City, CA). Unfiltered records were acquired at a frequency of 300 Hz and stored on a microcomputer using Axotape for analysis off-line. Blood samples were collected from the aorta and analyzed for glucose concentration (Beckman glucose analyzer 2; Beckman, Brea, CA), insulin concentration (Coat-A-Count; Diagnostic Products, Los Angeles, CA) and nonesterified fatty acids (FFAs) (half-micro test; Boehringer Mannheim, Mannheim, Germany).

For *in vitro* experiments, islets were isolated from the pancreas by dissection and fixed to the base of a recording chamber. Islets were superfused with a modified Krebs solution containing (in millimoles per liter) 120 NaCl, 5 KCl, 25 NaHCO₃, 2.5 CaCl₂, and 1.1 MgCl₂ and gassed with a mixture of 95% O₂ and 5% CO₂ (pH = 7.4). These experiments were carried out at 37°C with different glucose concentrations. All the reagents were of analytical grade from Sigma (Poole, U.K.) and Merck (Darmstadt, Germany).

RESULTS

Figure 1 shows examples of the electrical activity recorded *in vivo* from fed (Fig. 1*A*) and 48-h fasted (Fig. 1*B*) animals. The numbers above each record correspond to the glycemia measured simultaneously with the electrical recording, which is representative of electrical activity under steady-state conditions. In fed animals, at a blood glucose of 4.5 mmol/l the cell is below threshold and its membrane potential is hyperpolarized. At 5.4 mmol/l glucose, the cell shows oscillations in the electrical activity composed of alternating depolarized and hyperpolarized phases. At a higher blood glucose concentration (6.9 mmol/l), active phases of longer duration and shorter silent phases are seen. Finally, when blood glucose is 7.8 mmol/l, the cell displays continuous activity. This relationship between blood glucose and electrical activity is consistently found in *in vivo* electrical recordings (32).

Figure 1*B* shows that for a given glycemia, the activity is consistently higher in fasted animals compared with the control ones. For example, 4.5 mmol/l blood glucose, a concentration that is subthreshold in control animals, in fasted animals produces maximal activation of the cell. In fasted animals with lower glucose concentrations, 4.1 and 3.7 mmol/l, the electrical activity is progressively decreased. Only glycemias as low as 2.2 mmol/l are associated with cell hyperpolarization.

Figure 2 shows the relationship between blood glucose and the percentage of active phase (the time spent depolarized with respect to the total recording time) for fasted animals. In this figure, each open circle corresponds to a fasted animal recording ($n = 67$), such as those shown in Fig. 1*B*. The data points indicate a very steep relationship between blood glucose and electrical activity. The cells become active at glycemias in the range 2.5–3 mmol/l and show continuous




FIG. 2. Plot of the relationship between glycaemia and percentage of the active phase of electrical activity. Each open symbol represents a different experiment in fasted animals ($n = 67$), like those shown in Fig. 1B. The sigmoid curve, labeled "Fed," shows the fitting obtained in experiments on fed animals, like those shown in Fig. 1A (32). The EC_{50} (glycaemia at which the cells are in the active phase half of the time) for the control curve is 6.76 mmol/l (95% CI 6.59–6.94), and the slope is 5.57 (95% CI 4.67–6.50). A sigmoidal fitting to the data obtained in fasted animals gives an EC_{50} of 3.8 mmol/l (95% CI 3.5–4.1) and a slope of 4.6 (95% CI 2.9–6.2).

electrical activity above 5 mmol/l. In other words, in the narrow range from 2.5 to 5 mmol/l glucose, which are the values of glycemias in our fasted animals, the cells change from no activity to maximal activity. The continuous line labeled "Fed" shows the fitting of data obtained in previous experiments (32) and describes the relationship between blood glucose and electrical activity in fed animals, such as those shown in Fig. 1A. It is clear that in fasted animals, for a given blood glucose concentration, the cells are more active than those in the control situation. Thus, contrary to the expected results that β -cell electrical activity would decrease with the lower glycemias associated with fasting, we observed hyperactivity and a shift of the dose-response curve toward lower glucose levels.

One possible explanation for the hyperactivity observed in fasted animals could be the impairment of cell metabolism due to hypoglycemia. We considered the possibility that the K_{ATP} channels were blocked and unresponsive. To test this possibility, we looked at the effects of the K_{ATP} channel opener diazoxide on electrical activity in fasted animals.

Figure 3A shows the effect of diazoxide on electrical activity in a fasted animal with a glycaemia of 4.1 mmol/l. The injection of diazoxide leads to the transitory suppression of the active phases for a period of ~ 90 s, after which the cell reestablished the oscillatory pattern. Figure 3B shows the effect of diazoxide on a different cell, which showed continuous electrical activity. Again, diazoxide completely hyperpolarized the cell, in this case for a longer period of time. The effect was transitory, and the cell restarted electrical activity, in this case with oscillations of electrical activity. Essentially the same diazoxide modulatory effects have been described in control animals (35). These results suggest, therefore, that in fasted animals, the electrical activity of pancreatic β -cells is controlled by the K_{ATP} channel activity, although with a different glucose sensitivity.




FIG. 3. Effect of diazoxide on the electrical activity in fasted animals. In A, diazoxide was injected (50 μ l of a 15 mmol/l solution) at the time indicated by the arrow into an animal that shows oscillatory electrical activity. B shows the effect of the same diazoxide concentration in an animal with continuous electrical activity. The letter G denotes the time at which a blood sample was taken and the glycaemia measured. The figure is representative of six similar experiments.

The shift in the dose-response curve for glucose found in fasted animals could be due to an intrinsic change in the β -cell sensitivity to glucose. Alternatively, the factor responsible for the shift could be external to the β -cells. If the former is correct, isolated islets from fasted animals should show a different dose-response curve for glucose with respect to islets taken from control animals. To check this, we isolated islets from control and fasted animals and recorded the electrical activity at different glucose concentrations. The time taken from the isolation of the islets and the beginning of the recording ranged from 30 to 60 min. We found in both groups of islets the same dose-response relationship (not shown). These results suggested that there may be a factor in fasted animals whose presence changes β -cell glucose sensitivity.

TABLE 1

Glycaemia, insulin, and FFAs in the different groups of animals studied

	Glycaemia (mmol/l)	Insulin (μ U/ml)	FFAs (mmol/l)
Control	7 \pm 1.2 (73)	10.3 \pm 3.5 (5)	0.2 \pm 0.05 (11)
Fasted	4.7 \pm 0.8 (64)*	5.9 \pm 0.8 (7)†	0.65 \pm 0.37 (11)*
Control + NA	7.4 \pm 1.7 (8)	9.5 \pm 2.8 (5)	0.12 \pm 0.05 (5)
Fasted + NA	4.4 \pm 0.6 (13)*	7.4 \pm 2.5 (7)	0.15 \pm 0.07 (9)

Data are means \pm SD (n). Groups of animals studied include control (fed), 48-h fasted, control treated with nicotinic acid (NA), and fasted treated with NA. * $P < 0.01$; † $P < 0.05$ (fasted compared with control and fasted + NA compared with control + NA).




FIG. 4. Representative records of the electrical activity recorded in 48-h fasted animals (A) and fasted animals treated with nicotinic acid (B). The figure is representative of the effects obtained in 12 animals treated with nicotinic acid. The values above each record, denoted by the letter G, are the blood glucose concentrations measured during electrical recording. Each record is from a different animal.

The most notable change produced during fasting is an increase in circulating FFAs. As shown in Table 1, under our experimental conditions the FFA values rose from 0.2 mmol/l in control animals to 0.65 mmol/l in fasted animals. The treatment of fasted animals with the antilipolytic agent nicotinic acid (36) prevented the rise in FFAs during fasting, whereas it had no effect in control animals (Table 1).

Figure 4 shows the effect of treatment with nicotinic acid in fasted animals. The nicotinic acid treatment consists of daily intraperitoneal injections of 30 μ l (stock 100 mmol/l) of nicotinic acid. Figure 4A shows the electrical activity at two blood glucose concentrations (4.1 and 4.5 mmol/l) in fasted animals. As previously shown (Fig. 1), cells at these glycemia range from moderately activate at 4.1 mmol/l to continuously active at 4.5 mmol/l glucose. Figure 4B shows electrical activity in two animals treated with nicotinic acid. As can be appreciated, for comparable blood glucose concentrations the electrical activity is considerably lower in these animals and is very close to the activity recorded in control nonfasted animals. In fact, the percentage of active phase for a given glycemia in fasted animals treated with nicotinic acid is the same as the one obtained in control animals (represented in Fig. 2 by the continuous line).

In parallel experiments, we checked the effect of nicotinic acid in normally fed animals. As shown in Table 1, nicotinic acid treatment did not produce any significant change in blood glucose or insulin secretion and produced only a small decrease in FFAs. The electrical activity recorded from fed animals treated with nicotinic acid was indistinguishable from that recorded from fed nontreated animals (data not shown).

If the FFA levels are responsible for increased electrical activity in fasted animals, it should be possible to mimic the fasted state by the addition of fatty acids. Figure 5 shows the effect of palmitic acid injection on electrical activity in a fasted and nicotinic acid-treated animal. The initial part of the




FIG. 5. Effect of the injection of palmitate (25 μ l of a 5 mmol/l solution) on the electrical activity in fasted animals treated with nicotinic acid. Palmitate was injected at the end of the first part of the record, at the time indicated by the arrow. The second part of the record shows the electrical activity 15 min after the injection of palmitate. The letter G denotes the time at which a blood sample was taken and the glycemia measured. The figure is representative of six similar experiments.

record shows the electrical activity, which is equivalent to that of a fed animal, since lipolysis was prevented by nicotinic acid. At the time indicated by the arrow, palmitate was injected, leading to a slow increase in electrical activity. At 15 min after the injection, the cell was in continuous electrical activity, although the glucose concentration slightly decreased. In six experiments performed under similar conditions, palmitate produced an increase in the length of the active phase of electrical activity of $111 \pm 94\%$ (mean \pm SD). The glycemia values before and after injection of palmitate were 4.9 ± 0.5 and 4.8 ± 0.5 mmol/l, respectively.

DISCUSSION

The results presented in this paper show that for a given blood glucose concentration, electrical activity of pancreatic β -cells in fasted animals is increased with respect to control animals. This increase in activity seems to be produced by the increase in FFAs that occurs during fasting. These results put forward two important points. First, lipid metabolism exerts an important influence on pancreatic β -cell electrical activity. Second, in fasting animals, there is a dissociation between electrical activity and insulin secretion. Here, increased electrical activity is not associated with an increase in insulin secretion.

Fasting leads to a decrease in blood glucose levels and a corresponding decrease in circulating insulin. Taking into consideration previous electrophysiological recordings in vivo (32,35), a proportional decrease in electrical activity was expected; however, the opposite effect was found. For a given blood glucose concentration, the electrical activity is higher in fasted animals. The overstimulated electrical activity observed in 48-h fasted animals with respect to fed ones is due to a specific metabolic effect, not to impairment of β -cell status or gross alterations in the channels implicated in regulation of membrane potential. As in fed animals, the membrane potential of β -cells in fasted animals can be regulated by pharmacological interventions that modulate K_{ATP} channels. Thus, treatment with the K_{ATP} channel opener diazoxide led to hyperpolarization of the membrane and a decrease in the time spent by the cell in the active phase. Likewise, the K_{ATP} channel blocker tolbutamide increases electrical activity in hyperpolarized cells (data not shown). The same pharmacological modulation has been reported in fed animals (35). On the other hand, islets of Langerhans dissected

from fasted animals show the same glucose–electrical activity relationship as fed ones.

The increased electrical activity in fasted animals is a consequence of increased levels of FFAs. This conclusion is derived from the experiments with nicotinic acid and the effects of palmitic acid infusion. Thus, the effect of fasting on electrical activity can be blocked by treatment with the antilipolytic agent nicotinic acid (36), restoring the relationship between glucose and electrical activity toward the fed type. Conversely, infusion of palmitic acid leads to the apparition of increased electrical activity. The slow time course of the palmitic acid effect may indicate that it is not due to a direct effect on membrane channels, but to an indirect one dependent on metabolism.

It has been recently shown that long-chain acyl-CoA esters produce the activation of K_{ATP} channels (13). Such an effect should lead to a hyperpolarization rather than the increased electrical activity observed here. It is possible that differences in experimental conditions may explain such differences. It is also possible that the effects of lipids on K_{ATP} channels are different in short- and long-term exposures (see below). Further work is necessary to clarify this point.

There is a body of work suggesting the importance of lipid metabolism in β -cell insulin secretion. The effects of lipids have been implicated in the normal physiological response to glucose, as well as that under pathological conditions. Under physiological conditions, glucose metabolism leads to the diversion of fatty acids from the oxidative to the esterification pathway, leading to an increase in fatty acyl-CoA (15–18). This diversion is due to an increase in malonyl-CoA and its inhibitory action on carnitine palmitoyl-transferase I (CPTI). However, the exact mechanism(s) that links the increase in FA-CoA with insulin secretion is not clear and could be direct or mediated by increased pools of diacylglycerol or the effect on other enzymes proximal to insulin secretion. There is also evidence that the effect of fatty acids on insulin secretion depends on the length of exposure. Short-term exposure produces a stimulation (16,22,24,25), whereas long-term exposure produces an inhibition of insulin secretion (26–28). In this respect, the increased lipid levels in 48-h fasting mice has to be considered long-term exposure. Consistent with this, the fasted animals treated with nicotinic acid show a tendency to increase insulin levels (5.9 vs. 7.4 μ U/ml, fasted vs. fasted plus nicotinic acid) despite the fact that the values of glycemia still correspond to those of fasted animals (4.7 vs. 4.4 mmol/l, fasted versus fasted plus nicotinic acid). It has to be kept in mind that the increased insulin secretion in nicotinic acid-treated animals is associated with a decrease in the electrical activity. This further highlights the dissociation between electrical activity and insulin secretion in fasted animals.

The existence of a positive correlation among glucose concentration, electrical activity, and the concomitant increase in intracellular calcium and insulin secretion in models in vitro and in vivo is well documented. This correlation seems to be modified during starvation. Our experiments do not provide information to clarify the steps that may be altered. However, it is interesting to note that in isolated HIT cells, acyl-CoA esters cause an increase in calcium influx into a nonmitochondrial pool, consistent with a stimulatory effect on the endoplasmic reticulum Ca-ATPase. Therefore, it would be interesting to determine if overstimulation of electrical activity in fasted animals is associated with the expected increased

levels of intracellular calcium (37). In the case of a prevailing effect on the Ca-ATPase, the calcium increase due to electrical activity could be at least partially curtailed. In addition, it has been shown that fasted rats show reduced levels of mRNA for the calcium channel (38). Further experiments are necessary to check if the increased electrical activity is associated with increased levels of intracellular calcium.

Previous work has shown that during fasting, the presence of FFAs is necessary for an adequate insulin secretory response to an increase in glucose (21). Treatment with nicotinic acid leads to a decrease in such a response. From our experiments, it can be speculated that in nicotinic acid-treated animals, which have a fed-like glucose–electrical activity relationship, the secretory response to a glucose load would be decreased. This may indicate that in addition to electrical activity, there must be other factors directly implicated in insulin secretion. In any case, the final effects of FFAs are not easily predictable because of the multiplicity of their actions.

The possible participation of elevated levels of fatty acids in initial forms of type 2 diabetes, especially those associated with obesity, has been indicated. The experiments presented here raise the possibility that continuous overstimulation of β -cell electrical activity may lead to a deleterious effect on the β -cell, contributing to the later impairment of insulin secretion.

In conclusion, our results directly demonstrate the participation of lipids in the regulation of pancreatic β -cell electrical activity during fasting. Their effect has proven to be complex and linked to a dissociation in the conventional stimulus-secretion coupling chain in pancreatic β -cells. Among other issues, it remains to be determined whether the increased electrical activity and the depressed insulin secretion are causally related or are two independent manifestations of lipid metabolism.

ACKNOWLEDGMENTS

This work was supported by grants FIS 94/1345 (Fondo de Investigaciones de la Seguridad Social) and SAF 97/0195 (Comision Interministerial de Ciencia y Tecnologia, Spain).

We thank A. Perez-Vergara for technical assistance.

REFERENCES

- Ashcroft FM, Ashcroft SJH: Mechanism of insulin secretion. In *Insulin: Molecular Biology to Pathology*. Ashcroft FM, Ashcroft SJH, Eds. Oxford, U.K., Oxford University Press, 1992, p. 97–150
- Cook DL, Halles N: Intracellular ATP directly blocks K channels in pancreatic β -cells. *Nature* 311:271–273, 1984
- Ashcroft FM, Harrison DE, Ashcroft SJH: Glucose induces closure of single potassium channels in isolated rat pancreatic β -cells. *Nature* 312:446–448, 1984
- Meissner HP: Electrical characteristics of beta-cells in pancreatic islets. *J Physiol Paris* 72:757–767, 1976
- Prentky M, Matschinsky FM: Ca^{2+} , cAMP, and phospholipid-derived messengers in coupling mechanisms of insulin secretion. *Physiol Rev* 67:1185–1248, 1987
- Atwater I, Carroll P, Xu Li M: Electrophysiology of the pancreatic B-cell. In *Molecular and Cellular Biology of Diabetes Mellitus*. Vol 1. Draznin B, Melmed S, Le Roith D, Eds. New York, Alan R. Liss, 1989, p. 49–68
- Santos R, Rosario L, Nadal A, Garcia-Sancho J, Soria B, Valdeolmillos M: Widespread synchronous $[Ca^{2+}]_i$ oscillations due to bursting electrical activity in single pancreatic islets. *Pflügers Archiv* 418:417–422, 1991
- Gilon P, Shepherd R, Henquin J-C: Oscillations of secretion driven by oscillations of cytoplasmic Ca^{2+} as evidenced in single pancreatic islets. *J Biol Chem* 268:22265–22268, 1993
- Dukes ID, McIntyre MS, Mertz RJ, Philipson LH, Roe MW, Spencer B, Worley JF: Dependence on NADH produced during glycolysis for β -cell glucose signalling. *J Biol Chem* 269:10979–10982, 1994
- MacDonald MJ: Feasibility of a mitochondrial pyruvate malate shuttle in pan-

- cretic islets. *J Biol Chem* 270:20051–20058, 1995
1. Komatsu M, Schermerhorn T, Noda M, Straub G, Aizawa T, Sharp G: Augmentation of insulin release by glucose in the absence of extracellular Ca^{2+} . *Diabetes* 46:1928–1938, 1997
 2. Newgard CB, McGarry JD: Metabolic coupling factors in pancreatic β -cell signal transduction. *Annu Rev Biochem* 64:689–719, 1995
 3. Larsson O, Deeney JT, Bränström R, Berggren P-O, Corkey BE: Activation of the ATP-sensitive K channel by long chain Acyl-CoA. *J Biol Chem* 271:10623–10626, 1996
 4. Deeney JT, Tornheim K, Korchak HM, Prentki M, Corkey BE: Acyl CoA esters modulate intracellular calcium handling by permeabilized clonal pancreatic β cells. *J Biol Chem* 267:19840–19845, 1992
 5. Corkey BE, Glennon MC, Chen KS, Deeney JT, Matschinsky FM, Prentky M: A role for malonyl-CoA in glucose-stimulated insulin secretion from clonal pancreatic β -cells. *J Biol Chem* 264:21608–21612, 1989
 6. Prentky M, Vischer S, Glennon MC, Regazzi R, Deeney JT, Corkey BE: Malonyl-CoA and long-chain acyl-CoA esters as metabolic coupling factors in nutrient-induced insulin secretion. *J Biol Chem* 267:5802–5810, 1992
 7. Chen S, Ogawa A, Ohneda M, Unger RH, Foster DW, McGarry DJ: More direct evidence for malonyl-CoA-carnitine palmitoyltransferase I interaction as a key event in pancreatic β -cell signaling. *Diabetes* 43:878–883, 1994
 8. Brun T, Roche E, Assimacopoulos-Jeannet F, Corkey BE, Kim K-H, Prentki M: Evidence for an anaplerotic/malonyl CoA pathway in pancreatic β -cell nutrient signaling. *Diabetes* 45:190–198, 1996
 9. Malaisse WJ, Malisse-Lagae F, Wright PH: Effect of fasting upon insulin secretion in the rat. *Am J Physiol* 213:843–848, 1967
 0. Burch PT, Trus MD, Berner DK, Leontire A, Zawulich KC, Matschinsky FM: Adaptation of glycolytic enzymes: glucose use and insulin release in rat pancreatic islets during fasting and refeeding. *Diabetes* 30:923–928, 1981
 1. Stein DT, Esser V, Stevenson BE, Lane KE, Whiteside JH, Daniels MB, Chen S, McGarry D: Essentiality of circulating fatty acids for glucose-stimulated insulin secretion in the fasted rat. *J Clin Invest* 97:2728–2735, 1996
 2. Stein DT, Stevenson BE, Chester MW, Basit M, Daniels MB, Turley SD, McGarry D: The insulinotropic potency of fatty acids is influenced profoundly by their chain length and degree of saturation. *J Clin Invest* 100:398–403, 1997
 3. Hosokawa H, Corkey BE, Leahy JL: Beta-cell hypersensitivity to glucose following 24-h exposure of rat islets to fatty acids. *Diabetologia* 40:392–397, 1997
 4. Vara E, Tamarit-Rodriguez J: Glucose stimulation of insulin secretion in islets of fed and starved rats and its dependence on lipid metabolism. *Metabolism* 35:266–271, 1986
 5. Warnotte C, Gilon P, Nenquin M, Henquin M: Mechanisms of the stimulation of insulin release by saturated fatty acids: a study of palmitate effects in mouse β -cells. *Diabetes* 43:703–711, 1994
 6. Sako Y, Grill VE: A 48 h lipid infusion in the rat time-dependently inhibits glucose-induced insulin secretion and β -cell oxidation through a process likely coupled to fatty acid oxidation. *Endocrinology* 127:1580–1589, 1990
 7. Elks ML: Chronic perfusion of rat islets with palmitate suppresses glucose-stimulated insulin release. *Endocrinology* 133:208–214, 1993
 8. Zhou YP, Grill VE: Long term exposure to fatty acids and ketones inhibits β -cell function in human pancreatic islets of Langerhans. *J Clin Endocrinol Metab* 80:1584–1590, 1995
 9. Lee Y, Hiroshi H, Ohneda M, Johnson JH, McGarry JD, Hunger RH: β -cell lipotoxicity in the pathogenesis of non-insulin-dependent diabetes mellitus of obese rats: impairment in adipocyte- β -cell relationships. *Proc Natl Acad Sci USA* 91:10878–10882, 1994
 0. Prentki M, Corkey BE: Are the β -cell signaling molecules malonyl-CoA and cytosolic long-chain acyl-CoA implicated in multiple tissue defects of obesity and NIDDM? *Diabetes* 45:273–283, 1996
 1. Boden G: Role of fatty acids in the pathogenesis of insulin resistance and NIDDM. *Diabetes* 46:3–10, 1997
 2. Sanchez-Andres JV, Gomis A, Valdeolmillos M: The electrical activity of pancreatic β -cells recorded "in vivo" shows glucose-dependent oscillations. *J Physiol* 486:223–228, 1995
 3. Valdeolmillos M, Gomis A, Sanchez-Andres JV: In vivo synchronous membrane potential oscillations in mouse pancreatic β -cells: evidence of lack of coordination between different islets of Langerhans. *J Physiology* 493:9–18, 1996
 4. Gomis A, Sanchez-Andres JV, Valdeolmillos M: Oscillatory pattern of electrical activity in mouse pancreatic β -cells recorded in vivo. *Euro J Physiol* 432:510–515, 1996
 5. Gomis A, Valdeolmillos M: Regulation by tolbutamide and diazoxide of the electrical activity in mouse pancreatic β -cells recorded in vivo. *Br J Pharmacol* 123:443–448, 1998
 6. Balasse EO, Ooms HA: Role of plasma free fatty acids in the control of insulin secretion in man. *Diabetologia* 9:145–151, 1973
 7. Fernandez J, Valdeolmillos M: Glucose-dependent Ca oscillations in mouse pancreatic islets of Langerhans recorded in vivo (Abstract). *J Physiol* 491:91P, 1995
 8. Iwashima Y, Kondoh-Abico A, Seino S, Takeda J, Eto M, Polonsky K, Makino I: Reduced levels of messenger ribonucleic acid for calcium channel, glucose transporter-2, and glucokinase are associated with alterations in insulin secretion in fasted rats. *Endocrinology* 135:1010–1017, 1994

Author Queries (please see Q in margin and underlined text)

Q1: Is this the correct affiliation for both authors? If not, please provide affiliations.

Q2: Sentence beginning “Such studies have the...” OK as edited? If not, please reword for clarity.

Q3: Please provide location (city and state) for Dow Corning.

Q4: Please provide location for Axon Instruments.

Q5: Please provide locations for Diagnostic Products and Boehringer Mannheim. Please provide manufacturer and location for the Beckman glucose analyzer.

Q6: Please provide location for Sigma and Merck.

Q7: Correct that you are referring here to the sigmoid curve labeled “Fed” in Fig 2? “fed” OK for “control” here?

Q8: rewording of the definition of EC₅₀ OK?

Q9: “which” meant for “that” here?

Q10: “The values above each record...”: Is this what was meant by “Other details as in Fig. 1”? If not, please provide full figure legend.

Q11: “The letter G...”: Is this what was meant by “Other details as in Fig. 3”? If not, please provide full figure legend.

Q12: To avoid a sentence fragment, rewording of sentence beginning “First, lipid...” OK? If not, please advise.

Q13: Please spell out FA.

Q14: “which have” OK for “with”?

References

Ref. 1: Is Oxford, U.K. the correct place of publication? If not, please provide.

Ref. 37 does not appear to be correct. Please double-check information provided. Is this a one-page article, an abstract, or a letter?