

Use of Titanium Mesh in Lieu of a Fixation Screw to Stabilize an Autogenous Block Graft: A Case Report

Sudhindra Kulkarni, MDS^{1*}
 Srinath Thakur, MDS¹
 Sujatha Kamath, MDS²
 Sampath Kumar, MDS²

Localized alveolar augmentation is carried out either before, during, or sometimes after implant placement. The placement of autogenous graft as a block or a particulate alone or in combination with anorganic bone mineral has been practiced with a great deal of success. The block graft is secured in place with a screw and protected by a membrane. This case report describes the treatment of a female patient with a bucco-lingual bone deficiency grafted with autogenous block and a titanium mesh. Five months later, implant was placed in the grafted area. The definitive prosthesis was cemented 6 months later and followed up for 2 years.

Key Words: *autogenous bone graft, titanium mesh, implant*

INTRODUCTION

The treatment of complete and partial edentulism with implant-retained restorations is now an accepted reality. The root form dental implants and osseointegration are now a proven entity.¹⁻³ The availability of sufficient volumes of bone is one of the prerequisites for successful osseointegration. Conditions like trauma, periodontal disease, periapical cysts, and sometimes anatomic limitations compromise the amount of bone available for implant placement and also influence the final restorative outcome.⁴ Such cases necessitate replacement of the missing teeth by other modes of replacement or augmentation of the alveolar bone.

Localized ridge augmentation has become one of the most common modes of replacement of the lost bone, and it is carried out either before, during, or sometimes after implant placement.⁵⁻¹¹ Bone augmentation is carried out not only to provide the underlying base for implant placement but also to

create harmonious tissue contours in relation to the adjacent teeth.

Studies have shown that the implant survival in regenerated bone is predictable.⁷⁻¹³ Various types of grafts, including autogenous bone as a corticocancellous block or in a particulate form,^{6,14-20} allografts,^{17,21,22} xenografts,^{23,24} and alloplastic¹⁷ materials or a combination of these materials have been used by clinicians, and the overall perception has been that the procedure has been successful,¹¹


Autogenous blocks have been stabilized routinely using a fixation screw holding the block to the recipient bed.^{5,6,16,17,25} Though the use of collagen membrane or expanded polytetrafluoroethylene membrane over the graft is routinely carried out, the problems of membrane collapse, local infection after membrane exposure, and incomplete bone formation in the space provided by the membrane haunt practitioners.^{5,6,26-28}

Titanium mesh provides a unique advantage in that it is malleable, making it adaptable to the underlying graft, which helps it achieve a firm shape and secure the graft in situ. It also prevents the collapse of the overlying tissue into the defect area. The current report presents a case of a localized alveolar defect treated with a corticocancellous block graft stabilized with a titanium mesh.

¹ Department of Periodontics and Implantology, SDM College of Dental Sciences and Hospital, Dharwad, India.

² Department of Prosthodontics and Implantology, SDM College of Dental Sciences and Hospital, Dharwad, India.

* Corresponding author, e-mail: drsudhindrak@gmail.com
 DOI: 10.1563/AAID-JOI-D-11-00006



FIGURES 1–5. **FIGURE 1.** Preoperative view. **FIGURE 2.** Outline of the graft at the donor site. **FIGURE 3.** Titanium mesh tucked under the palatal flap and the recipient bed decorticated. **FIGURE 4.** Titanium mesh contoured and adapted over the graft. **FIGURE 5.** Primary closure attained at the donor and the recipient sites.

CASE REPORT

An 18-year-old female patient was referred to the Department of Implantology for replacement of her missing upper right lateral incisor (tooth #7). The patient’s history revealed trauma at the age of 10 years and avulsion of the lateral incisor, which was treated by reimplantation and splinting of the same tooth. The tooth underwent discoloration and root resorption and was eventually extracted when the patient was 15 years old. For the previous 3 years the patient had been wearing a treatment partial denture but now desired a fixed replacement. Clinical examination revealed discolored tooth #8 and ridge deficiency in the tooth #7 area (Figure 1).

Surgical technique

A midcrestal incision was made in the tooth #7 area, extending mesially and distally to the gingival crevices of the adjacent teeth. A vertical release incision was placed distally, exposing the underlying bone. The recipient bed was de-corticated and shaped to receive a corticocancellous bone graft. Corticocancellous autogenous block graft was harvested from the mandibular parasymphiseal area using piezosurgery (Figure 2).

A titanium mesh screen (Synthes, West Chester, Pa) of 1.0-mm sized pores was cut and shaped to

the defect and tucked under the palatal flap (Figure 3). The block was placed in the recipient area and was stabilized into the defect shaped to receive it. Bovine anorganic bone mineral (Bio-Oss, Geitslich Biomaterials, Wolhusen, Switzerland) was grafted around the block. The mesh was contoured and adapted around the graft, which ensured stabilization of the graft and the mesh. No fixation screws were used to hold the graft or the mesh. The flaps were approximated and closed with 3-0 resorbable sutures (Vicryl, Ethicon, Johnson and Johnson, Aurangabad, India) (Figures 4 through 6). Five months later the site was reentered, and the mesh was removed. It was noticed that the graft was well adapted and integrated into the site and did not display much resorption.

Sequential drilling was carried out, and a 3.5 diameter Nobel Replace Select Tapered (Nobel Biocare AB, Goteborg, Sweden) implant was placed in the osteotomy (Figure 7). The implant was uncovered 4 months later and was temporized for a period of 2 months (Figures 8 through 10). Porcelain fused to metal (PFM) crowns were fabricated and cemented on the implant and the central incisor (Figures 11 and 12). The patient has now been followed up for 2 years. The soft tissue and the bone levels around the implants have been maintained (Figures 13 and 14).

DISCUSSION

Esthetic rehabilitation of missing anterior teeth is a challenging proposition, and the unavailability of adequate bone volume at the site adds to the challenge. Though bone grafting procedures around dental implants have been shown to be predictable, no single procedure has been shown to be better than the other. Autogenous bone has been said to be the gold standard because of its osteogenic potential and compatibility.¹⁷


The graft harvested from the mandibular parasymphysis has shown good clinical outcomes.^{6,18,19,27} The block has usually been fixed with mini screws to secure it in place so as to prevent fibrous growth into the graft, which could lead to failure of integration of the graft.^{16,25,26}

The decortications of the recipient site with perforations created an ideal bed for the placement of the corticocancellous autogenous block, which was then secured with the help of a titanium mesh. Once contoured, the mesh stays as shaped and remains firm.^{22,24,28-30} The mesh was well contoured and adapted to the graft, thus avoiding the placement of a fixation screw. However, in most studies, the mesh has been stabilized with screws.^{24,28-30} This is probably the first case report wherein the block graft is secured by the mesh only and without the use of the fixation screws. This, however, does not mean that even in large defects one can avoid the use of a screw to stabilize a block or the mesh. The fact that stabilization of the graft at the recipient site is the key to success and has to be achieved irrespective of the mode of fixation cannot be undermined.

In the present report, no membrane was placed over the mesh, as it has been observed that a pseudo-periosteum forms around the mesh, which may act as a protective barrier.³⁰

Graft resorption is a very common occurrence, irrespective of the type of the graft or coverage by a membrane.³¹ In the present patient no significant resorption of the block was noticed. Studies have shown that whether or not a membrane is used, there is some amount of graft resorption, which necessitates overcorrection. So the use of a membrane is not mandatory as done in the present case.

The nonplacement of the fixation screw to hold the graft might have been one of the reasons there was little resorption of the graft as it avoided an



FIGURES 6–8. **FIGURE 6.** Radiograph showing the mesh and the graft. **FIGURE 7.** Implant placed at the grafted sites 5 months after the surgery. **FIGURE 8.** Second-stage procedure done and the final abutment placed.



FIGURES 9–14. **FIGURE 9.** Temporary restoration placed. **FIGURE 10.** Well-healed tissues before cementation. **FIGURE 11.** Final restorations placed. **FIGURE 12.** Radiographic image immediately after the cementation of the final restorations. **FIGURE 13.** Clinical picture at the 2-year follow-up. **FIGURE 14.** Radiographic image 2 years after the restorations.

additional trauma to the block caused by drilling; however, more studies are needed to establish this as a fact and not a one-time occurrence.

The temporary restoration placed at the second-stage surgery provided ideal soft tissue contours over a 2-month period, and thus, an ideal emergence profile was achieved. The final restorative outcome was very well accepted by the patient.

In conclusion, it can be stated that titanium mesh can be used to secure the autogenous block graft and result in predictable graft integration.

ABBREVIATION

PEM: porcelain fused to metal

REFERENCES

1. Adell R, Eriksson B, Lekholm U, Branemark PI, Jemt T. Long-term follow-up study of osseointegrated implants in the treatment of totally edentulous jaw. *Int J Oral Maxillofac Implants.* 1990;85: 347–359.
2. Lekholm U, van Steenberghe D, Herrmann I, et al. Osseointegrated implants in the treatment of partially edentulous

- jaws. A prospective 5-year multicentre study. *Int J Oral Maxillofac Implants.* 1994;9:627–635.
3. Buser D, Mericske-Stern R, Bernard JP, et al. Long term evaluation of non-submerged ITI Implants. Part I: 8 year life table analysis of a prospective multicentre study with 2359 implants. *Clin Oral Implants Res.* 1997;8:161–172.
4. van Arx T, Cochran DL, Hermann JS, Schenk RK, Higginbottom FL, Buser D. Lateral ridge augmentation and Implant placement. An experimental study evaluating implant osseointegration in different augmentation materials in the canine mandible. *Int J Oral Maxillofac Implants.* 2001;16:343–354.
5. Misch CM, Misch CE, Resnik RR, Ismail YH. Reconstruction of maxillary alveolar defects with mandibular symphysis grafts for dental implants. A preliminary procedural report. *Int J Oral Maxillofac Implants.* 1992;7:360–366.
6. Urbani G, Lombardo G, Santi E, Tarnow D. Localised ridge augmentation with chin grafts and resorbable pins. Case reports. *Int J Periodontics Restorative Dent.* 1998;18:363–375.
7. Nevins M, Mellonig JT, Clem DS, Reiser GM, Buser DA. Implants in regenerated bone: long term survival. *Int J Periodontics Restorative Dent.* 1998;18:35–48.
8. Dahlin C, Linde A, Gottlow J, Nyman S. Healing of bone defects by guided tissue regeneration. *Plast Reconstr Surg.* 1988;81: 672–676.
9. Dahlin C, Sennerby L, Lekholm U, Linde A, Nyman S. Generation of new bone around dental implants using a membrane technique: an experimental study in rabbits. *Int J Oral Maxillofac Implants.* 1989;4:19–25.
10. Jovanovic S, Spiekermann H, Richter EJ. Bone regeneration around dental implants in dehiscence defect sites. A clinical study. *Int J Oral Maxillofac Implants.* 1992;7:233–245.

11. Aghaloo T, Moy P. Which hard tissue augmentation techniques are the most successful in furnishing bony support for implant placement. *Int J Oral Maxillofac Implants.* 2007; 22(suppl):49–70
12. Simion M, Jovanovich SA, Tinti C, Benfenati SP. Long term evaluation of osseointegrated implants inserted at the time or after vertical ridge augmentation. A retrospective study on 123 implants with 1–5 year follow up. *Clin Oral Implant Res.* 2001;12:35–45
13. Zitzmann NU, Scharer P, Marinello CP. Long term results of implants treated with guided bone regeneration: a 5-year prospective study. *Int J Oral Maxillofac Implants.* 2001; 16:355–366.
14. Misch CM. Comparison of intraoral donor sites for onlay grafting prior to implant placement. *Int J Oral Maxillofac Implants.* 1997;12:767–776.
15. Goldenberg V, Stevenson S. Natural history of autografts and allografts. *Clin Orthop Relat Res.* 1987;225:7–16.
16. Esposito M, Grusovin MG, Coulthard P, Worthington HV. The efficacy of various bone augmentation procedures for dental implants: a Cochrane systematic review of randomised controlled clinical trials. *Int J Oral Maxillofac Implants.* 2006;5:696–710.
17. Hallman M, Mordenfeld A, Strandkvist T. Bone replacement following dental trauma prior to implant surgery-present status. *Dent Traumatol.* 2009;25:2–11
18. Cordaro L, Amadé DS, Cordaro M. Clinical results of alveolar ridge augmentation with mandibular block bone grafts in partially edentulous patients prior to implant placement. *Clin Oral Implants Res.* 2002;13:103–111.
19. Balaji SM. Management of deficient anterior maxillary alveolus with mandibular parasymphiseal bone graft for implants. *Implant Dent.* 2002;11:363–369.
20. Kent JN, Quinn JH, Zide MF, Guerra LR, Boyne PJ. Alveolar ridge augmentation using non-resorbable hydroxyapatite with or without autogenous cancellous bone. *J Oral Maxillofac Surg.* 1983; 41:629–642.
21. Fonseca RJ, Nelson JF, Clark PJ, Frost DE, Olson RA. Revascularization and healing of onlay particulate allogenic bone grafts in primates. *J Oral Maxillofac Surg.* 1983;41:153–162.
22. van Arx T, Kurt B. The TIME technique. A new method for localized alveolar ridge augmentation prior to placement of dental implants. *Int J Oral Maxillofac Implants.* 1996;11:387–394.
23. Skoglund A, Hising P, Young C. A clinical and histological examination in humans of the osseous response to implant natural bone mineral. *Int J Oral Maxillofac Implants.* 1997;12:194–199.
24. Proussaefs P, Lozada J, Kleinman A, Rohrer MD, McMillan PJ. The use of titanium mesh in conjunction with autogenous bone graft and inorganic bovine bone mineral (Bio-Oss) for localized alveolar ridge augmentation. A human study. *Int J Periodontics Restorative Dent.* 2003;23:185–195.
25. Phillips JH, Rahn BA. Fixation effects on membranous and endochondral onlay bone graft revascularisation and bone deposition. *Plast Reconstr Surg.* 1990;6:891–897
26. Von Arx T, Buser D. Horizontal ridge augmentation using autogenous block grafts and the guided bone regeneration technique with collagen membranes: a clinical study with 42 patients. *Clin Oral Implant Res.* 2006;17:359–366
27. Buser D, Dula K, Hirt HP, Schenk RK. Lateral ridge augmentation using autografts and barrier membranes. A clinical study with 40 partially edentulous patients. *J Oral Maxillofac Surg.* 1996;54:420–432.
28. Lozada JL, Proussaefs PT. Clinical radiographic and histologic evaluation of maxillary bone reconstruction by using a titanium mesh and autogenous iliac graft. A case report. *J Oral Implantol.* 2002;28:9–14.
29. Malchiodi L, Scarano A, Quaranta M, Piattelli A. Rigid fixation by means of titanium mesh in edentulous ridge expansion for horizontal ridge augmentation in the maxilla. *Int J Oral Maxillofac Implants.* 1998;13:701–705.
30. von Arx T, Kurt B. Implant placement and simultaneous ridge augmentation using autogenous bone and microtitanium mesh. A prospective clinical study with 20 implants. *Clin Oral Implants Res.* 1999;10:24–33.
31. Chiapasco M, Abati S, Romeo E, Vogel G. Clinical outcome of autogenous bone blocks or guided bone regeneration with e-PTFE membranes for the reconstruction of narrow edentulous ridges. *Clin Oral Implants Res.* 1999;10:278–288.