

Immediate Loading of Maxillary and Mandibular Implant-Supported Fixed Complete Dentures: A 1- to 10-Year Retrospective Study

Ting-Jen Ji, DDS, MSD^{1*}
 Joseph Y. K. Kan, DDS, MS²
 Kitichai Rungcharassaeng, DDS, MS³
 Phillip Roe, DDS, MS²
 Jaime L. Lozada, DMD¹

This study evaluated the success rates of 50 full-arch maxillary and/or mandibular implant-supported fixed complete dentures. After a mean follow-up time of 42.1 months, 269 implants remained in function, which corresponded to cumulative implant success rates of 85.2% and an absolute success rate of 90.6% (269/297 implants). This study suggested that higher implant failure rates might be associated with a dental history of bruxism (29.3%) vs no history of bruxism (4.6%) and surgeons with limited experience (≤ 5 years; 12.2%) vs surgeons with experience (2.4%).

Key Words: *implant-supported fixed complete denture, marginal bone change, immediate loading, surgical experience, immediate implant placement, bruxism*

INTRODUCTION

Prosthetic rehabilitation with an implant-supported fixed complete denture (ISFCD) has been well documented and is a predictable treatment modality for edentulous patients.¹ Traditional guidelines for successful osseointegration have included a healing period of 3 to 6 months without functional loading.^{2,3} However, the use of an interim removable prosthesis during the course of treatment can be an inconvenience to patients.⁴ Recently, the immediate loading of ISFCDs was advocated, and comparable success rates have been reported.⁵⁻¹⁴ The placement of a provisional prosthesis immediately after implant placement can

provide immediate esthetics and function, decrease the number of patient visits, and reduce morbidity of a second surgical intervention.^{4,15}

This retrospective study evaluated the implant success rates of immediately loaded maxillary and/or mandibular ISFCD. Factors affecting implant success rates and prosthetic survival rates were also assessed.

MATERIALS AND METHODS

This study was approved by the institutional review board of Loma Linda University and was conducted in the Center for Implant Dentistry, Loma Linda University School of Dentistry, Loma Linda, California. Treatment records were reviewed for patients who received a full-arch maxillary and/or mandibular ISFCD between January 1998 and December 2008. Patients were included if periapical radiographs were available from the time of surgery (with the attached prosthesis; T0) and at the last follow-up (T1). Implant positions were represented

¹ Advanced Education in Implant Dentistry, Loma Linda University School of Dentistry.

² Department of Restorative Dentistry, Loma Linda University School of Dentistry.

³ Department of Orthodontics and Dentofacial Orthopedics, Loma Linda University School of Dentistry.

* Corresponding author, e-mail: tingjen_ji@hotmail.com

DOI: 10.1563/AAID-JOI-D-11-00027

by their corresponding tooth number. Patient demographics (age and gender), date of implant placement, and date of the final follow-up appointment were recorded.

Implant success was evaluated using a modification of criteria proposed by Spiekermann and Jansen¹⁶ where applicable:

- No implant loss.
- No mesial or distal vertical bone loss >4 mm as assessed on periapical radiograph.

Marginal bone level (MBL) was measured using periapical radiographs taken immediately after implant surgery (T0) and at the last follow-up examination (T1). The apical corner of the implant neck was used as the reference line (RL) (Figure). The marginal bone level was the distance between the RL and the implant-bone contact.¹⁷ The value was zero when the implant-bone contact point was at or more coronal to the RL, and negative when the implant-bone contact was more apical to the RL. Measurements were made on the mesial and distal aspects of each implant to the nearest 1 mm, and the MBL changes between T0 and T1 were calculated. A marginal bone loss greater than 4 mm at any individual implant site was indicative of a failure.

The intraexaminer reliability of the MBL measurements was determined by using double assessments of MBL measured 3 months apart by one examiner (T.J.) and expressed as the intraclass correlation coefficient. The number for marginal bone level measurements made in this study was 0.99.

Prosthesis failure included any provisional or definitive prostheses that were deemed nonfunctional and removed because of extensive implant loss. Implant failure rates were determined for the following categories:

- Type of prosthesis: provisional or definitive prosthesis
- Surgeons' surgical experience: >5 years or ≤5 years
- Surgical site: implant placement in previously grafted healed sites (socket preservation, sinus graft, socket preservation + sinus graft, guide bone regeneration (GBR) + sinus graft, iliac crest block + GBR + sinus graft or Ti-Mesh GBR + sinus




FIGURE. Measurement of marginal bone level (MBL). Reference line (RL) is at the apical corner of the implant neck. Zero and positive values were given when the MBLs were at or coronal and apical to the RL. Negative values were given when the MBLs were apical to the RL.

graft), implant placement in non-grafted healed sites, or immediate implant placement

- Smoking
- Diabetes
- Bruxism
- Implant locations: maxillary anterior (MxA), maxillary posterior (MxP), mandibular anterior (MdA), or mandibular posterior (MdP)
- Implant diameter: 3.0 mm to 3.5 mm, 3.6 mm to 4.5 mm, or 4.6 mm to 5.0 mm
- Implant length: 8.0 mm to 10.0 mm, 10.1 mm to 14.0 mm, or 14.1 mm to 16.0 mm
- Opposing dentition: natural dentition, implant-tissue-supported overdenture, implant-supported overdenture, ISFCD, complete denture, natural dentition with removable partial denture, or full-arch implant-supported fixed partial denture.

Prosthetic complications included any provisional or definitive prosthesis, such as acrylic resin base fracture, broken denture teeth, screw loosening, screw fracture, and/or framework misfit. Framework misfit was assessed at the time of prosthesis placement using a panoramic radiograph.

STATISTICAL ANALYSIS

Life table analysis was used to estimate the progress of implant success over time. The implant failure rates associated with each recorded parameter were represented using descriptive statistics.

TABLE 1
Life table analysis

Time Interval (months)	Total No. of Implants	No. of Implant Successes	No. of Implant Failures	Absolute Implant Success Rate	Cumulative Implant Success Rate
0–12	284	284	13	0.956	0.956
13–24	241	241	0	1.000	0.956
25–36	204	194	10	0.951	0.909
37–48	111	109	2	0.982	0.892
49–60	67	64	3	0.955	0.852
61–72	40	40	0	1.000	0.852
73–84	18	18	0	1.000	0.852
85–96	6	6	0	1.000	0.852
97–108	6	6	0	1.000	0.852
109–126	6	6	0	1.000	0.852
Overall (0–126)	297	269	28	0.906	0.852

RESULTS

A total of 45 patients received 50 full-arch immediately loaded maxillary and/or mandibular ISFCDs. The subject population comprised 18 male and 27 female patients between the ages of 25 and 88 years (mean age = 61.5 years). Eight patients were smokers and one patient had diabetes. Five patients were treated with a maxillary and mandibular ISFCD, and the remaining 40 patients received a single-arch ISFCD. A total of 297 implants were placed in 50 jaws (147 implants in 21 maxillary arches, 150 implants in 29 mandibular arches). This included 233 TiUnite (168 NobelReplace Tapered Groovy; 38 Replace Select Tapered; 27 NobelSpeedy Replace, Nobel Biocare, Yorba Linda, Calif), 54 HA-coated (7 Steri-Oss Hex-Loc, Nobel Biocare; 47 Tapered Screw-Vent, Zimmer Dental, Carlsbad, Calif), and 10 grit-blasted, acid-etched threaded (10 Xive, Dentsply Friadent, Mannheim, Germany) implants.

After a mean follow-up time of 42.1 months (range = 1 to 125.5 months), 269 implants remained in function with <4 mm of MBL change corresponding to the cumulative implant success rates of 85.2% and absolute success rate of 90.6% (269/297) (Table 1).

Tables 1 through 10 display the number and percentage of failed implants in relation to the various recorded parameters.

DISCUSSION

Short- and long-term studies involving full-arch immediately loaded maxillary and/or mandibular ISFCDs have reported success/survival rates of 92.7% to 100.0%.^{5–14} Comparatively, a lower cumulative implant success rate of 85.2% and an absolute success rate of 90.6% (269 of 297) (Table 1) were observed after a mean follow-up time of 42.1 months (range = 1 to 125.5 months) in this study.

In this study, 15 of 81 implants (18.5%) supporting provisional prostheses and 13 of 216 implants (6.0%) supporting definitive prostheses failed (Table 2). It is interesting to note that similar implant failure rates were observed when comparing all resin (18.3%) and metal-resin (19.0%) provisional prostheses. Regardless of the incidence of implant failure, no prosthesis failure was observed in this study.

The experience of the surgeon and the complexity of the surgical procedure have a strong

TABLE 2
Implant failures during provisionalization vs after definitive prosthesis placement

	No. of Implants	Implant Failures		No. of Prostheses	Prostheses Failures	
		No.	%		No.	%
Provisional prosthesis	81	15	18.5	11	0	0
All resin prosthesis	60	11	18.3	8	0	0
Metal-resin prosthesis	21	4	19.0	3	0	0
Definitive prosthesis	216	13	6.0	39	0	0

TABLE 3
Implant failures according to surgeons' experience

Surgeons' Experience	No. of Implants	Implant Failures	
		No.	%
>5 years	85	2	2.4
≤5 years	212	26	12.2

influence on future implant success.^{18,19} Lambert et al¹⁸ found that implants placed by surgeons with less experience (<50 implants) failed twice as often as those placed by surgeons with more experience (>50 implants). Similarly, in this study, the implant failure rate for 2 surgeons with >5 years of surgical experience was 2.4% (2 of 85 implants), whereas the remaining 18 surgeons, that is, those with ≤5 years of surgical experience, incurred an implant failure rate of 12.2% (26 of 212 implants) (Table 3).

It has been suggested that immediately loaded implants placed in fresh extraction sites pose a high risk for implant failure compared with those placed in healed sites.²⁰⁻²² This can result from reduced implant stability as extraction sites do not allow the implant to engage bone circumferentially throughout the length of the implant and/or the presence of residual infection from the failing tooth.²³⁻²⁷ De Bruyn and Collaert²⁷ reported significantly higher implant failure rates (39%) in extraction sites compared with healed sites (0.7%). Higher implant failure rates were observed in this study in the immediate implant placement groups (7/31 = 22.5%) compared with non-grafted healed-site

groups (14/180 = 7.8%) and the grafted-healed site groups (7/86 = 8.1%) (Table 4).

Various augmentation techniques and materials have been used to regenerate the alveolar process in preparation for implant placement.^{28,29} Furthermore, implants placed in augmented sites have been reported to have reasonable success and predictability.³⁰⁻³⁶ In this study, a high implant failure rate (3/6 = 50%) was observed in the iliac crest block + GBR + sinus graft group (Table 4). It could be speculated that bone maturation could be a contributing factor for high failure rates.²⁸

The effects of patient-related risk factors and their influence on implant success and failure have been evaluated.³⁷⁻⁴¹ Moy et al³⁸ reported that smoking, diabetes, head and neck radiation, and postmenopausal estrogen therapy were correlated with an increased rate of implant failure. Similarly, Goodacre et al³⁷ reported a high incidence of implant failure associated with patients with diabetes (9%) and smokers (11%). In this study, a higher implant failure rate was observed in patients with diabetes (28.6%) vs patients who did not have diabetes (8.5%), and comparable failure rates were observed in both smokers (9.7%) and nonsmokers (9.4%). These results differ from those of previous studies that reported significantly higher implant failure rates in smokers (11%) than in nonsmokers (5%) (Table 5).^{37,39}

It has been reported that bruxism may cause unfavorable occlusal forces to implants, ultimately leading to bone loss and implants failures.⁴² In a recent review of the literature, Lobbezoo et al⁴³

TABLE 4
Implant failures according to surgical procedure*

Surgical Procedure	No. of Implants	Implant Failures	
		No.	%
Implant placement in previously grafted healed sites	86	7	8.1
Socket preservation only†	7	0	0
Sinus graft only†	16	2	12.5
GBR only†	0	0	0
Socket preservation + sinus graft†	8	0	0
GBR + sinus graft†	42	2	4.8
Iliac crest block + GBR + sinus graft†	6	3	50.0
Ti-Mesh GBR + Sinus graft†	7	0	0
Implant placement in non-grafted healed sites	180	14	7.8
Immediate implant placement	31	7	22.5
Total	297	28	9.4

*GBR indicates guide bone regeneration; Ti, Titanium.
†Subset of implant placement in grafted healed site.

TABLE 5

Implant failures according to patient's habits and medical and dental status

		No. of Implants	Implant Failures	
			No.	%
Smoking status	Yes	62	6	9.7
	No	235	22	9.4
Diabetes	Yes	14	4	28.6
	No	283	24	8.5
Bruxism	Yes	58	17	29.3
	No	239	11	4.6
Total		297	28	9.4

stated that although there is insufficient evidence to support or refute the relationship between bruxism and implant failure, a careful approach is still recommended. Guidelines to minimize implant failures in patients with bruxism have included increasing the number of implants, improving the occlusion scheme, and incorporating an occlusal stabilization splint after treatment.⁴³ In this study, a higher implant failure rate was reported with patients with bruxism (29.3%) compared with patients without bruxism (4.6%) (Table 5). Because of the nature of this study, it was difficult to evaluate whether these guidelines had been applied during patient treatment. The results from this study suggested that patients with a history of bruxism might be contraindicated for immediately loaded ISFCDs.

With respect to bone quality, studies have shown that implant failure rates are higher in the maxilla than in the mandible, and the area of the lowest failure rate is in the anterior mandible and the highest failure rate is in the posterior maxilla.^{37,44-46} Although the implant failure rates in this study were highest in the MxP region (11.4%), the differences, compared with the failure rates at other regions (7.4%, 9.2%, and 9.5% for MxA, MdA, and MdP regions, respectively; Table 6), were not as great as previously reported in the literature.^{37,44,45} Furthermore, the higher implant failure rate in MxP reported in this study may be attributed to the fact that most of these implants were placed in grafted sites (7/9). These results suggest that the implant prognosis should not be made solely according to the implant location in relation to different sextant of the mouth.

The use of short implants (≤ 10 mm) has been controversial.⁴⁷⁻⁵⁵ Some studies have reported higher failure rates associated with short implants (≤ 10 mm),^{37,47-50} whereas others have reported

TABLE 6

Implant failures according to implant location*

Implant Location	No. of Implants	Implant Failures	
		No.	%
MxA	68	5	7.4
MxP	79	9	11.4
MdA	76	7	9.2
MdP	74	7	9.5
Total	297	28	9.4

*MxA indicates maxillary anterior; MxP, maxillary posterior; MdA, mandibular anterior; MdP, mandibular posterior.

good success.⁵¹⁻⁵⁴ Potential factors contributing to the failure of short implants include implant surface,⁵⁴ implant geometry,⁵⁵ bone quality,^{51,52} and biomechanical stress.⁵⁴ When these factors are appropriately selected and managed, a predictable outcome can be achieved with short implants.^{53,54} Therefore, implant length may not be a primary factor to distribute prosthetic loading to the bone-implant interface.⁵⁴ In this study, it is interesting to note that implants with a length of 14-16 mm had a higher failure rate (20.4%) than the shorter implants (Table 7). However, the failures in the group with implant lengths of 14-16 mm were all distributed among 4 patients with bruxism. Consequently, the implant failures appeared to be affected by patient-related risk factors (bruxism) rather than implant-related risk factors (implant length).

Finite-element analysis studies have suggested that implant diameter has a more significant impact on stress distribution than implant length.⁵⁶ Increasing implant diameter appears to reduce strain to the surrounding crestal bone, thereby preventing further bone remodeling.⁵⁷ The results of this study

TABLE 7

Implant failures according to the implant diameter and length

Implant Size (mm)	No. of Implants	Implant Failures	
		No.	%
Diameter			
3.0-3.5 mm	34	6	17.6
3.6-4.5 mm	232	20	8.6
4.6-5.0 mm	31	2	6.5
Length			
8.0-10.0 mm	62	4	6.5
10.1-14.0 mm	186	14	7.5
14.1-16.0 mm	49	10	20.4

TABLE 8
Implant failures according to the opposing dentition

Opposing Dentition	No. of Implants	Implant Failures	
		No.	%
Natural dentition	133	18	13.5
Implant-tissue-supported overdenture	8	0	0
Implant-supported overdenture	5	0	0
Implant-supported fixed complete denture	108	7	6.5
Implant-supported full-arch fixed partial denture	8	2	25.0
Complete denture	19	1	5.3
Natural dentition with removable partial denture	16	0	0
Total	297	28	9.4

indicate that implant failures were inversely correlated to the implant diameter. (Table 7) Therefore, when it is possible, the use of implants with a diameter ≤ 3.5 mm should be avoided for immediate-loading situations.

It has been reported in the literature that the maximum bite force associated with natural teeth and/or an implant-supported prosthesis are higher than with a removable prosthesis.⁵⁸ The general agreement seems to indicate a correlation between high masticatory forces and increased rates of implant failure.⁵⁹ This concurs with the results of this study, where a higher implant failure rate (27/262 = 10.9%; Table 8) was observed when the ISFCD was opposing natural teeth and/or an implant-supported prosthesis rather than when it was opposing a removable prosthesis (1/35 = 2.1%; Table 8). It should be noted that proper management of the occlusal scheme and occlusal contacts of the prosthesis may influence treatment outcome; however, these factors were not accounted for in this study.

Prosthetic-related complications, such as acrylic resin fracture, screw loosening, screw fracture, and

prosthetic misfit, are common and have been reported in the literature.⁶⁰⁻⁶² In this study, similar incidence rates of prosthetic complications were observed in most categories for both provisional and definitive prostheses (Tables 9 and 10). The only exception was the incidence of prosthesis misfit, which occurred at the highest frequency among all complications during the provisional stage (24%; Table 9) and the lowest in definitive prosthesis (0%; Table 10). Of the 12 provisional prostheses that experienced misfit, 27% (9/33) of these implants failed. It is important to note that a small number of the prostheses placed immediately after implant placement were definitive. Therefore, special care should be taken when fabricating and fitting the prosthesis (provisional or definitive) when performing an immediately loaded ISFCD procedure.

CONCLUSIONS

After a mean follow-up period of 42.1 months, the cumulative implant success rate was 85.2%, with an

TABLE 9
Incidence of prosthetic complications with interim prosthesis

Prosthetic Complication	No. of Prostheses (No. of Incidences)/ No. of Prostheses		Incidence Rate (%)
	Misfit	12 (12)/50	
Acrylic resin base fracture	8 (20)/50	16.0	
Broken denture teeth	6 (15)/50	12.0	
Screw loosening	2 (2)/50	4.0	
Screw fracture	0 (0)/50	0	
Total	18* (49)/50	36.0	

*Indicates that more than one prosthetic complication occurred in one prosthesis.

TABLE 10
Incidence of prosthetic complications with definitive prosthesis

Prosthetic Complication	No. of Prostheses (No. of Incidence)/ No. of Prostheses		Incidence Rate (%)
	Broken denture teeth	5 (9)/35	
Acrylic resin base fracture	3 (5)/35	8.6	
Screw fracture	1 (1)/35	2.9	
Screw loosening	0 (0)/35	0	
Misfit	0 (0)/35	0	
Total	6* (15)/35	17.1	

*Indicates that more than one prosthetic complication occurred in one prosthesis.

absolute success rate of 90.6%. Within the confines of this retrospective study, the results of this study suggest that higher implant failure rates might be associated with patients reporting a history of bruxism (29.3%) vs those without bruxism (4.6%) and surgeons with limited experience (≤ 5 years; 12.2%) vs experienced surgeons (2.4%). However, further controlled studies are needed for more meaningful and definitive conclusions.

ABBREVIATIONS

GBR: guide bone regeneration
 ISFCD: implant-supported fixed complete denture
 MBL: marginal bone level
 MdA: mandibular anterior
 MdP: mandibular posterior
 MxA: maxillary anterior
 MxP: maxillary posterior
 RL: reference line

ACKNOWLEDGMENTS

The authors would like to thank the residents and faculty members at the Loma Linda University School of Dentistry, Center for Implant Dentistry, for their involvement with patient care and Lie Hong Chen for her assistance in the statistical analysis.

NOTE

This manuscript was in partial fulfillment of an MSD degree for Ting-Jen Ji, DDS, MSD.

REFERENCES

- Adell R, Eriksson B, Lekholm U, Brånemark PI, Jemt T. Long-term follow-up study of osseointegrated implants in the treatment of totally edentulous jaws. *Int J Oral Maxillofac Implants.* 1990;5:347–359.
- Adell R, Lekholm U, Rockler B, Brånemark PIA. 15-year study of osseointegrated implants in the treatments of the edentulous jaw. *Int J Oral Surg.* 1981;10:387–416.
- Brånemark PI. Osseointegration and its experimental background. *J Prosthet Dent.* 1983;50:399–410.
- Chee W, Jivraj S. Efficiency of immediately loaded mandibular full-arch implant restorations. *Clin Implant Dent Relat Res.* 2003;5:52–56.
- Schnitman PA, Wohrle PS, Rubenstein JE, DaSilva JD, Wang NH. Ten-year results for Brånemark implants immediately loaded with fixed prostheses at implant placement. *Int J Oral Maxillofac Implants.* 1997;12:495–503.
- Tarnow DP, Emtiaz S, Classi A. Immediate loading of threaded implants at stage 1 surgery in edentulous arches: ten consecutive case reports with 1- to 5-year data. *Int J Oral Maxillofac Implants.* 1997;12:319–324.
- Jaffin RA, Kumar A, Berman CL. Immediate loading of implants in partially and fully edentulous jaws: a series of 27 case reports. *J Periodontol.* 2000;71:833–838.
- Bashi SF, Wolfinger GJ, Balshi TJ. A prospective study of immediate functional loading, following the Teeth in a Day protocol: a case series of 55 consecutive edentulous maxillas. *Clin Implant Dent Relat Res.* 2005;7:24–31.
- Brånemark PI, Engstrand P, Ohrenell LO, et al. Brånemark Novum: a new treatment concept for rehabilitation of the edentulous mandible. Preliminary results from a prospective clinical follow-up study. *Clin Implant Dent Relat Res.* 1999;1:2–16.
- Ganeles J, Rosenberg MM, Holt RL, Reichman LH. Immediate loading of implants with fixed restorations in the completely edentulous mandible: report of 27 patients from a private practice. *Int J Oral Maxillofac Implants.* 2001;16:418–426.
- Grunder U. Immediate functional loading of immediate implants placed in edentulous arches: 2-year results. *Int J Periodont Restor Dent.* 2001;21:545–551.
- Collaert B, De Bruyn H. Immediate functional loading of TiOblast dental implants in full-arch edentulous maxillae: a 3-year prospective study. *Clin Oral Implants Res.* 2008;19:1254–1260.
- Ibanez JC, Tahhan MJ, Zamar JA, et al. Immediate occlusal loading of double acid-etched surface titanium implants in 41 consecutive full-arch cases in the mandible and maxilla: 6- to 74-month results. *J Periodontol.* 2005;76:1972–1981.
- van Steenberghe D, Molly L, Jacobs R, Vandekerckhove B, Quirynen M, Naert I. The immediate rehabilitation by means of a ready-made final fixed prosthesis in the edentulous mandible: a 1-year follow-up study on 50 consecutive patients. *Clin Oral Implant Res.* 2004;15:360–365.
- Norton MR. A short-term clinical evaluation of immediately restored maxillary TiOblast single-tooth implants. *Int J Oral Maxillofac Implants.* 2004;19:274–281.
- Spiekermann H, Jansen VK, Richter EJA. 10-year follow-up study of IMZ and TPS implants in the edentulous mandible using bar-retained overdentures. *Int J Oral Maxillofac Implants.* 1995;10:231–243.
- Rungcharassaeng K, Lozada JL, Kan YK, Kim JS, Campagni WV, Munoz CA. Peri-implant tissue response of immediately loaded, threaded, HA-coated implants: 1-year results. *J Prosthet Dent.* 2002;87:173–181.
- Lambert PM, Morris HF, Ochi S. Positive effect of surgical experience with implants on second-stage implant survival. *J Oral Maxillofac Surg.* 1997;55(suppl):12–18.
- Melo MD, Shafie H, Obeid G. Implant survival rates for oral and maxillofacial surgery residents: a retrospective clinical review with analysis of resident level of training on implant survival. *J Oral Maxillofac Surg.* 2006;64:1185–1189.
- Ribeiro FS, Pontes AE, Marcantonio E, Piattelli A, Neto RJ, Marcantonio E Jr. Success rate of immediate non-functional loaded single-tooth implants: immediate versus delayed implantation. *Implant Dent.* 2008;17:109–117.
- Esposito M, Grusovin MG, Polyzos IP, Felice P, Worthington HV. Interventions for replacing missing teeth: dental implants in fresh extraction sockets (immediate, immediate-delayed and delayed implants). *Cochrane Database Syst Rev.* 2010;8: CD005968.
- Chaushu G, Chaushu S, Tzohar A, Dayan D. Immediate loading of single-tooth implants: immediate versus non-immediate implantation. A clinical report. *Int J Oral Maxillofac Implants.* 2001;16:267–272.
- Chen ST, Wilson TG Jr, Hämmerle CH. Immediate or early placement of implants following tooth extraction: review of biological basis, clinical procedures, and outcomes. *Int J Oral Maxillofac Implants.* 2004;19(suppl):12–25.
- Rosenquist B, Grenthe B. Immediate placement of implants

into extraction sockets: implant survival. *Int J Oral Maxillofac Implants*. 1996;11:205–209.

25. Grunder U, Polizzi G, Goené R, et al. 3-year prospective multicenter follow-up report on the immediate and delayed-immediate placement of implants. *Int J Oral Maxillofac Implants*. 1999;14:210–216.

26. Polizzi G, Grunder U, Goené R, et al. Immediate and delayed implant placement into extraction sockets: a 5-year report. *Clin Implant Dent Relat Res*. 2000;2:93–99.

27. De Bruyn H, Collaert B. Early loading of machined-surface Brånemark implants in completely edentulous mandibles: healed bone versus fresh extraction sites. *Clin Implant Dent Relat Res*. 2002;4:136–142.

28. Aghaloo TL, Moy PK. Which hard tissue augmentation techniques are the most successful in furnishing bony support for implant placement? *Int J Oral Maxillofac Implants*. 2007;22(suppl):49–70.

29. Chiapasco M, Zaniboni M, Boisco M. Augmentation procedures for the rehabilitation of deficient edentulous ridges with oral implant. *Clin Oral Implants Res*. 2006;17(suppl):136–159.

30. Simion M, Fontana F, Rasperini G, Maiorana C. Long-term evaluation of osseointegrated implants placed in sites augmented with sinus floor elevation associated with vertical ridge augmentation: a retrospective study of 38 consecutive implants with 1- to 7-year follow-up. *Int J Periodont Restor Dent*. 2004;24:208–221.

31. Johansson B, Wannfors K, Ekenbäck J, Smedberg JI, Hirsch J. Implants and sinus-inlay bone grafts in a 1-stage procedure on severely atrophied maxillae: surgical aspects of a 3-year follow-up study. *Int J Oral Maxillofac Implants*. 1999;14:811–818.

32. Williamson RA. Rehabilitation of the resorbed maxilla and mandible using autogenous bone grafts and osseointegrated implants. *Int J Oral Maxillofac Implants*. 1996;11:476–488.

33. Isaksson S, Alberius P. Maxillary alveolar ridge augmentation with onlay bone-grafts and immediate endosseous implants. *J Craniomaxillofac Surg*. 1992;20:2–7.

34. Lekholm U, Wannfors K, Isaksson S, Adielsson B. Oral implants in combination with bone grafts. A 3-year retrospective multicenter study using the Brånemark implant system. *Int J Oral Maxillofac Surg*. 1999;28:181–187.

35. Chaushu G, Mardinger O, Calderon S, Moses O, Nissan J. The use of cancellous block allograft for sinus floor augmentation with simultaneous implant placement in the posterior atrophic maxilla. *J Periodontol*. 2009;80:422–428.

36. Valentini P, Abensur D. Maxillary sinus floor elevation for implant placement with demineralized freeze-dried bone and bovine bone (Bio-Oss): a clinical study of 20 patients. *Int J Periodont Restor Dent*. 1997;17:232–241.

37. Goodacre CJ, Bernal G, Rungcharassaeng K, Kan JY. Clinical complications with implant and implant prostheses. *J Prosthet Dent*. 2003;90:121–132.

38. Moy PK, Medina D, Shetty V, Aghaloo TL. Dental implant failure rates and associated risk factors. *Int J Oral Maxillofac Implants*. 2005;20:569–577.

39. Bain CA, Moy PK. The association between the failure of dental implants and cigarette smoking. *Int J Oral Maxillofac Implants*. 1993;8:609–615.

40. Kan JY, Rungcharassaeng K, Lozada JL, Goodacre CJ. Effects of smoking on implant success in grafted maxillary sinuses. *J Prosthet Dent*. 1999;82:307–311.

41. Balshi TJ, Wolfinger GJ. Dental implants in the diabetic patient: a retrospective study. *Implant Dent*. 1999;8:355–359.

42. Glauser R, Ree A, Lundgren A, Gottlow J, Hammerle CH, Scharer P. Immediate occlusal loading of Branemark implants applied in various jawbone regions: a prospective, 1-year clinical study. *Clin Implant Dent Relat Res*. 2001;3:204–213.

43. Lobbezoo F, Van Der Zaag J, Naeije M. Bruxism: its multiple

causes and its effects on dental implants—an updated review. *J Oral Rehabil*. 2006;33:293–300.

44. Jaffin RA, Berman CL. The excessive loss of Branemark fixtures in type IV bone: a 5-year analysis. *J Periodontol*. 1991;62:2–4.

45. Buser D, Mericske-Stern R, Bernard JP, et al. Long-term evaluation of non-submerged ITI implants. Part 1: 8-year life table analysis of a prospective multi-center study with 2359 implants. *Clin Oral Implants Res*. 1997;8:161–172.

46. Kacer CM, Dyer JD, Kraut RA. Immediate loading of dental implants in the anterior and posterior mandible: a retrospective study of 120 cases. *J Oral Maxillofac Surg*. 2010;68:2861–2867.

47. Bahat O. Treatment planning and placement of implants in the posterior maxillae: report of 732 consecutive Nobelpharma implants. *Int J Oral Maxillofac Implants*. 1993;8:151–161.

48. Lekholm U, Gunne J, Henry P, et al. Survival of the Brånemark implant in partially edentulous jaws: a 10-year prospective multicenter study. *Int J Oral Maxillofac Implants*. 1999;14:639–645.

49. van Steenberghe D, Lekholm U, Bolender C, et al. Applicability of osseointegrated oral implants in the rehabilitation of partial edentulism: a prospective multicenter study on 558 fixtures. *Int J Oral Maxillofac Implants*. 1990;5:272–281.

50. Winkler S, Morris HF, Ochi S. Implant survival to 36 months as related to length and diameter. *Ann Periodontol*. 2000;5:22–31.

51. ten Bruggenkate CM, Asikainen P, Foitzik C, Krekeler G, Sutter F. Short (6-mm) nonsubmerged dental implants: results of a multicenter clinical trial of 1 to 7 years. *Int J Oral Maxillofac Implants*. 1998;13:791–798.

52. Tawil G, Younan R. Clinical evaluation of short, machined-surface implants followed for 12 to 92 months. *Int J Oral Maxillofac Implants*. 2003;18:894–901.

53. Anitua E, Orive G, Aguirre JJ, Andía I. Five-year clinical evaluation of short dental implants placed in posterior areas: a retrospective study. *J Periodontol*. 2008;79:42–48.

54. Misch CE, Steingra J, Barboza E, Misch-Dietsh F, Cianciola LJ, Kazor C. Short dental implants in posterior partial edentulism: a multicenter retrospective 6-year case series study. *J Periodontol*. 2006;77:1340–1347.

55. Hagi D, Deporter DA, Pilliar RM, Arenovich T. A targeted review of study outcomes with short (< or = 7 mm) endosseous dental implants placed in partially edentulous patients. *J Periodontol*. 2004;75:798–804.

56. Himmlová L, Dostalova T, Kacovsky A, Konvickova S. Influence of implant length and diameter on stress distribution: a finite element analysis. *J Prosthet Dent*. 2004;91:20–25.

57. Hsu JT, Fuh LJ, Lin DJ, Shen YW, Huang HL. Bone strain and interfacial sliding analyses of platform switching and implant diameter on an immediately loaded implant: experimental and three-dimensional finite element analyses. *J Periodontol*. 2009;80:1125–1132.

58. Fontin-Tekamp FA, Slagter AP, Van Der Bilt A, et al. Biting and chewing in overdenture, full dentures, and natural dentitions. *J Dent Res*. 2000;79:1519–1524.

59. Misch CE, Suzuki JB, Misch-Dietsh FM, Bidez MW. A positive correlation between occlusal trauma and peri-implant bone loss: literature support. *Implant Dent*. 2005;14:108–116.

60. Allen PF, McMillan AS, Smith DG. Complications and maintenance requirements of implant-supported prostheses provided in a UK dental hospital. *Br Dent J*. 1997;182:298–302.

61. Naert I, Quirynen M, van Steenberghe D, Darius P. A study of 589 consecutive implants supporting complete fixed prostheses. Part II: prosthetic aspects. *J Prosthet Dent*. 1992;68:949–956.

62. Hemmings KW, Schmitt A, Zarb GA. Complications and maintenance requirements for fixed prostheses and overdentures in the edentulous mandible: a 5-year report. *Int J Oral Maxillofac Implants*. 1994;9:191–196.

