

Relative Abundance of *Ceratitis capitata* and *Anastrepha fraterculus* (Diptera: Tephritidae) in Diverse Host Species and Localities of Argentina

DIEGO F. SEGURA,¹ M. TERESA VERA,^{1, 2} CYNTHIA L. CAGNOTTI,¹ NORMA VACCARO,³ OLGA DE COLL,⁴ SERGIO M. OVRUSKI,⁵ AND JORGE L. CLADERA¹

Ann. Entomol. Soc. Am. 99(1): 70–83 (2006)

ABSTRACT Two fruit fly species (Diptera: Tephritidae) of economic importance occur in Argentina, the Mediterranean fruit fly, *Ceratitis capitata* (Wiedemann), and *Anastrepha fraterculus* (Wiedemann). Here, we compared the relative abundance of these fruit pests in 26 fruit species sampled from 62 localities of Argentina in regions where *C. capitata* and *A. fraterculus* coexist. In general, *C. capitata* was predominant over *A. fraterculus* (97.46% of the emerged adults were *C. capitata*), but not always. Using the number of emerged adults of each species, we calculated a relative abundance index (RAI) for each host in each locality. RAI is the abundance of *C. capitata* relative to the combined abundance of *A. fraterculus* and *C. capitata*. Some families of fruit species were more prone to show high (Rutaceae and Rosaceae) or low (Myrtaceae) RAI values, and also native plants showed lower RAI values than introduced plants. RAI showed high variation among host species in different localities, suggesting a differential use of these hosts by the two flies. There were localities where *A. fraterculus* was not found in spite of suitable temperature and the presence of hosts. Most host species showed little variation in RAI among localities, usually favoring *C. capitata*, but peach, grapefruit, and guava showed high variation. This suggests that these fruit species are suitable for both fruit flies but more favorable to one or the other, depending on local environmental conditions (e.g., relative humidity and degree of disturbance) of each locality.

KEY WORDS Tephritidae, relative abundance index, geographical distribution, Mediterranean fruit fly

THE FAMILY TEPHRITIDAE INCLUDES some of the most important fruit pests worldwide (White and Elson-Harris 1992). The economic damage caused by these flies is two-fold: direct damage to the fruit (larval activity) and limited access to potential markets because of quarantine restrictions imposed by countries that are free of these pests (Malavasi et al. 1994). In the American Continent, flies within *Anastrepha*, *Ceratitis*, *Rhagoletis*, and *Toxotrypana* cause the most economic damage (Landolt 1985, Enkerlin et al. 1989, Aluja 1994).

In Argentina, there are two quarantine species of fruit flies: the Mediterranean fruit fly, *Ceratitis capitata* (Wiedemann), and *Anastrepha fraterculus* (Wiedemann) (Aruani et al. 1996). *C. capitata* is native to Africa and has a wide distribution, covering many

tropical, subtropical, and temperate regions of the world (Copeland et al. 2002). This species shows a high adaptability to diverse climates as well as a large number of host fruit species (>350; Liquido et al. 1991). Its presence in Argentina was first recorded at the beginning of the 20th century in orchards located in the vicinity of Buenos Aires city (Vergani 1952). Later, it was reported in commercial orchards of northeastern and northwestern regions of the country. The last region in which it was reported was northern Patagonia (southern Argentina), where *C. capitata* was first detected in 1952 (Rial 1997). *A. fraterculus* is native to South America and is distributed from Mexico to Argentina, but there is morphological and genetic evidence indicating that there are many cryptic species (Steck 1991, Hernández-Ortiz et al. 2004) and that not all the species within this species complex are pests (Aluja et al. 2003). In Argentina, *A. fraterculus* is mainly distributed in regions with tropical and subtropical climate (Ovruski et al. 2003). It is also a polyphagous species that attacks different families of fruit species, but the number of hosts cited is smaller than that for *C. capitata* (≈80 species; Norrbom 2004). Both species cause significant annual losses to the fruit

¹ Instituto de Genética "E.A. Favret", INTA Castelar, Argentina.

² Estación Experimental Agroindustrial Obispo Colombes, Tucumán, Argentina.

³ Estación Experimental Agropecuaria Concordia (INTA), Entre Ríos, Argentina.

⁴ Estación Experimental Agropecuaria Montecarlo (INTA), Misiones, Argentina.

⁵ PROIMI Biotecnología-CONICET, División Control Biológico de Plagas, Tucumán, Argentina.

production of Argentina and constitute a major barrier to the expansion of this market (Ovruski et al. 1999).

In spite of the importance of these fruit fly species, there is little information published on the relative abundance of *C. capitata* and *A. fraterculus* in areas of Argentina where they coexist. Other than Vergani (1956), the only map available showing the distribution of the two species at the national level is hypothetical (Ortiz 1999). Variability in the relative abundance of the two species among different regions was reported by Vattuone et al. (1995) based on average values among different hosts, but the relationship between the two species strongly depended on the fruit considered. Comparisons based on trapping data (FAO 1989, Segade and Polack 1999, Vattuone et al. 1999) are biased by the degree of attraction that adult flies of the two species show to the bait used in the traps. Some local studies provide data based on fruit sampling (Costilla 1967, FAO 1989, Putruele 1993, Vattuone et al. 1999). In Tucumán province, Costilla (1967) found a greater proportion of *C. capitata* than *A. fraterculus* in citrus orchards of grapefruit, *Citrus paradisi* L., and orange, *Citrus sinensis* (L.) Osbeck. Schliserman and Ovruski (2004), working in areas of the same province with much native vegetation, found mainly *C. capitata* in bitter orange, *Citrus aurantium* L. Ovruski et al. (2003) also reported proportions of *C. capitata* and *A. fraterculus* adults and larval infestation levels in wild or commercially grown plants, both native and introduced, and emphasized the importance of *Citrus* spp. as hosts of *C. capitata* and native fruit species as hosts of *A. fraterculus*. In Entre Ríos province, Putruele (1993) recovered both species from grapefruit; peach, *Prunus persica* L.; fig, *Ficus carica* L.; and mandarin, *Citrus reticulata* Blanco; but only in grapefruit was *A. fraterculus* more abundant than *C. capitata*. In the same province, apple, *Malus domestica* Borkh., and pear, *Pyrus communis* L., were infested only by *C. capitata*, whereas pomegranate, *Punica granatum* L., and quince, *Cydonia oblonga* Mill., were attacked only by *A. fraterculus* (Putruele 1993). In Catamarca province, Vattuone et al. (1999) reported that orange, peach, grapefruit, mandarin, and kumquat, *Citrus aurantium* variety *myrtifolia* Ker-Gawl., were infested only by *C. capitata*. In La Rioja province, Nasca et al. (1996) recorded greater infestation of *C. capitata* than *A. fraterculus* in fig; persimmon, *Diospyros kaki* L.; quince; pomegranate; plum, *Prunus domestica* L.; and apricot, *Prunus armeniaca* L. Although these studies are valuable, there has been no comprehensive study comparing the relative abundances of both species along the variety of hosts and regions present in Argentina.

The aim of this study was to provide these comparative data for *C. capitata* and *A. fraterculus*. We describe the relative abundance of these flies through the analysis of data obtained from fruit samplings of many different host species. Specifically, we were interested in comparing the possible variability in the abundances of the two fly species 1) among different host species from the same locality and 2) among different localities for the same host. We also exam-

ined the influence of plant taxonomy and origin (native or introduced) as well as variations in the climate on the relative abundance of these two species.

Materials and Methods

Sampling. Fruit sampling was carried out in those political provinces of Argentina where the two fruit flies are reported to coexist (see Appendix 1). The localities ranged from undisturbed areas with most of the vegetation being native to very disturbed systems, such as suburban areas or agricultural landscapes. The climatic characteristics, specifically rainfall and relative humidity, also varied widely.

Sampling was performed mainly during the fruiting season of 1999–2000. Nevertheless, because of inherent logistical complexities faced in a study covering such a large geographical area, sampling data derived from the 2000–2001 season in some localities (e.g., Concordia), and from the 1998–1999 fruiting season in others (e.g., Posadas). To include as much area as possible, we also present fruit-collecting data from 1993 for three localities in the province of Catamarca. For the localities that were sampled during more than one fruiting season, no substantial differences between years were found in the relative abundance index (RAI) values (not shown), so these data were pooled. Descriptions of each locality are provided in Appendix 1.

All fruit species sampled have been previously cited as hosts for both fruit fly species (Liquido et al. 1991, Norrbom 2004), and they included native and introduced plants. In each locality, at least 10 fruit were sampled for each host species. Only fruit with evident signs of infestation by fruit flies was collected. The sampled fruit was placed in plastic trays over a layer of sand (or vermiculite), which was used by larvae as a pupation substrate after leaving the fruit. Pupae were separated from the sand using a sieve and then transferred to new containers, where they were maintained until emergence. Adults were identified, and the number of individuals of each fruit fly species was recorded. For inclusion in the analysis, at least 10 adults were required for a given host species and locality.

Data Analysis. To describe the relationship between the abundances of the two fruit fly species, a relative abundance index (RAI) was calculated for each host species in each locality according to the following formula: $RAI_{xy} = Cc / (Cc + Af)$, where *Cc* and *Af* are the number of emerged adults of *C. capitata* and *A. fraterculus*, respectively, for host *X* in locality *Y*. RAI ranges from 0 (exclusive presence of *A. fraterculus*) to 1 (exclusive presence of *C. capitata*). The RAI value for a given host (RAI_x) was estimated as the mean value for all of the localities, where that particular host was sampled. According to the RAI value obtained, hosts and localities were assigned to one of five categories: exclusive presence of one or the other species ($RAI = 0$ or $RAI = 1$), both species present but higher abundance of one or the other ($0 < RAI < 0.33$ or $0.66 < RAI < 1$), and intermediate cases ($0.33 \leq RAI \leq 0.66$).

Table 1. Host species sampled

Common name	Scientific name	Family	Native/exotic
Albarillo	<i>Ximenia americana</i> L.	Olacaceae	N
Apple	<i>Malus domestica</i> Borkh	Rosaceae	E
Asian pear	<i>Pyrus pyrifolia</i> (Burn. F.) Nakai	Rosaceae	E
Bitter orange	<i>Citrus aurantium</i> L.	Rutaceae	E
Feijoa	<i>Feijoa sellowiana</i> Berg	Myrtaceae	N
Fig	<i>Ficus carica</i> L.	Moraceae	E
Grapefruit	<i>Citrus paradisi</i> L.	Rutaceae	E
Guava	<i>Psidium guajaba</i> L.	Myrtaceae	N
Kiwi fruit	<i>Actinidia chinensis</i> Planch.	Actinidiaceae	E
Kumquat	<i>Citrus aurantium</i> var. <i>myrtifolia</i> Ker-Gawl	Rutaceae	E
Loquat	<i>Eriobotrya japonica</i> (Thunb.) Lindl	Rosaceae	E
Mandarine	<i>Citrus reticulata</i> Blanco	Rutaceae	E
Orange	<i>Citrus sinensis</i> (L.) Osbeck	Rutaceae	E
Papaya	<i>Carica papaya</i> L.	Caricaceae	N
Peach	<i>Prunus persica</i> L.	Rosaceae	E
Pear	<i>Pyrus communis</i> L.	Rosaceae	E
Persimmon	<i>Diospiros kaki</i> L.f.	Ebenaceae	E
Plum	<i>Prunus insititia</i> L	Rosaceae	E
Quince	<i>Cydonia oblonga</i> Mill	Rosaceae	E
Ubajay	<i>Hexachlamys edulis</i> (O. Berg)	Myrtaceae	N

For every host–locality combination, we compared the observed frequencies with expected frequencies estimated multiplying the number of cases by the probability that in a random sample of 10 pupae: all of them were *C. capitata*; all of them were *A. fraterculus*; seven to nine pupae were of one of the two flies (and three to one of the other); and four to six pupae were of either of them. A chi-square goodness-of-fit test was performed to compare observed and expected frequencies (StatSoft, Inc. 2000).

A Mann–Whitney test was performed to compare the RAI values recorded for native and introduced fruit species (StatSoft, Inc. 2000). To compare the RAI values found for the main taxonomic families of fruit species (Myrtaceae, Rosaceae, and Rutaceae), a non-parametric analysis of variance (ANOVA) (Kruskal–Wallis test) was performed. When this analysis showed significant differences, nonparametric multiple comparison tests were performed (Analytical Software 2000).

Results

We obtained fruit samples from 62 localities in 12 of the 24 political provinces of Argentina, covering six ecological regions (see *Appendix 1*) and representing the complete area shared by both species of fruit flies in the country. In total, 30,354 fruit were collected, belonging to 20 host species from eight families of plants. Three of these families are native to South America (Myrtaceae, Caricaceae, and Olaceae), and five are introduced (Actinidiaceae, Ebenaceae, Moraceae, Rosaceae, and Rutaceae) (Table 1). Of 43,142 pupae recovered from the fruit, a total of 27,301 adults emerged: 23,608 (97.46%) were *C. capitata*, and 3,693 (2.54%) were *A. fraterculus*. We recovered parasitoids in only seven sampling sites, and in those cases the percentage of parasitism was never higher than 5%.

Thirty-two localities and 12 host species were sampled in northwestern Argentina (NWA) (Fig. 1;

Appendix 1), and 30 localities and 18 host species were sampled in northeastern Argentina (NEA) (Fig. 2; *Appendix 1*). A general predominance of *C. capitata* was observed, which was higher in NWA than in NEA ($U = 1096.5$, $P = 0.042$; Mann–Whitney test). The proportion of samples in which $RAI = 1$ was 72% in the NWA and 50% in NEA (Table 2).

In some cases, RAI showed a marked variation among hosts in the same locality, e.g., in Yuto from NWA and in Posadas and Concordia from NEA (Figs. 1 and 2; Table 2). In those cases, some hosts had a RAI that was very high (mandarin from Yuto; persimmon from Posadas; and peach, guava, and orange from Concordia) or very low (guava, *Psidium guajava* L., from Yuto; grapefruit and guava from Posadas; and feijoa from Concordia), whereas other hosts showed intermediate values. In other localities, the variation among hosts was considerably lower. For example, in Chilecito (Fig. 1) and in Saenz Peña and San Pedro (Fig. 2), *C. capitata* was the main fruit fly species found, whereas in Montecarlo (Fig. 2) *A. fraterculus* was more abundant than *C. capitata* in almost all host species.

Irrespective of locality, many fruit species showed values of RAI closer to 1 than to 0, e.g., orange and fig. No host species consistently had values of RAI near 0 (Table 3). Other fruit species, such as grapefruit, peach, and guava showed a wide range of RAI values (Table 3).

Native host plants showed significantly lower values of RAI than introduced hosts (Table 4). Significant differences were also found among the three most abundant families of host species (Table 4). Nonparametric multiple comparisons showed differences between Myrtaceae and Rutaceae ($df = 1$, $P < 0.05$), whereas host species belonging to the family Rosaceae gave intermediate values of RAI that did not differ statistically from the other two.

The number of cases in which the two species of fruit flies shared a host in equal abundance (Fig. 3) was




Fig. 1. RAI for each locality and host species sampled in NWA.

lower than expected, whereas the cases in which only one fruit fly species was recovered from a host was much higher than expected ($\chi^2 = 326.39$, $P < 0.001$).

Discussion

In some localities, we sampled fruit suitable for both *C. capitata* and *A. fraterculus*, and only one fruit fly species was recovered, or one of them occurred in extremely low abundance (Sáenz Peña, Monte Case-

ros, Montecarlo, San Pedro, and Bella Vista in Fig. 2; Chilecito and Campo Santo in Fig. 1). This fact suggests that factors other than availability of suitable hosts are limiting the establishment of one fruit fly species and not the other. Among these factors, we can postulate abiotic factors, such as temperature, humidity, or rainfall, and biotic factors, such as the duration of periods without mature suitable fruit, the competition between the two fruit fly species (Celedonio-Hurtado et al. 1995, and references




Fig. 2. RAI for each locality and host species sampled in NEA.

therein), and the degree of environmental disturbance. Parasitism could bias the relative abundances of these fruit fly species favoring one of the two

species (Ovruski et al. 2004); however, the numbers recovered were so low that parasitism did not significantly affect the RAI values obtained here.

Table 2. RAI values per localities and hosts

Province	Locality	Host	Sampled fruit	Infestation level ^a	RAI
Buenos Aires	Mercedes	Plum	500	1.31	0.33
		Peach	41	2.66	1.00
	Pilar	Kumquat	181	1.15	0.36
	Gob. Castro	Orange	265	0.72	1.00
	Doyle	Grapefruit	676	0.14	1.00
	San Pedro	Orange	211	1.76	1.00
		Peach	1425	3.64	0.99
		Mandarin	1685	0.57	0.99
		Plum	791	0.51	1.00
		Apple	120	0.60	1.00
		Asian pear	150	2.20	1.00
		Fig	1457	2.90	1.00
		Pear	30	1.67	1.00
		Persimmon	495	2.13	0.99
		Kiwi fruit	230	0.73	1.00
		Feijoa	143	1.92	0.97
		Peach	15	3.00	0.79
	Villa Adelina	Feijoa	50	2.92	0.03
Catamarca	Ituzaingó	Peach	103	1.77	0.83
	S. F. V. Catamarca	Quince	151	1.09	1.00
	Capayan	Peach	91	1.16	1.00
	Chumbicha	Orange	36	1.36	1.00
	San José	Fig	88	0.68	1.00
		Peach	41	1.56	0.88
		Mandarin	27	1.96	1.00
		Orange	30	1.27	1.00
	Sta. Rosa	Grapefruit	30	1.37	1.00
		Kumquat	41	2.15	1.00
		Fig	50	6.04	1.00
		Orange	35	0.69	1.00
	Sumalao	Peach	57	1.79	0.93
	San Isidro	Peach	30	4.13	0.90
Corrientes	Villa Dolores	Grapefruit	38	1.24	1.00
	9 de Julio	Grapefruit	31	3.35	0.86
	Bella Vista	Kumquat	123	0.66	1.00
	Ituzaingó	Guava	286	1.11	1.00
		Ubajay	70	5.03	0.01
		Orange	129	1.25	1.00
		Mandarin	108	0.99	1.00
	Monte Caseros	Guava	101	2.18	0.98
		Grapefruit	60	1.18	0.97
		Grapefruit	59	1.93	0.92
Chaco	Virasoro	Papaya	26	1.54	1.00
	Saenz Peña	Guava	87	1.64	1.00
		Grapefruit	12	1.25	1.00
		Guava	375	4.77	0.06
Entre Ríos	Palmar	Peach	567	2.76	1.00
		Feijoa	1468	0.60	0.50
		Guava	987	1.18	0.99
		Mandarin	743	0.58	0.95
	Concordia	Orange	568	0.17	1.00
		Mandarin	252	0.43	1.00
		Mandarin	153	1.40	1.00
		Mandarin	80	1.46	1.00
	Chajarí	Mandarin	21	6.57	1.00
	Humaitá	Guava	33	6.00	0.99
	La Criolla	Guava	173	5.85	1.00
	Pto Yerúa	Orange	219	0.49	1.00
	Villa Adela	Guava	461	3.36	0.28
	Villa Zorraquín	Mandarin	53	0.68	1.00
		Bitter orange	36	3.47	1.00
Jujuy	Ledesma	Orange	76	1.04	1.00
	Perico	Orange	195	0.75	1.00
La Rioja	Chilecito	Albarillo	31	0.81	1.00
		Plum	23	3.17	0.96
		Peach	1296	0.89	1.00
		Fig	183	4.91	1.00
	Los Dorados	Orange	36	1.69	1.00
		Fig	25	3.72	1.00
		Peach	170	0.92	1.00
		Peach	25	2.68	1.00
	Guandacol				
	Los Sarmientos				

(continued)

Table 2. Continued

Province	Locality	Host	Sampled fruit	Infestation level ^a	RAI
Misiones	Caraguatay	Guava	79	1.85	0.00
		Peach	28	2.36	0.00
		Guava	106	3.01	0.00
		Mandarin	591	0.36	0.35
		Grapefruit	45	0.49	0.00
		Peach	64	0.95	0.07
		Feijoa	377	3.00	0.00
		Loquat	761	0.19	0.00
		Persimmon	80	2.46	0.99
	Posadas	Guava	3272	0.99	0.46
		Mandarin	153	0.78	0.85
		Grapefruit	169	0.83	0.00
		Persimmon	26	2.88	1.00
		Peach	31	5.52	1.00
		Papaya	14	1.93	1.00
Salta	Taruma	Grapefruit	22	1.18	1.00
		Grapefruit	131	1.36	1.00
	Campo Santo	Bitter orange	18	0.72	1.00
		Guava	225	0.40	0.00
	Cnia. Santa Rosa	Bitter orange	55	2.24	1.00
		Grapefruit	31	1.77	1.00
		Mandarin	11	3.73	1.00
		Peach	23	3.43	0.00
San Luis	Quines	Orange	16	2.25	1.00
	Arocena	Bitter orange	32	1.38	1.00
		Peach	405	1.46	1.00
Santa Fé	Reconquista	Grapefruit	103	1.50	1.00
	Monje	Bitter orange	57	1.79	1.00
	Famailla	Guava	3371	0.53	0.06
		Guava	106	2.54	0.00
Tucumán	La Rinconada	Bitter orange	12	3.25	1.00
	Lules	Peach	186	4.06	0.01
	San Javier	Peach	160	3.86	0.98
	Tafi Viejo	Peach	325	5.38	0.95
	Taficillo	Orange	615	0.52	1.00
	Timbó				

^a Infestation level is the number of pupae per fruit.

Consideration of specific localities offers some insight into the factors affecting fly distribution. For example, in Saenz Peña, *A. fraterculus* was absent from all the host species sampled, including guavas, one of the primary host for this species (Putruele 1993, Aluja et al. 2000, Selivon 2000). The thermal regime of Saenz Peña is suitable for this species, but the annual relative humidity is close to 50%, suggesting that this site is too

Table 3. Number of localities in each RAI categories for each host species

Host species	RAI = 0	0 < RAI < 0.33	0.33 < RAI < 0.66	0.66 < RAI < 1	RAI = 1	n	RAI range
Albarillo	0	0	0	0	1	1	1.00–1.00
Apple	0	0	0	0	1	1	1.00–1.00
Asian Pear	0	0	0	0	1	1	1.00–1.00
Bitter orange	0	0	0	0	6	6	1.00–1.00
Feijoa	1	1	1	1	0	4	0.00–0.97
Fig	0	0	0	0	5	5	1.00–1.00
Grapefruit	2	0	0	3	8	13	0.00–1.00
Guava	4	3	1	3	3	14	0.00–1.00
Kiwi fruit	0	0	0	0	1	1	1.00–1.00
Kumquat	0	0	1	0	2	3	0.36–1.00
Loquat	1	0	0	0	0	1	0.00–0.00
Mandarin	0	0	1	3	8	12	0.35–1.00
Orange	0	0	0	0	13	13	1.00–1.00
Papaya	0	0	0	0	2	2	1.00–1.00
Peach	3	1	0	7	9	20	0.00–1.00
Pear	0	0	0	0	1	1	1.00–1.00
Persimmon	0	0	0	2	1	3	0.99–1.00
Plum	0	1	0	1	1	3	0.33–1.00
Quince	0	0	0	0	1	1	1.00–1.00
Ubajay	0	1	0	0	0	1	0.01–0.01

RAI = 0 indicates only AF found; 0.01 < RAI < 0.33 indicates more AF than CC; 0.33 < RAI < 0.66 indicates same amt of AF as CC; 0.66 < RAI < 0.99 indicates more CC than AF; RAI = 1 indicates only CC present.

Table 4. RAI values by origin and botanic family of host plants

Classification by	RAI ^a	Q25 ^b	Q75 ^c	n	Nonparametric test
Origin					
Introduced	1.000	0.974	1.000	84	Mann–Whitney test: $U = 490.00, P < 0.001$
Native	0.370	0.004	1.000	22	
Host family					
Myrtaceae	0.060a	0.000	0.982	19	Kruskal–Wallis test: $H = 24.11, df = 2, P < 0.001, n = 94$
Rosaceae	0.986ab	0.810	1.00	28	
Rutaceae	1.000b	1.000	1.000	47	

^a Medians followed by a different letter differed statistically ($P < 0.05$; multiple comparison Dunn’s test).

^b First quartile.

^c Third quartile.

dry for *A. fraterculus*. Orán, by contrast, has a similar thermal regime but a higher annual relative humidity, and here guavas were heavily infested by *A. fraterculus*. Environmental disturbance supposedly favors *C. capitata* (Putruele 1997, Malavasi et al. 2000, Ovruski et al. 2003). This seems to be the case for Saenz Peña with a very extensive agricultural landscape and very little native vegetation, but not for Orán [although currently there is a heavy trend to deforest the Yungas and start soybean, *Glycine max* (L.) Merr., plantations]. However, in Montecarlo, where the original environment also has been disturbed, there is high abundance of *A. fraterculus*, indicating that the local environment may have a stronger impact than the disturbance on the relative abundance of this fruit fly. Montecarlo, in Misiones province, with subtropical climate and high relative humidity exhibits a landscape with dense vegetation and backyards with fruit fly host plants, many of them native, that provide excellent refuges for *A. fraterculus*. In San Pedro, the thermal regime and the relative humidity seem appropriate for the development of *A. fraterculus*; however, RAI values for this locality are high because this species is present in a small number of hosts and in low abundance. The absence of suitable hosts for *A. fraterculus* from late autumn to middle spring (Segura et al. 2004) is the most likely explanation for this pattern.

The great variation in RAI among hosts (see Posadas, Concordia, and others) indicates a pattern of differential use of the available hosts. The possible

explanations are 1) a different pattern of host preference in adults of each fruit fly species; 2) asymmetric interspecific competition (the result of which depends on the fruit species; Fitt 1989); 3) differential mortality of eggs, larvae, or both in each host species (Carey 1984); or 4) differential ability of each fruit fly species to find and infest different fruit species.

Citrus spp. were found to be better hosts for *C. capitata* than for *A. fraterculus*, regardless of the inter-locality variation in biotic and abiotic factors that may favor one of the other species, at least in Argentina (Tables 2 and 3). Mandarin, orange, and bitter orange showed lower variation in RAI than the other fruit, and the most abundant species was always *C. capitata*. In agreement with other observations (Malavasi and Morgante 1980, Putruele 1993, Nasca et al. 1996, Vaccaro 2000, Ovruski et al. 2003) we found that grapefruit was the only *Citrus* with values of RAI favoring *A. fraterculus*. Several studies (Nascimento et al. 1984, da Silva-Branco et al. 2000, Aluja et al. 2003) have shown the low suitability of *Citrus* spp. as host for *Anastrepha* spp., but in the laboratory, forced development on grapefruit, orange, and lemon, *Citrus limon* L., showed better recovery of *A. fraterculus* pupae in the case of grapefruit (Gramajo 2004). Ovruski et al. (2003) proposed a stronger attraction of *C. capitata* toward infochemicals (sensu Dicke and Sabelis 1988) produced by *Citrus* as another explanation for the high RAI values found in these host species (for example, Howse and Knapp (1996) suggested that some components of male *C. capitata* pheromone are similar to




Fig. 3. Number of expected and observed cases (for each combination of host species and locality) for each classification category of RAI (see text for delimitation of the RAI categories). The observed frequencies differed statistically from the expected frequencies (chi-square test: $\chi^2 = 326.39, P < 0.001$).

volatiles emitted by *Citrus* trees and fruit). Asymmetrical larval competition favoring *C. capitata* also could be postulated (but then, it is not clear why this asymmetry would be reversed for grapefruit).

From apple and pear, we recovered pupae of *C. capitata* in San Pedro (Buenos Aires province), as did previous studies in the neighboring province of Entre Ríos (FAO 1989, Putruele 1996). Nasca et al. (1996) recorded *A. fraterculus* pupae from pear collected in Antinaco-Los Colorados Valley (La Rioja province). Several studies carried out in Brazil report the presence of *A. fraterculus* in these two fruit species (Malavasi et al. 1980, Kovaleski et al. 2000, Nora et al. 2000). However, Ovruski et al. (2003) did not find infestation by any fruit fly in apple and pear (157 and 196 fruit sampled, respectively) in NWA, questioning the status of these fruit species as hosts for the two fly species. Surveys of these two fruit species should be expanded, with emphasis in NWA.

C. capitata was more abundant in plant species belonging to the family Rutaceae, whereas *A. fraterculus* was predominant in plants of the family Myrtaceae. Species belonging to the family Rosaceae showed intermediate values of RAI, roughly corresponding to the comparison of RAI between introduced and native species: *C. capitata* dominates in introduced plants (Rutaceae and Rosaceae, among others), whereas in native plants (Myrtaceae among others) *A. fraterculus* shows higher abundance (Table 4). Various Brazilian authors also found this (Malavasi and Morgante 1980, de Souza Filho et al. 2000, Malavasi et al. 2000, Veloso et al. 2000). Eskafi and Kolbe (1990) described the same pattern in Guatemala, although their samplings also included different *Anastrepha* species. Ovruski et al. (2003) also reported that the introduced fruit species favor *C. capitata* and that the native species serve as a reservoir for *A. fraterculus* (with two exceptions discussed below). The fact that almost 85% of all fruit species sampled are exotic in Argentina could be responsible, at least to some extent, for the high predominance of *C. capitata* in our samplings.

As noted above, *A. fraterculus* showed better yields in native plants, probably because of their common evolutionary history. *C. capitata*, being such a polyphagous species with such high reproductive capacity (Liquido et al. 1991), may gain advantage in introduced hosts with which *A. fraterculus* has had less contact in its evolutionary history. Moreover, plant species introduced to the Americas from the same region as *C. capitata*, for example, coffee, *Coffea arabica* L., constitute good hosts for this fly (Harris and Lee 1989, Malavasi et al. 2000). Interestingly, Copeland et al. (2002) did not find *C. capitata* in guavas sampled in Kenya (in a sample of 84 fruit), where this fly is native, and *P. guajava* is an introduced species. In our study, *C. capitata* was found infesting guavas in 10 of 14 localities sampled (Table 2; Figs. 1 and 2). Another exception, mentioned by Ovruski et al. (2003), might be peach and plum (both introduced species). But, if we calculate the RAIs from their published data (0.16 and 0.25, respectively) they differ markedly from those found in the current study (average RAI of 0.77

and 0.76 for peach and plum, respectively). In Ovruski et al. (2003), however, the sampling of these two host species occurred in forest areas, scattered among native vegetation, whereas in our study, they occurred in highly disturbed areas, illustrating the strong influence of the environment on the RAI values. RAI values lower than 0.10 for peach, in Itacuruzú, Montecarlo, and San Javier, located in areas with native vegetation and subtropical climate are good examples supporting our explanation.

We found that both species tended to occur alone many more times than they occurred together sharing one host in one locality. We could interpret this pattern as competitive exclusion of one species by the other. It has been suggested that, when two or more fruit fly species coexist, some form of competition for hosts could arise (Duyck et al. 2004). Most examples of interspecific competition among tephritids derive from situations in which a new species has been introduced into a given environment (Duyck et al. 2004). For example, interspecific competition was proposed to explain the displacement of *C. capitata* by *Bactrocera dorsalis* (Hendel) (Fitt 1989) in Hawaii and by *Bactrocera tryoni* (Froggatt) in Australia (Allman 1939, Andrewartha and Birch 1954, Christenson and Foote 1960, Bateman 1971, Fitt 1989). Owing to the short period of coexistence (≈ 100 yr), mechanisms that tend to minimize the competition for resources ("avoidance" sensu Díaz-Fleischer et al. 2000, Duyck et al. 2004, Sivinski et al. 2004) have probably not yet evolved. However, the patterns of relative abundance give only indirect evidence for the existence of interspecific competition. Fitt (1989) suggests looking for direct evidence of competition, as a modification in the abundance of one species after manipulating the abundance of the other.

In conclusion, this first attempt to analyze the relative abundance of *C. capitata* and *A. fraterculus* covering different regions and different hosts in Argentina proved that both species coexist here in several areas and exhibit similar ecological requirements. Therefore, we should expect strong competition between them in habitats where the resources are scarce, as in wild or urban habitats where the density of host plants is usually low. These habitats serve as refuges for small populations that are usually neglected by traditional pest control efforts and may be *foci* where reinfestation starts. Future studies of interspecific competition between *C. capitata* and *A. fraterculus* should focus on these habitats to produce valuable information for area-wide management of these pests. *C. capitata* is the major fruit fly pest in almost all regions in Argentina. The sterile insect technique (Knippling 1955), successfully implemented in La Rioja, Mendoza, and San Juan provinces, and the Patagonia region (De Longo et al. 2000, Frissolo et al. 2001, Sánchez et al. 2001), aims at the eradication of *C. capitata*. It would be very useful to be able to predict the response of *A. fraterculus* populations to a marked decrease in the density of *C. capitata* and identify areas likely to experience an increase in the *A. fraterculus* population, thereby avoiding outbreaks of this pest.

Acknowledgments

We thank to M. de la Vega, D. Figueroa, L. Hansen, F. Milla, J. Mousques, N. Petit-Marty, A. Polack, L. Santinoni, G. Segade, and E. Willink, for collaboration during the fruit collection. We also thank E. Vattuone for help during the bibliographic search and D. O. McInnis, M. Viscarret, and S. López for comments on a previous version of this manuscript. We are grateful for revisions by T. Shelly and two anonymous reviewers. This work was supported by Foncyt PID 615 to J.L.C. and PIP-CONICET n° 04973 to S.M.O.

References Cited

- Allman, S. L. 1939. The Queensland fruit fly – observations on breeding and development. *Agric. Gaz. N.S.W.* 50: 499–501, 547–549.
- Analytical Software. 2000. Statistix for Windows (computer program manual). Analytical Software, Tallahassee, FL.
- Andrewartha, H. G., and L. C. Birch. 1954. Distribution and abundance of animals. University of Chicago Press, Chicago, IL.
- Aluja, M. 1994. Bionomics and management of *Anastrepha*. *Annu. Rev. Entomol.* 39: 155–178.
- Aluja, M., J. Piñero, M. López, C. Ruiz, A. Zúñiga, E. Piedra, F. Díaz-Fleischer, and J. Sivinski. 2000. New host plant and distribution records in Mexico for *Anastrepha* spp., *Toxotrypana curvicauda* Gerstaecker, *Rhagoletis zoqui* Bush, *Rhagoletis* sp., and *Hexachaeta* sp. (Diptera: Tephritidae). *Proc. Entomol. Soc. Wash.* 102: 802–815.
- Aluja, M., D. Pérez-Staples, R. Macías-Ordoñez, J. Piñero, B. A. McPheron, and V. Hernández-Ortiz. 2003. Non-host status of *Citrus sinensis* cv. Valencia and *C. paradisi* cv. Ruby Red to Mexican *Anastrepha fraterculus* (Diptera: Tephritidae). *J. Econ. Entomol.* 96: 1693–1703.
- Anonymous. 1992. Atlas de la República Argentina. Editorial El Ateneo, Buenos Aires, Argentina.
- Aruani R., A. Ceresa, J. C. Granados, G. Taret, P. Peruzzoti, and G. Ortiz. 1996. Advances in the national fruit fly control and eradication program in Argentina, pp. 521–530. In B. McPheron and G. J. Steck [eds.], *Fruit flies pest: a world assessment of their biology and management*. St. Lucie Press, Delray Beach, FL.
- Bateman, M. A. 1971. The ecology of fruit flies. *Annu. Rev. Entomol.* 17: 493–518.
- Cabrera, A. L. 1976. Regiones Fitogeográficas Argentinas. Enciclopedia Argentina de Agricultura y Jardinería. Enciclopedia Argentina de la Agricultura y Jardinería. 2nd ed. vol. 2. Section 1. Editorial Acme, Buenos Aires, Argentina.
- Cabrera, A. L., and A. Willink. 1973. Biogeografía de América Latina. Monografía 13. Serie Biología. OEA. Secretaría General de la OEA, Washington, DC.
- Carey, J. R. 1984. Host-specific demographic studies of the Mediterranean fruit fly *Ceratitis capitata*. *Ecol. Entomol.* 9: 261–270.
- Celedonio-Hurtado, H., M. Aluja, and P. Liedo. 1995. Adult population fluctuations of *Anastrepha* species (Diptera: Tephritidae) in tropical orchard habitats of Chiapas, México. *Environ. Entomol.* 24: 861–869.
- Christenson, L. D., and R. H. Foote. 1960. Biology of fruit flies. *Annu. Rev. Entomol.* 5: 171–192.
- Copeland, R. S., R. A. Wharton, Q. Luke, and M. de Meyer. 2002. Indigenous hosts of *Ceratitis capitata* (Diptera: Tephritidae) in Kenya. *Ann. Entomol. Soc. Am.* 95: 672–694.
- Costilla, M. A. 1967. Importancia de la mosca del mediterráneo (*Ceratitis capitata* Wied.) en los citrus de Tucumán y su control. *Boletín Est. Exp. Agr. Tucumán* 105: 1–12.
- da Silva-Branco, E., J. D. Vendramin, and F. Denardi. 2000. Resistencia às moscas-das-frutas em fruteiras, pp. 161–167. In A. Malavasi and R. A. Zucchi [eds.], *Moscas-das-frutas de importância econômica no Brasil: conhecimento básico e aplicado*. Holos Editora, Ribeirão Preto, Brazil.
- De Longo, O., A. Colombo, P. Gómez-Riera, and A. Bartolucci. 2000. The use of massive SIT for the control of the medfly, *Ceratitis capitata* (Wied.), strain SEIB 6-96, in Mendoza, Argentina. 351–359. In K. H. Tan [ed.], *Area-wide management of fruit flies and other insect pests*. Universiti Sains Malaysia Press, Penang, Malaysia.
- de Souza Filho, M. F., A. Raga, and R. A. Zucchi. 2000. São Paulo, pp. 277–283. In A. Malavasi and R. A. Zucchi [eds.], *Moscas-das-frutas de importância econômica no Brasil: conhecimento básico e aplicado*. Holos Editora, Ribeirão Preto, Brazil.
- Díaz-Fleischer, F., D. Papaj, R. Prokopy, A. Norrbom, and M. Aluja. 2000. Evolution of fruit fly oviposition behavior, pp. 811–841. In M. Aluja and A. Norrbom [eds.], *Fruit flies (Tephritidae): phylogeny and evolution of behavior*. CRC, Boca Raton, FL.
- Duyck, P., P. David, and S. Quilici. 2004. A review of relationships between interspecific competition and invasions in fruit flies (Diptera: Tephritidae). *Ecol. Entomol.* 29: 511–520.
- Dicke, M., and M. W. Sabelis. 1988. Infochemical terminology: should it be based on cost-benefit analysis rather than origin of compounds? *Funct. Ecol.* 2: 131–139.
- Enkerlin, D., L. García, and F. López M. 1989. Pest status: Mexico, Central and South America, pp. 83–90. In A. S. Robinson and G. Hooper [eds.], *Fruit flies: their biology, natural enemies and control*, vol. 3A. Elsevier, Amsterdam, The Netherlands.
- Eskafi, F. M., and M. E. Kolbe. 1990. Infestation patterns of commonly cultivated, edible fruit species by *Ceratitis capitata* and *Anastrepha* spp. (Diptera. Tephritidae) in Guatemala and their relationship to environmental factors. *Environ. Entomol.* 19: 1371–1380.
- [FAO]. Food and Agriculture Organization 1989. Avances en las investigaciones sobre moscas de las frutas en el litoral del río Uruguay. Organización de las Naciones Unidas para la Agricultura y la Alimentación. Boletín Técnico del proyecto “Control integrado de moscas de la fruta”.
- Fitt, G. P. 1989. The role of interspecific interactions in the dynamics of tephritid populations, pp. 281–300. In A. S. Robinson and G. Hooper [eds.], *Fruit flies*. Elsevier, Amsterdam, The Netherlands.
- Frissolo, M. S., C. Ambrosius, G. Muñoz, and M. Caimi. 2001. Fruit flies eradication program in La Rioja, Argentina. In *Book of Abstracts of the 4th Meeting of the Working Group on Fruit Flies of the Western Hemisphere*, 25–30 November 2001. Iscamen, Mendoza, Argentina.
- Gramajo, M. C. 2004. Comparación de la sensibilidad al frío entre *Anastrepha fraterculus* (Wied.) y *Ceratitis capitata* (Wied.) (Diptera: Tephritidae) en fruta cítrica. M.S. thesis, Facultad de Ciencias Naturales e Instituto Miguel Lillo, Universidad de Tucumán, Argentina.
- Harris, E. J., and C. Y. Lee. 1989. Development of *Ceratitis capitata* (Diptera: Tephritidae) in coffee in wet and dry habitats. *Environ. Entomol.* 18: 1042–1049.
- Hernández-Ortiz, V., J. A. Gómez-Anaya, A. Sánchez, B. A. McPheron, and M. Aluja. 2004. Morphometric analysis of Mexican and South American populations of the *Anastrepha fraterculus* complex (Diptera: Tephritidae) and recognition of a distinct Mexican morphotype. *Bull. Entomol. Res.* 94: 487–499.

- Howse, P. E., and J. J. Knapp. 1996. Pheromone of Mediterranean fruit fly: presumed mode of action and implications for improved trapping techniques, pp. 91–99. In B. A. McPherson and G. J. Steck [eds.], *Fruit fly pests: a world assessment of their biology and management*. St. Lucie Press, Delray Beach, FL.
- Knipling, G. F. 1955. Possibilities of insect control of eradication through the use of sexually sterile males. *J. Econ. Entomol.* 48: 459–462.
- Kovaleski, A., R. L. Sugayama, K. Uramoto, and A. Malavasi. 2000. Rio Grande do Sul, pp. 285–290. In A. Malavasi and R. A. Zucchi [eds.], *Moscas-das frutas de importancia economica no: conhecimento básico e aplicado*. Holos Editora, Ribeirao Preto, Brazil.
- Liquido, N. J., L. A. Shinoda, and R. T. Cunningham. 1991. Host plants of the Mediterranean fruit fly (Diptera: Tephritidae): an annotated world review. *Misc. Publ. Entomol. Soc. Am.* 77: 1–52.
- Landolt, P. J. 1985. Papaya fruit fly eggs and larvae (Diptera: Tephritidae) in field-collected papaya. *Fla. Entomol.* 68: 354–356.
- Malavasi, A., and J. S. Morgante. 1980. Biologia de “Moscas-das-frutas” (Diptera, Tephritidae). II: Indices de infestação em diferentes hospedeiros e localidades. *Rev. Bras. Biol.* 40: 17–24.
- Malavasi, A., J. S. Morgante, and R. A. Zucchi. 1980. Biologia de “Moscas-das-frutas” (Diptera, Tephritidae). I: Lista de hospedeiros e ocorrência. *Rev. Bras. Biol.* 40: 9–16.
- Malavasi, A., G. G. Rohwer, and D. S. Campbell. 1994. Fruit fly free areas: strategies to develop them, pp. 165–180. In C. O. Calkins, W. Klassen, and P. Liedo [eds.], *Fruit flies and the sterile insect technique*. CRC, Boca Raton, FL.
- Malavasi, A., R. A. Zucchi, and R. L. Sugayama. 2000. Biogeografia, pp. 93–98. In A. Malavasi and R. A. Zucchi [eds.], *Moscas-das frutas de importancia economica no Brasil: conhecimento básico e aplicado*. Holos Editora, Ribeirao Preto, Brazil.
- Nasca, A. J., J. A. Zamora, L. E. Vergara, and H. E. Jaldo. 1996. Hospederos de moscas de los frutos en el Valle de Antinaco-Los Colorados, provincia de La Rioja, Republica Argentina. *CIRPON Rev. Investig.* 10: 19–24.
- Nascimento, A. S., A. L. M. Mesquita, and R. A. Zucchi. 1984. Parasitism of pupae of *Anastrepha* spp. (Dip., Tephritidae) by *Doryctobracon areolatus* (Szépligeti, 1911) (Hymn., Braconidae) in citrus and tropical fruits, pp. 239–246. In 4th Japan-Brazil Symposium on Science and Technology, 7–9 August 1984, Annals, vol. 2, Academia de Ciências e Tecnologia do Estado de São Paulo, Sp., Brazil.
- Nora, I., E. R. Hickel, and H. F. Prando. 2000. Santa Catarina, pp. 271–275. In A. Malavasi and R. A. Zucchi [eds.], *Moscas-das frutas de importancia economica no Brasil: conhecimento básico e aplicado*. Holos Editora, Ribeirao Preto, Brazil.
- Norrbom, A. L. 2004. Fruit fly (Tephritidae) host plant database, version Nov. 2004 (<http://www.sel.barc.usda.gov:591/diptera/Tephritidae/TephHosts/search.html>).
- Ortiz, G. 1999. Potential use of the sterile insect technique against the South American fruit fly, pp. 121–130. In *The South American fruit fly, Anastrepha fraterculus* (Wied.): advances in artificial rearing, taxonomic status and biological studies. International Atomic Energy Agency, IAEA Tech-Doc 1064, Austria.
- Ovruksi, S., P. Schliserman, and M. Aluja. 2003. Native and introduced host plants of *Anastrepha fraterculus* and *Ceratitis capitata* (Diptera: Tephritidae) in northwestern Argentina. *J. Econ. Entomol.* 96: 1108–1118.
- Ovruksi, S. M., P. Schliserman, and M. Aluja. 2004. Indigenous parasitoids (Hymenoptera) attacking *Anastrepha fraterculus* and *Ceratitis capitata* (Diptera: Tephritidae) in native and exotic host plants in Northwestern Argentina. *Biol. Control* 29: 43–57.
- Putruele, M. T. G. 1993. Las moscas de las frutas. Huéspedes comprobados en el NE de Entre Ríos. *G Enfermedades, Plagas y su Control* 24: 1–4.
- Putruele, M. T. G. 1996. Hosts for *Ceratitis capitata* and *Anastrepha fraterculus* in the northeastern province of Entre Ríos, Argentina, pp. 343–345. In B. A. McPherson and G. J. Steck [eds.], *Fruit fly pests: a world assessment of their biology and management*. St. Lucie Press, Delray Beach, FL.
- Putruele, M. T. G. 1997. Dinámica poblacional de la Mosca del Mediterráneo (*C. capitata* Wied.) en el área citrícola de Concordia. *Rev. de la Asociación de Citricultores de Concordia* 34: 11–14.
- Rial, E. J. 1997. Distribución temporal y abundancia relativa de la mosca del Mediterráneo (*Ceratitis capitata*, Wied.) en el Alto Valle de Río Negro y Neuquén, p. 7. In *Memorias del I Taller de trabajo sobre los avances de investigación y apoyo científico al Programa de Control y Erradicación de Moscas de los Frutos*. Buenos Aires, Argentina.
- Sánchez, R. A., E. J. Rial, and A. P. Mongabure. 2001. Advances in the pogramme for the eradication of the Mediterranean fruit fly (*Ceratitis capitata*, Wied.) in the Patagonian region, Argentina. 2001. In *Book of Abstracts of the 4th Meeting of the Working Group on Fruit Flies of the Western Hemisphere*, 25–30 November 2001. Iscamen, Mendoza, Argentina.
- Schliserman, P., and S. M. Ovruksi. 2004. Incidencia de moscas de la fruta de importancia económica sobre *Citrus aurantium* (Rutaceae) en Tucumán, Argentina. *Manejo Integrado de Plagas y Agroecología* 72: 52–61.
- Segade, G., and L. A. Polack. 1999. Monitoreo de moscas de la fruta (Diptera: Tephritidae) en montes de duraznero y de cítricos en el Partido de San Pedro, Provincia de Buenos Aires, p. 23. In *Memorias del I Taller Internacional sobre moscas de los frutos*. PROCEN, SENASA. Buenos Aires, Argentina.
- Segura, D. F., M. T. Vera, and J. L. Cladera. 2004. Fluctuación estacional en la infestación de diversos hospedadores por la mosca del Mediterráneo, *Ceratitis capitata* (Diptera: Tephritidae), en la provincia de Buenos Aires. *Ecol. Aust.* 14: 3–17.
- Selivon, D. 2000. Relações com as plantas hospedeiras, pp. 87–91. In A. Malavasi and R. A. Zucchi [eds.], *Moscas-das frutas de importancia economica no Brasil: conhecimento básico e aplicado*. Holos Editora, Ribeirao Preto, Brazil.
- Sivinski, J., M. Aluja, J. Piñero, and M. Ojeda. 2004. Novel analysis of spatial and temporal patterns of resource use in a group of tephritid flies of the genus *Anastrepha*. *Ann. Entomol. Soc. Am.* 97: 504–512.
- StatSoft, Inc. 2000. *Statistica for Windows* (computer program manual). Statsoft, Inc., Tulsa, OK.
- Steck, G. J. 1991. Biochemical systematics and population genetic structure of *Anastrepha fraterculus* and related species (Diptera: Tephritidae). *Ann. Entomol. Soc. Am.* 84: 10–28.
- Vaccaro N. 2000. Relevamiento de *Anastrepha fraterculus* Wied. en distintos sitios del país para estudios morfológicos, p. 42. In *Memorias del II Taller Internacional sobre moscas de los frutos*. PROCEN, SENASA. Buenos Aires, Argentina.

- Vattuone, E., J. J. Cólica, E. J. Córdoba, and C. N. Palmieri. 1995. Relevamiento de las especies de "moscas de los frutos" (Diptera: Tephritidae), su distribución y fluctuación poblacional en Andalgalá, provincia de Catamarca (República Argentina.), p. 93. *In* Memorias del III Congreso Argentino de Entomología. Mendoza, Argentina.
- Vattuone, E., C. Palmieri, and R. Berlanda. 1999. Fluctuación poblacional de *C. capitata* Wied. (Diptera: Tephritidae) en plantaciones cítricas (*Citrus* spp.) en el departamento Santa Rosa, provincia de Catamarca, p. 264. *In* Memorias de la X Jornada Fitosanitarias Argentinas. Jujuy, Argentina.
- Veloso, V.R.S., P. M. Fernández, and R. A. Zucchi. 2000. Moscas-das-frutas nos estados brasileiros Goiás, pp. 247–258. *In* A. Malavasi and R. A. Zucchi [eds.], Moscas-das-frutas de importancia economica no Brasil: conhecimento básico e aplicado. Holos Editora, Ribeirao Preto, Brazil.
- Vergani, A. R. 1952. La mosca del Mediterráneo *Ceratitidis capitata*. Boletín de Sanidad Vegetal 8, N°22.
- Vergani, A. R. 1956. Distribución geográfica de las "moscas de los frutos" en la Argentina. IDIA 99: 1–5.
- White, I. M., and M. M. Elson-Harris. 1992. Fruit flies of economic significance: their identification, and bionomics. CAB International, Wallingford, United Kingdom.

Received 7 October 2004; accepted 29 August 2005.

Appendix 1. Ecological regions (Cabrera and Willink 1973, Cabrera 1976, Anonymous 1992), geographical coordinates, and altitude for each locality sampled.

Ecological region			Geographic coordinates			Climatic and ecological conditions
Domain	Province	Localities	Latitude (S)	Longitude (W)	Altitude (m)	
Amazonico	Las Yungas	Famaillá	27.03	65.25	363	Hot and wet climate; rainfall in summer and frosts during winter; 2,500–3,000 mm per year; 19.1–29.8°C mean maximum temperature; 10.1–15.6°C mean minimum temperature; vegetation of subtropical cloud forest, rich in Lauraceae and Myrtaceae.
		Horco Molle	26.91	65.08	466	
		La Rinconada	26.85	65.31	425	
		Lules	26.93	65.35	415	
		San Javier	26.85	65.33	514	
		Tafi Viejo	26.75	65.35	592	
		Taficillo	26.77	65.27	619	
		Timbó	26.70	65.13	531	
		Colonia Santa Rosa	23.45	64.36	322	
		Metán	25.48	64.95	803	
		Orán	23.13	64.33	337	
		Pichanal	23.33	64.23	303	
		Yuto	23.63	64.47	346	
		Ledesma	23.83	64.77	413	
		Perico	24.38	65.10	897	
Chaqueño	Paranaense	Caraguatay	26.66	54.75	193	Hot and rainy climate; rainfall throughout the year, 1,600–2,000 mm per year; 27.4–28.5°C mean maximum temperature; 13.0–15.3°C mean minimum temperature; vegetation of subtropical rainforest, rich in Leguminosae, Lauraceae, Meliaceae, and Myrtaceae.
		Itacuruzú	26.90	55.13	96	
		Montecarlo	26.57	54.78	161	
		Posadas	27.38	55.88	133	
		Taruna	26.70	54.66	150	
		Iruyaingó (Corrientes)	27.60	56.66	63	
		Sombrito	27.70	58.70	65	
		Embarcación	23.22	64.10	274	
		Campo Santo	24.70	65.08	850	
		Saenz Peña	26.80	60.45	60	
	Chaqueña	S.F. Valle de Catamarca	28.50	65.78	505	Hot continental climate; summer rainfalls, varying between 200 mm annually in the east to 500 mm in the west; 20.5°C–28.6°C mean maximum temperature; 13.6°C–16.1°C mean minimum temperature; vegetation characteristic of deciduous xerophytic forests, with an herbaceous stratum compose by Gramineae, Cactaceae and terrestrial Bromelaceae; also palm trees, savannahs and bushy steppes.
		Capayán	28.77	66.07	358	
		Chumbicha	28.87	66.23	377	
		San José	26.78	66.06	984	
		Santa Rosa	28.45	65.71	512	
		Sumalao	28.45	65.62	518	
		San Isidro	28.45	65.76	514	
		Villa Dolores	28.43	65.73	530	
		Luján	32.37	65.95	589	
		Merlo	32.35	65.03	808	
		Quines	32.23	65.80	482	
		de Julio	28.83	58.83	55	
		Bella Vista	28.52	59.05	58	
		Reconquista	29.11	59.42	53	
	Espinal	Monte Caseros	30.28	57.63	35	
		Chajart	30.77	57.98	55	
		Virasoro	28.50	56.17	112	

Monte	Chilecito	29.17	67.50	1080	Hot and dry climate in the north, dry and colder in the south; summery rainfalls in the north and winter rainfalls in teh south, varying from 80 to 200 mm per year; 20.4-27.0°C mean maximum temperature; 5.7-10.4°C mean minimum temperature; vegetation characteristic of deciduous xerophytic forests, palm groves, grassy savannahs, grassy steppes, and bushy steppes.
	Los Dorados	29.35	67.47	1063	
	Guandacol	29.56	68.56	1053	
	Los Sarmitos	29.15	67.52	1072	
Pampeana	Palmar	32.16	58.28	44	Hot-temperate climate: rainfall throughout the year, diminishing from north to south and from east to west, varying from 600 to 1,200 mm per year; 20.1-24.2°C mean maximum temperature; 6.9-11.8°C mean minimum temperature; vegetation of grassy steppe and xeric woodland.
	Concordia	31.38	58.02	22	
	Humaitá	31.78	58.30	44	
	La Criolla	31.15	58.06	38	
	Pto. Yerúá	31.53	58.03	35	
	Villa Adela	31.10	58.20	40	
	Villa Zorraquín	31.32	58.05	47	
	Arocena	32.08	60.97	14	
	Monje	32.38	60.93	15	
	Mercedes	34.65	59.43	117	
	Pilar	34.45	58.90	19	
	Gob. Castro	33.76	59.88	52	
	Doyle	33.46	60.10	20	
	San Pedro	33.66	59.72	15	
	Villa Adelinea	34.53	58.55	24	
	Ituzaingó (Buenos Aires)	34.67	58.67	31	