

First record of the rotifer *Lecane shieli* Segers & Sanoamuang, 1994 from Australia

Tsuyoshi Kobayashi¹, Russell J. Shiel² and Hendrik Segers³

¹Department of Environment and Climate Change NSW, PO Box A290 Sydney South, New South Wales, Australia 1232; ²Ecology and Evolutionary Biology, DX650312, University of Adelaide, South Australia, Australia; ³Royal Belgian Institute of Natural Sciences, Freshwater Biology, Vautierstraat 29, B-1000 Brussels, Belgium

Corresponding author: Yoshi.Kobayashi@environment.nsw.gov.au

ABSTRACT

The monogonont rotifer *Lecane shieli* is recorded for the first time from Australia (Bora Channel in the Macquarie Marshes, New South Wales: 30.65872° S /147.53144° E). Until recently, *L. shieli* was known only from Thailand, where the species was first described. This Australian record of a Thai 'endemic' is interesting in considering possible dispersal and biogeographical distribution.

Key words: zooplankton, monogonont rotifer, *Lecane shieli*, Macquarie Marshes, biodiversity, migratory waterbird

Wetland (or river-floodplain) systems provide important habitats for a variety of terrestrial and aquatic biota (Junk *et al.* 1989; Junk and Wantzen 2004 and references therein). An understanding of biodiversity and productivity in relation to water availability is important in ecological management of such systems. River and creek channels that run through the floodplains become an important conduit and refuge for aquatic biota in dry times (Junk and Wantzen 2004).

The Macquarie Marshes (total area: ~210,000 ha), in the north-west end of the Macquarie River catchment, form one of the largest floodplain wetlands in the Murray-Darling Basin of south-eastern Australia (Keith 2004). The Marshes comprise a mosaic of inland floodplain swamps, shrub lands, and woodlands as well as inland riverine forests (Keith 2004), and provide an important habitat for many waterbirds including colonially-nesting waterbirds (Kingsford and Thomas 1995; Kingsford and Auld 2005). The Macquarie Marshes Nature Reserve was listed as an internationally significant wetland under the Ramsar Convention in 1986 (Kingsford and Thomas 1995). Because of extended droughts exacerbated by regulation/extraction of the Macquarie River that supplies the Marshes, the presence of aquatic biota in the Marshes is confined primarily to river and creek channels where there is water.

As part of an ecological study of the Macquarie Marshes, zooplankton samples were collected in November 2006. Here we report the first record of the monogonont rotifer *Lecane shieli* from Australia. The specimen was found in a sample collected from Bora Channel in the Macquarie Marshes.

Methods

Sampling was conducted between 7 and 10 November 2006 at multiple sites along the Macquarie River, Monkeygar Creek and Bora Channel that run through the Macquarie Marshes from the southern to the northern end (Fig. 1). Duplicate water samples (10 L each) were collected at each site by submerging a plastic container in the mid channel. Zooplankton samples were concentrated by filtering through a 35 µm mesh sieve, and preserved in a 4% buffered formaldehyde solution. In the laboratory, zooplankton specimens were examined and identified (Segers 1995; Shiel 1995) under a Leica Diaplan compound microscope at a magnification of ×100 to ×320, with an image analysis system consisting of Leica DFC480 digital camera and Leica IM Version 4.0 digital imaging software (Leica Microsystems, Germany). In the field, water temperature and dissolved oxygen were measured using a YSI Model 5100 Dissolved Oxygen/Temperature Meter (YSI Inc., Ohio, USA). Duplicate water samples were also collected for measurements of conductivity (ORION Model 160 conductivity meter, Orion Research Inc., Massachusetts, USA) and turbidity (HACH 2100AN turbidimeter, Hach Company, Colorado, USA), and for analysis of nutrients in the laboratory. Nutrient analysis methods followed Greenberg *et al.* (2005).

Lecane shieli Segers & Sanoamuang, 1994

A single parthenogenetic female specimen of *Lecane shieli* was found in the sample collected on 10 November 2006 from Bora Channel of the Macquarie Marshes within the Northern Nature Reserve (30.65872° S/147.53144° E) (Figs. 2 and 3). The morphological characteristics of the specimen accord with descriptions by Segers and




Figure 1. Location of Macquarie Marshes and the sampling site on Bora Channel from which *Lecane shieli* was collected. Δ

Sanoamuang (1994). Physico-chemical conditions of the Bora Channel water were: water temperature at 0745 and 1800 hrs: 18.7 and 27.3 °C; dissolved oxygen: 5.9-6.0 mg L⁻¹; conductivity: 514-515 μ S cm⁻¹; turbidity: 8.3-8.6 NTU; total N: 980-1329 μ g L⁻¹; oxidized N: 1.8-2.0 μ g L⁻¹; ammonia N: 5.5-6.3 μ g L⁻¹; total P: 158-216 μ g L⁻¹; and filterable reactive P: 95.1-95.4 μ g L⁻¹.

The rotifer *Lecane shieli* was originally recorded and described from Thailand but never recorded elsewhere. *Lecane shieli* now occurs in Turkey (Altindag et al., pers. comm., 2007) and appears to live in southern and northern hemispheres. The species occurs in various freshwater habitats including a floodplain wetland in Thailand, with warm-water preference (Segers and Sanoamuang 1994; Savatentalinton and Segers 2005; L. Sanoamuang, pers. comm. 2007, Khon Kaen University, Thailand). Its habitats have conductivity reported in the range of 85 – 515 μ S cm⁻¹, and water temperature up to 27 °C (Segers and Sanoamuang 1994 and present study).

Several rotifer species (e.g. *Brachionus kostei*, *B. lyratus* and *Lecane batillifer*) that were previously considered endemic to Australia also occur in Thailand (Sanoamuang 1998). This first Australian record of





Figure 2. Photomicrograph of *Lecane shieli* collected from Bora Channel in Macquarie Marshes.



Figure 3. Sampling site on Bora Channel from which *Lecane shieli* was collected. View towards upstream.

Lecane shieli raises intriguing questions about possible dispersal and biogeographical distribution. Migratory waterbirds may be important potential vectors of microfauna (Santamaría and Klaassen 2002; Green and Figuerola 2005). Movements of wild birds into Australia from Asia occur every year with migratory shorebirds arriving in spring (Tracey *et al.* 2004). The Macquarie Marshes are used intermittently by at least ten species of migratory shorebirds (Kingsford and Thomas 1995). Five of them (Greenshank *Tringa nebularis*, Marsh Sandpiper *Tringa stagnatilis*, Red-necked Stint *Calidris ruficollis*, Black-tailed Godwit *Limosa limosa* and Bar-

tailed Godwit *Limosa lapponica*) occur in Thailand (Li and Mundkur 2007, pp.106-107). Aquatic invertebrates including rotifers can be transported in the gut of waterbirds, or on their feet and feathers (Frisch *et al.* in press). A recent study by Green *et al.* (in press) showed that three rotifer taxa (*Brachinous* sp., *Cephalodella* sp. and bdelloid) emerged from droppings of nomadic waterbirds (Eurasian Coot *Fulica atra* and Black Swan *Cygnus atratus*) in the Macquarie Marshes. This type of microfauna dispersal mechanism could explain the occurrence of zooplankton in the Macquarie Marshes with close affinity with south-east Asia.

Acknowledgments

We thank Jeff Kelleway, Tim Ralph and Debashish Mazmuder (Australian Nuclear Science & Technology Organisation) for help in field work; Ray Jones, Bill Johnson and Neil Saintilan for logistic support; Richard Kingsford (University of NSW), Jennifer Spencer, Rachael Thomas, Stuart Halse (Bennelongia P/L), Daniel Lunney, Renee Shepherd and Paul Rendell for useful suggestions and comments. Thanks are offered to Shannon Simpson for producing figure 1; Jeff Kelleway for providing a

photograph of the sampling site (figure 3); and Ed Czobik for conducting nutrient analyses. This work was funded by the NSW Wetland Recovery Program, which is jointly funded by the NSW Government and the Australian Government's Water Smart Australia programme, and is delivered in partnership by the DECC and the Department of Water and Energy, in association with the Central West and Border Rivers-Gwydir Catchment Management Authority.

References

- Frisch, D., Green, A.J. and Figuerola, J. (in press). High dispersal capacity of a broad spectrum of aquatic invertebrates via waterbirds. *Aquatic Sciences*
- Green, A.J. and Figuerola, J. 2005. Recent advances in the study of long-distance dispersal of aquatic invertebrates via birds. *Diversity and Distributions* 11: 149-156.
- Green, A.J., Jenkins, K.M., Bell, D., Morris, P.J. and Kingsford, R.T. (in press). The potential role of waterbirds in dispersing invertebrates and plants in arid Australia. *Freshwater Biology*
- Greenberg, A.E., Clesceri, L.S., Rice, E.W. and Franson, M.H. (eds) 2005. *Standard Methods for Examination of Water & Wastewater*. Centennial Edition, 21st Ed. American Public Health Association.
- Junk, W. J., Bayley, P.B. and Sparks, R.E. 1989. The flood pulse concept in river-floodplain systems. *Canadian Special Publication of Fisheries and Aquatic Sciences* 106: 110-127.
- Junk, W.J. and Wantzen, K.M. 2004. The Flood Pulse Concept: New Aspects, Approaches, and Applications - an Update. Pp. 117-140 in Proceedings of the Second International Symposium on the Management of Large Rivers for Fisheries Volume 2, edited by R.L. Welcomme and T. Petr. Food and Agriculture Organization & Mekong River Commission. FAO Regional Office for Asia and the Pacific, Bangkok. RAP Publication 2004/17.
- Keith, D.A. 2004. *Ocean Shores to Desert Dunes: the Native Vegetation of New South Wales and the ACT*. Department of Environment and Conservation, Hurstville, NSW.
- Kingsford, R.T. and Auld, K.M. 2005. Waterbird breeding and environmental flow management in the Macquarie Marshes, arid Australia. *River Research and Applications* 21: 187-200.
- Kingsford, R.T. and Thomas, R.F. 1995. The Macquarie Marshes in arid Australia and their waterbirds: a 50 year history of decline. *Environmental Management* 19: 867-878.
- Li, Z.W.D. and Mundkur, T. 2007. *Numbers and Distribution of Waterbirds and Wetlands in the Asia-Pacific Region. Results of the Asian Waterbird Census: 2002-2004*. Wetlands International, Kuala Lumpur, Malaysia.
- Sanoamuang, L. 1998. Rotifera of some freshwater habitats in the floodplain of the River Nan, northern Thailand. *Hydrobiologia* 387/388: 27-33.
- Santamaría, L. and Klaassen, M. 2002. Waterbird-mediated dispersal of aquatic organisms: an introduction. *Acta Oecologica* 23: 115-119.
- Savatenalinton, S. and Segers, H. 2005. Rotifers from Kalasin Province, Northeast Thailand, with notes on new and rare species. *Zoological Studies* 44: 361-367.
- Segers, H. 1995. *Rotifera. Volume 2: The Lecanidae (Monogonota). Guides to the Identification of the Microinvertebrates of the Continental Waters of the World* 6. SPB Academic Publishing, The Hague.
- Segers, H. and Sanoamuang, L. 1994. Two more new species of *Lecane* (Rotifera, Monogononta) from Thailand. *Belgian Journal of Zoology* 124: 39-46.
- Shiel, R.J. 1995. *A Guide to Identification of Rotifers, Cladocerans and Copepods from Australian Inland Waters*. Co-operative Research Centre for Freshwater Ecology Identification Guide no. 3. Murray-Darling Freshwater Research Centre, Albury, NSW.
- Tracey, J.P., Wood, R., Roshier, D., West, P. and Saunders, G.R. 2004. The role of wild birds in the transmission of avian influenza for Australia: an ecological perspective. *Emu* 104: 109-124.