



**Testi AM, Pession A, Diverio D, et al. Risk-adapted treatment of acute promyelocytic leukemia: results from the International Consortium for Childhood APL. *Blood*. 2018;132(4):405-412.**

In the visual abstract, for high-risk (HR) patients in the ICC-APL-01 trial, the information for consolidation course 3 should read "ATRA + IDA + ARA-C," not "ATRA + IDA." The corrected visual abstract is shown below.



Page 407: In Figure 1, idarubicin (IDA) should not have been listed in consolidation 1 for standard-risk and high-risk patients, cytarabine (ARA-C) should not have been listed in consolidation 2 for standard-risk and high-risk patients, and ARA-C (1 g/m<sup>2</sup>/12 hours on days 1, 2, and 3) should have been listed in consolidation 3 for high-risk patients. Also, in consolidation 2 for both standard-risk and high-risk patients, "IDA 10mg/m<sup>2</sup>/d day 4" should read "IDA 5mg/m<sup>2</sup>/d days 1,3,5." The corrected Figure 1 is shown below.



**Figure 1. ICC-APL-01 study protocol design.** 6-MP, 6-mercaptopurine; IDA, idarubicin; MTX, methotrexate; MTZ, mitoxantrone.

<https://doi.org/10.1182/blood.2022017903>

© 2023 by The American Society of Hematology