

LEUKEMIC STEM CELLS

Myeloproliferative neoplasm stem cells

Adam J. Mead¹ and Ann Mullally²

¹Medical Research Council (MRC) Molecular Haematology Unit, MRC Weatherall Institute of Molecular Medicine and NIHR Biomedical Research Centre, University of Oxford, Oxford, United Kingdom; and ²Division of Hematology, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA

Myeloproliferative neoplasms (MPNs) arise in the hematopoietic stem cell (HSC) compartment as a result of the acquisition of somatic mutations in a single HSC that provides a selective advantage to mutant HSC over normal HSC and promotes myeloid differentiation to engender a myeloproliferative phenotype. This population of somatically mutated HSC, which initiates and sustains MPNs, is termed MPN stem cells. In >95% of cases, mutations that drive the development of an MPN phenotype occur in a mutually exclusive manner in 1 of 3 genes: *JAK2*, *CALR*, or *MPL*. The

thrombopoietin receptor, *MPL*, is the key cytokine receptor in MPN development, and these mutations all activate *MPL*-*JAK*-*STAT* signaling in MPN stem cells. Despite common biological features, MPNs display diverse disease phenotypes as a result of both constitutional and acquired factors that influence MPN stem cells, and likely also as a result of heterogeneity in the HSC in which MPN-initiating mutations arise. As the MPN clone expands, it exerts cell-extrinsic effects on components of the bone marrow niche that can favor the survival and expansion of MPN stem cells

over normal HSC, further sustaining and driving malignant hematopoiesis. Although developed as targeted therapies for MPNs, current *JAK2* inhibitors do not preferentially target MPN stem cells, and as a result, rarely induce molecular remissions in MPN patients. As the understanding of the molecular mechanisms underlying the clonal dominance of MPN stem cells advances, this will help facilitate the development of therapies that preferentially target MPN stem cells over normal HSC. (*Blood*. 2017;129(12):1607-1616)

Introduction

Myeloproliferative neoplasm (MPN) stem cells are defined as a clonal population of rare cells within the bone marrow that harbor an MPN-initiating somatic mutation, are capable of indefinite self-renewal, and that, through a combination of cell-intrinsic and cell-extrinsic effects undergo clonal expansion. In this review, we focus on BCR-ABL⁻negative MPNs and highlight 5 main features of MPN stem cells: (1) MPN disease-initiating somatic mutations, (2) the effects of MPN somatic mutations on hematopoietic stem cell (HSC) function, (3) the role of MPN stem cells in the heterogeneity of disease phenotype in MPNs, (4) the MPN stem cell niche, and (5) therapeutic targeting of MPN stem cells.

MPN disease-initiating somatic mutations

The molecular basis of MPNs has been defined in almost all cases. In >95% of cases of MPNs, the mutations that drive the development of an MPN phenotype are accounted for by somatic mutations in 3 genes: *JAK2*, *CALR*, or *MPL*, and notably these mutations occur in a mutually exclusive manner.¹ Mutations in *JAK2* and *MPL* occur as gain-of-function point mutations (ie, *JAK2*V617F and *MPL*W515L/K, respectively), whereas the mutations in *CALR* occur as +1 base-pair frameshifts in the last coding exon of *CALR*, which result in the generation of a novel C-terminus.^{1,2} Recent work indicates that *CALR* mutations confer a neomorphic function on mutant *CALR* that results in activation of *MPL* signaling.³⁻⁶ Additional germ line and somatic

mutations in *JAK2* or *MPL* were recently identified in ~19% of the so-called “triple-negative” MPN cases.⁷ Previously, negative regulators of the *JAK*-*STAT* signaling pathway, including *LNK*,⁸ *c-CBL*,^{9,10} and *SOCS*,¹¹ were also shown to be somatically inactivated at low frequency in MPNs, highlighting the primacy of the *JAK*-*STAT* signaling pathway in MPN pathogenesis.

Evidence for an MPN disease-initiating role for *JAK2*, *CALR*, and *MPL* somatic mutations has been provided by retroviral bone marrow transplant assays, where ectopic expression of each mutation *alone* is sufficient to engender MPNs in mice^{5,6,12}; however, it is important to note that the MPN is polyclonal in all these models. Recent work using a transgenic mouse model where human *JAK2*V617F is expressed from the endogenous human *JAK2* promoter demonstrated that although MPN can be initiated by transplanting a single *JAK2*V617F-expressing long-term (LT)-HSC into a lethally irradiated wild-type recipient mouse, MPNs developed in only a minority of recipients in whom long-term reconstitution occurred,¹³ supporting the concept that *JAK2*V617F is not always disease initiating when modeled at the level of a single LT-HSC. The question of whether *JAK2*V617F is disease initiating in MPNs was also recently raised in the context of human hematopoiesis following a series of studies in which *JAK2*V617F mutations were detected in the peripheral blood of normal individuals who do not have any apparent hematological disease.¹⁴⁻¹⁶ This phenomenon, where clonally restricted somatic mutations in genes associated with hematological malignancies (including *JAK2*V617F) are found in normal individuals, has been termed clonal hematopoiesis of indeterminate potential (CHIP)

Submitted 24 October 2016; accepted 22 November 2016. Prepublished online as *Blood* First Edition paper, 3 February 2017; DOI 10.1182/blood-2016-10-696005.

© 2017 by The American Society of Hematology

and is strongly associated with increasing age.¹⁷ JAK2V617F is among the most common CHIP-associated mutations and in most cases the *JAK2* mutations are isolated events that occur in the absence of other hematological malignancy-associated mutations, suggesting that JAK2V617F *alone* is sufficient to engender clonal hematopoiesis. However, intriguingly, the prevalence of JAK2V617F-positive MPNs is significantly lower than that of JAK2V617F-positive CHIP, suggesting that in many cases JAK2V617F *alone* may be sufficient to engender clonal hematopoiesis but insufficient to induce MPNs. This discrepancy may be explained in part by the observation from the Copenhagen General Population Study that higher JAK2V617F allele burden is associated with development of clinical MPNs, and that a minimum threshold JAK2V617F allele burden appears to be required for the development of overt disease.¹⁸ However, because the presence of concomitant somatic mutations was not assessed in the Copenhagen study, definitive conclusions regarding the sufficiency of JAK2V617F *alone* to cause MPNs cannot be made.

It is also important to note that other somatic genetic alterations, for example, *TET2* loss-of-function mutations, can precede the acquisition of JAK2V617F as reported in the original description of *TET2* mutations in myeloid malignancies¹⁹ and later validated in subsequent studies.^{20,21} Indeed, *TET2* is also a common CHIP gene, as are other epigenetic genes, such as *DNMT3A*.¹⁴⁻¹⁶ Somatic mutations, clonally expanded HSC in which additional somatic mutations subsequently occur to drive the development of a disease phenotype have been termed “preleukemic” HSC, and their role in the development of myeloid malignancies has now been clearly established, validating the original description of *TET2* mutations preceding JAK2V617F acquisition in MPNs.¹⁹ It has also been shown that *TET2* mutations can follow JAK2V617F, and the order in which these mutations are acquired can impact both the age of onset and clinical features of MPNs.²¹

The acquisition of *TET2* mutations in established MPNs has also been associated with leukemic transformation, as have mutations in other epigenetic genes such as *ASXL1*, in splicing genes such as *SRSF2*, and in additional genes including *IDH1/2*, *TP53*, *NRAS*, and *RUNX1*.^{22,23} Interestingly, acute myeloid leukemia (AML) that arises out of JAK2V617F-mutant MPNs retains the JAK2V617F allele only ~50% of the time,^{24,25} suggesting that the acquisition of additional somatic mutations in MPN stem cells that harbor a phenotypic driver mutation (ie, *JAK2*, *CALR*, or *MPL*) is not the only route to AML. Leukemic transformation that emerges from a pre-MPN clone (lacking an MPN phenotypic driver mutation) may be promoted by inflammatory cytokines secreted by the MPN clone, particularly in the context of MF (see “The MPN stem cell niche”).

What is the impact of MPN disease-initiating somatic mutations on the cell of origin in MPNs?

A number of lines of evidence support the concept that the original target cell that acquires an MPN-initiating mutation is an LT-HSC that resides at the apex of the hematopoietic hierarchy (Figure 1). In MPN patients, the JAK2V617F mutation is detectable in immunophenotypically defined (CD34⁺ CD38⁻) LT HSC²⁶ and in all mature cell lineages,^{27,28} indicating that the mutation is acquired by a multipotent cell. By genotyping hematopoietic colonies from MPN patients, *CALR* mutations were also demonstrated to be present in the earliest phylogenetic node.¹ Additional evidence from genetic mouse models further support the idea that Jak2V617F disease-initiating cells reside exclusively in the immunophenotypically defined LT-HSC compartment,

and that Jak2V617F multipotent progenitors lack disease-propagating potential.^{29,30} Following the acquisition of an MPN-initiating mutation by a single HSC, the development of an overt MPN phenotype requires clonal expansion, conferred through a selective advantage of MPN stem cells over their normal counterparts (Figure 2). In contrast to acute leukemia, where myeloid progenitor cells can acquire aberrant self-renewal,^{31,32} disease initiation and propagation in MPNs can only be sustained by cells residing within the immunophenotypically defined LT-HSC compartment.^{30,33}

The impact of MPN phenotypic driver mutations on the expansion of the HSC pool can be inferred from the somatic mutant allele burden in granulocytes at diagnosis. In this regard, it is informative to compare the JAK2V617F mutant allele burden in chronic phase MPNs (ie, polycythemia vera [PV] and essential thrombocythemia [ET]) with that of more advanced phase disease MPNs (ie, myelofibrosis [MF]). In PV, the JAK2V617F-mutant allele burden is often low at diagnosis,³⁴ and in ET, JAK2V617F heterozygous clones can remain stable over years,³⁵ with recent evidence suggesting that homozygous JAK2V617F clones do not necessarily expand.³⁶ Concordant with this observation, in a study of PV, ET, and MF patients, expansion of the CD34⁺ CD38⁻ HSC compartment was not present in PV and ET patients but was present in MF patients.³⁷ A separate study showed that JAK2V617F allele burden was higher in the CD34⁺ cell compartment of patients with MF as compared with PV and ET.³⁸ Interestingly, this expansion of *JAK2*-mutant cells in the CD34⁺ compartment of patients with MF was found to be independent of JAK2V617F homozygosity, suggesting that clonal expansion in the CD34⁺ compartment in MF is driven by other somatic genetic alterations enriched in MF as compared with PV or ET (for example, epigenetic loss-of-function mutations). In contrast, the mutant *CALR* allele fraction measured in granulocytes is typically 40% to 50% at the time of ET diagnosis, suggesting that heterozygous mutant *CALR* HSC become clonally dominant quickly.³⁹ Consistent with this observation, the age of onset of MPNs is younger in *CALR*-mutant patients as compared with *JAK2*-mutant patients.^{39,40}

The impact of MPN-initiating mutations on HSC function has been studied in vivo using both genetic mouse models and patient-derived xenograft models. JAK2V617F in particular has been the subject of intensive study, and Jak2V617F genetic mouse models have been reviewed in detail elsewhere.^{12,41} Although these models have demonstrated that JAK2V617F alone is sufficient to engender MPNs and have shown that oncogenic JAK2V617F signaling does not confer self-renewal capacity upon non-self-renewing hematopoietic cells, the impact of JAK2V617F on HSC self-renewal (as assayed by competitive repopulation) has been highly variable, depending on the genetic targeting approach taken to generate the particular mouse model. Recently, using a transgenic JAK2V617F mouse model, the level of JAK2V617F expression in single LT-HSC was shown to influence reconstitution capacity, with high JAK2V617F-expressing single LT-HSC demonstrating an exhaustion phenotype in a serial transplantation assay, as compared with low JAK2V617F-expressing single LT-HSC, which retained their long-term repopulation capacity.¹³ Both *Tet2* and *Ezh2* loss have been shown to enhance the repopulating activity of Jak2V617F disease-propagating MPN stem cells in genetic mouse models⁴²⁻⁴⁶; in the case of *Ezh2* loss, the enhanced repopulation was mediated through reactivation of a fetal *lin28b* program.⁴⁵

MPN studies in xenografts have found that JAK2V617F-mutant CD34⁺ cells from patients with PV and ET engraft relatively poorly,⁴⁷ whereas CD34⁺ cells from either the peripheral blood or the spleens of patients with MF demonstrate sustained engraftment,^{48,49} and in the case of splenic CD34⁺ cells, serial transplantation into secondary recipients was achieved.⁴⁹ The fact that JAK2V617F severe combined




Figure 1. Key somatic mutations and growth factor receptors important for MPN development. (A) Simplified "roadmap" of hematopoietic development. (B) Distribution of key growth factor receptors in different stem, progenitor, and precursor cell populations. For each population, potential impact of *JAK2V617F*, *CALR*, or *MPL* mutation is indicated.

immunodeficiency (SCID) repopulating cells (SRC) do not gain a proliferative advantage over wild-type SRC over time in CD122-depleted NOD/SCID mice⁵⁰ suggests that *JAK2V617F* is not a strong driver of clonal expansion at the level of the HSC. However, 1 important caveat in the interpretation of these results is that incompatibilities between human cytokine receptors (expressed on transplanted CD34⁺ cells) and murine cytokines (produced by the recipient murine bone marrow) may significantly impact *JAK2V617F* SRC activity in these studies. Comutation of *TET2* with *JAK2V617F* has been shown to increase the NOD-SCID repopulating capacity of CD34⁺ cells, when compared with CD34⁺ cells harboring *JAK2V617F* alone.¹⁹ In an effort to overcome issues with species incompatibility in xenografts, a humanized bone marrow ossicle transplantation model was recently employed to study MPN stem cells.³³ In this study, the CD34⁺ compartment was fractionated into CD34⁺CD38⁺ and CD34⁺CD38^{low/-} subfractions, and engraftment was observed exclusively in ossicles transplanted with CD34⁺CD38^{low/-} cells, confirming that the MPN-initiating population is contained solely in the immunophenotypically defined LT-HSC fraction.³³

Finally, the impact of cytokine receptor signaling in MPN stem cells has been assessed using various mouse models. In the HSC compartment, the thrombopoietin receptor, *Mpl*, is a key growth factor receptor that is required for normal HSC function,⁵¹ and importantly, all MPN-initiating mutations activate *MPL* signaling (Figure 1). Using genetic mouse models, a critical role for *Mpl* in *Jak2V617F*-driven MPN initiation has been demonstrated.⁵² A similar requirement for *Mpl* in mutant *CALR*-driven MPNs initiation was recently demonstrated using a retroviral bone marrow transplant model.⁶ Conversely, *IL3Rb* signaling has been shown to be expendable for MPN initiation,⁵³ and in the case of erythropoietin signaling, restricting *Jak2V617F* expression

to erythropoietin receptor (EpoR) expressing erythroid precursor cells has been found to result in a markedly attenuated MPN phenotype, supporting the absence of a role for the EpoR signaling in *Jak2V617F*-driven MPN initiation.³⁰ Although these data firmly establish *MPL* as the crucial growth factor receptor for MPN development, we still have an incomplete understanding of the molecular events downstream of activated *MPL* signaling that allow MPN stem cells to gain a clonal advantage in the bone marrow.

How do MPN stem cells contribute to heterogeneity of disease phenotype in MPNs?

Although MPNs have many clinical and biological features in common, including activation of JAK-STAT signaling, they also exhibit a range of distinct clinical phenotypes. Understanding the factors that influence the impact of an MPN disease-initiating mutation on the HSC that acquires it, and on its clonal progeny, is of fundamental importance in understanding heterogeneity of disease phenotype in MPNs, and these factors are summarized in Figure 3. First, it is clear that MPN phenotype is partly determined by the specific phenotypic driver mutation that is acquired. For example, *CALR* and *MPL* mutations are almost always associated with an ET or MF phenotype, but not a PV phenotype. The association of distinct MPN phenotypes with *CALR* or *MPL* mutations partly relates to the mechanism by which these mutations induce MPNs, namely through activating signaling through *MPL*,^{3-6,54} which is selectively expressed on HSC and cells in the megakaryocyte differentiation pathway (Figure 1). Activation of this pathway consequently promotes megakaryopoiesis and a




Figure 2. Key steps during MPN development from normal hematopoiesis following acquisition of an MPN-initiating mutation in a single HSC. The mutant HSC acquires a selective advantage over normal HSC and also promotes myeloid differentiation, eventually leading to a myeloproliferative phenotype. The expanded, abnormal myeloid clone disrupts the bone marrow microenvironment, promoting a self-reinforcing malignant niche that favors MPN stem cells over normal HSC and leads to eventual mobilization of MPN HSC into the peripheral blood (PB).

thrombocytosis phenotype. Specific signaling events downstream of a driver mutation can also influence MPN phenotype, through selectively promoting specific lineage differentiation, for example, *JAK2* exon 12 mutations, which activate *JAK2* signaling are associated with an isolated erythrocytosis phenotype both in MPN

patients⁵⁵ and in mouse models.⁵⁶ Whether this lineage specificity occurs as a result of instructive signaling in multipotent progenitors, for example, by differentially enhancing erythropoietin signaling,⁵⁷ or as a consequence of selective signaling in committed erythroid precursor cells, remains unclear.⁵⁸




Figure 3. Summary of the factors that influence phenotypic heterogeneity in MPNs.

In contrast to the specific clinical phenotypes associated with *CALR*, *MPL*, and *JAK2* exon 12 mutations, an intriguing aspect of MPN biology relates to the striking heterogeneity in clinical phenotype associated with the more common *JAK2V617F* mutation, which is associated with PV, ET, and MF phenotypes in humans and also in mouse models.^{12,41} An unresolved question is why a *JAK2V617F* mutation acquired by a single HSC can result in such a variety of clinical phenotypes. Furthermore, recent studies identifying the presence of CHIP in normal individuals¹⁴⁻¹⁶ indicate that many individuals who acquire a *JAK2V617F* mutation in an HSC will never develop MPN. A number of possible explanations have been proposed for *JAK2V617F*-associated clinical heterogeneity. First, evidence from MPN patients and mouse models indicates that phenotypic diversity in *JAK2V617F*-driven MPNs relates to both quantitative differences in *JAK2V617F* allele burden^{36,59} and qualitative differences in signaling downstream of *JAK2V617F*, including levels of STAT1 activation.⁶⁰ Second, other modifier genes, through effects either on HSC or on the bone marrow microenvironment, could influence disease phenotype, for example, constitutional differences in MYB expression.⁶¹ It is interesting to note, however, that the same germ-line predisposition loci are associated with both *JAK2V617F*-positive CHIP and *JAK2V617F*-positive MPNs.⁶² Third, the presence of other somatic mutations, for example, in *TET2*, and the order in which the mutations are acquired in HSC, have been shown to influence the resulting disease phenotype.²¹ Fourth, cell-extrinsic environmental factors could influence eventual MPN phenotype. Finally, phenotypic heterogeneity could relate, at least in part, to the specific cell of origin of MPNs.

A number of lines of evidence suggest that HSC are highly heterogeneous in their self-renewal capacity and their intrinsic lineage bias.⁶³ For example, platelet-biased HSC that reside at the apex of the hematopoietic hierarchy have been characterized.⁶⁴ Furthermore, single-cell transplantation studies in mice have identified myeloid lineage-restricted, self-renewing progenitors,⁶⁵ and a similar cell population has also been described in the context of native non-transplant hematopoiesis.⁶⁶ Thus, characteristics of the specific individual HSC cell of origin that acquires a *JAK2V617F* mutation could also contribute to the phenotypic heterogeneity of *JAK2V617F*-positive MPNs and may provide an explanation, at least in part, for the development of *JAK2V617F*-positive CHIP without progression to MPNs. With respect to phenotypic heterogeneity, a *JAK2V617F* mutation arising in a platelet-biased HSC might result in an ET phenotype (Figure 4A), whereas the same mutation arising in a myeloid lineage-balanced HSC might promote a PV phenotype (Figure 4B). A recent study using single-cell transplantation approaches to analyze a *JAK2V617F* transgenic mouse model provides some evidence in support of this hypothesis.¹³ Single-stem/progenitor cells from genetically identical donor mice were found to be capable of generating MPNs in vivo with markedly variable phenotypes. There is also evidence to indicate that some HSC possess more limited self-renewal capability than others,⁶³ and it may be that a *JAK2V617F* mutation arising in such a cell could result in CHIP (Figure 4C), with a clinical phenotype emerging only upon acquisition of a second mutation, for example in *TET2*, that enhances the self-renewal of the antecedent *JAK2V617F*-mutant clone⁴² (Figure 4D). Acquisition of a *JAK2V617F* mutation in an HSC already harboring a *TET2* mutation could influence the clinical phenotype,^{21,67} in addition to the age of MPN onset²¹ (Figure 4E). Given the increased prevalence of CHIP in older individuals, it is interesting to note that aging has significant effects on HSC heterogeneity, including an increase in myeloid-biased differentiation, a decreased output of differentiated cells, and a reduced capacity for self-renewal in long-term secondary transplantation assays.^{68,69}

The MPN stem cell niche

The bone marrow niche provides the microenvironmental signals that are essential for HSC function,⁷⁰ and progress in understanding the cellular composition of the HSC niche has facilitated investigation of how these different components are perturbed during the development of myeloid neoplasms.⁷¹ Indeed, a number of lines of evidence indicate that targeted disruption of the bone marrow niche *alone* is sufficient to induce an MPN phenotype in vivo.⁷²⁻⁷⁵

Although the initiating MPN mutation acquired by a single HSC is likely to occur in the context of an unperturbed niche, as the MPN progresses, the expanding malignant clone exerts cell-extrinsic effects on components of the bone marrow niche, and evidence from mouse models suggests that this niche disruption can favor the survival and expansion of MPN stem cells over normal HSC, resulting in a so-called malignant “self-reinforcing” niche.⁷⁶ For example, *JAK2V617F*-positive HSC have been shown to induce bone marrow neural damage through aberrant production of interleukin-1b, which in turn leads to a reduction in mesenchymal stem cell numbers and acceleration of MPN progression,⁷⁷ culminating in the development of a clinical trial to test the efficacy of β 3-adrenergic agonists in MPNs. Indeed, more advanced MPNs are characterized by the development of bone marrow fibrosis that is promoted by proinflammatory cytokines, including transforming growth factor β and tumor necrosis factor α (TNF α), which are produced by abnormal populations of myelomonocytic cells and megakaryocytes.^{76,78} Both transforming growth factor β and TNF α are also known negative regulators of normal HSC,^{79,80} and in a retroviral *JAK2V617F* model, TNF α was demonstrated to facilitate the preferential expansion of *JAK2V617F*-mutant hematopoietic cells.⁸¹ The bone marrow niche is also perturbed in MPNs as a result of enhanced angiogenesis as evidenced by increased bone marrow microvessel density in MPN patients,⁸² possibly occurring as a consequence of increased vascular endothelial growth factor expression.⁸³ The disrupted niche can also lead to mobilization of HSC into peripheral blood and spleen, resulting in splenic extramedullary hematopoiesis, which is a cardinal feature of more advanced MPNs.⁴⁹ Furthermore, the microenvironmental disruption of the bone marrow in MPNs might also promote leukemic transformation as demonstrated by an MF-xenograft study in which a high frequency of AML of *mouse origin* was noted. In this study, murine AML development was confined to the MF-xenograft group (and was not seen in xenografts of normal bone marrow or cord blood cells), suggesting that paracrine signaling arising from the MF cells drove leukemogenesis in normal mouse cells in vivo.⁸⁴ Recent work has highlighted lipocalin as an important paracrine factor in promoting DNA damage and disease evolution in MPNs.^{85,86}

Finally, a number of pathways mediating bone marrow fibrosis, inflammatory signals, and niche dysregulation are potentially amenable to therapeutic targeting. New approaches to modeling niche disruption such as those using humanized bone marrow ossicles in xenograft assays will be helpful in further exploring the underlying mechanisms of niche disruption in MPNs and in testing novel therapeutic interventions.³³

Therapeutic targeting of MPN stem cells

The finding that MPNs originate from and are propagated by rare self-renewing stem cells that closely resemble normal LT-HSC phenotypically represents both a challenge and a therapeutic opportunity. The opportunity lies in the fact that effectively targeting MPN




Figure 4. Disease heterogeneity following acquisition of the JAK2V617F mutation in a single HSC. X-axis represents time following acquisition of JAK2V617F indicated by the arrow. Y-axis represents relative contribution from this HSC clone to each lineage, indicated by color. (A) JAK2V617F occurs in a platelet-biased HSC resulting in ET. (B) JAK2V617F occurs in a lineage-balanced HSC resulting in PV with trilineage myeloproliferation. (C) JAK2V617F occurs in an HSC with limited self-renewal capability resulting in CHIP. (D) JAK2V617F precedes acquisition of a TET2 mutation resulting in a PV phenotype. (E) TET2 precedes acquisition of a JAK2V617F mutation resulting in an ET phenotype. (A-C) represent hypotheses to explain how HSC heterogeneity may influence MPN phenotype. (D-E) represent an interpretation of published data demonstrating that the order in which JAK2V617F and TET2 mutations are acquired influences MPN phenotype.

disease-propagating stem cells is not only required but is also likely to be sufficient to eradicate the disease and achieve a cure. However, in myeloid malignancies, stem cell populations have been demonstrated to be selectively resistant to therapies that are able to induce cytogenetic remissions⁸⁷ and even the most successful targeted therapies in chronic myelogenous leukemia (CML) fail to eradicate leukemia stem cells in the majority of patients.⁸⁸ This selective therapeutic resistance of leukemia stem cell populations is, at least in part, related to their marked quiescence in comparison with progenitor populations.^{89,90} Nevertheless, as somatic mutations confer a competitive advantage to MPN stem cells, the pathways that mediate this selective advantage should, in principle, be amenable to therapeutic targeting.

The only current curative therapy for MPNs, and therefore the only treatment that is able to eradicate MPN stem cells, is allogeneic HSC transplant. However, even following allogeneic transplant, relapse is unfortunately not an infrequent occurrence,⁹¹ indicating that the MPN stem cell populations can be resistant even to high-dose conditioning chemotherapy and alloimmunity-mediated clearance. Furthermore, due to the toxicity associated with the transplantation procedure, this approach is currently restricted to a minority of younger patients with advanced MPNs.⁹² Recently, CRISPR/Cas9 gene editing has emerged as a promising new approach with the potential to treat human disease, and although an enormous amount of work will be required to demonstrate the safety and feasibility of gene-editing as a therapeutic

approach in patients, the recent demonstration that the causative mutation for sickle cell disease could be repaired in HSC from patients with sickle cell disease⁹³ is an important advance. However, whether such a strategy could be used to treat clonal disorders such as MPNs by repairing JAK2V617F or MPLW515L/K mutations in HSC is uncertain at this point. Clearly, the challenges of applying this approach to a clonal disorder are considerably greater, as disease eradication cannot be achieved unless all MPN stem cells are successfully repaired.

The clinical development of JAK2 inhibitor therapies heralded a new era of molecular targeted therapy for BCR-ABL–negative MPNs. Ruxolitinib, the first clinically approved JAK2 inhibitor, has considerable efficacy in patients with PV and MF, with improvements in blood counts, spleen size, and symptoms. Despite this success, JAK2 inhibitors have been disappointing in their ability to induce molecular remissions in MPN patients, indicating that JAK2 inhibitors do not preferentially target MPN cells over normal cells, as evidenced by a lack of significant reduction in the JAK2V617F allele burden in patients undergoing treatment. Recently, the 4-year data from the COMFORT-I study, a phase 3 trial in MF, was reported. Only 12% of patients showed a >50% decrease in JAK2V617F allele burden, and <2% of patients achieved complete molecular remission.⁹⁴ A retrospective analysis of *CALR*-mutant patients treated with ruxolitinib on the COMFORT-II study (another phase 3 trial in MF) was recently reported.⁹⁵ Although *CALR*-mutant patients demonstrated comparable clinical responses to *JAK2*-mutant patients, there was no significant change in the *CALR*-mutant allele burden after a median of 60 weeks of treatment with ruxolitinib (n = 18). Underlying the failure of JAK2 inhibitors to induce molecular remission in BCR-ABL–negative MPNs is the lack of efficacy of these agents to target MPN stem cells, which is supported by evidence from both mouse models²⁹ and MF patient samples.⁹⁶ Encouragingly, complete inhibition of JAK2 in a retroviral MPLW515L MPN mouse model using a genetic knockout approach resulted in a superior therapeutic response than treatment with a JAK2 inhibitor.⁹⁷ A new generation of type II JAK2 inhibitors are able to induce more effective JAK2 inhibition,⁹⁸ and if developed for clinical use, such agents may be more successful at targeting MPN stem cells. However, JAK2 signaling is also essential for normal HSC function, as evidenced by several studies that have shown that hematopoietic-specific conditional genetic deletion of *Jak2* in adult mice results in severe cell-intrinsic defects in HSC function, impaired hematopoiesis, and reduced survival.^{99–101} Although total loss of JAK2 protein as occurred in the case of genetic knockout mouse models may be different than more potent JAK2 kinase inhibition, these murine studies have raised concerns regarding the potential for on-target hematological toxicity from more potent JAK2 inhibitors. Recent work focused on the crystal structure and biochemical properties of the pseudokinase domain of JAK2, which is the domain in which the V617F mutation is found, may help to overcome this issue by advancing the development of JAK2V617F-mutant specific inhibitors.^{102,103}

Interferon has been used in the treatment of MPNs for decades,¹⁰⁴ and more recently, it has been recognized that a minority of interferon-treated patients achieve molecular remissions with interferon treatment, with reemergence of polyclonal hematopoiesis in some.^{105,106} Interferon treatment is not a “molecularly targeted” therapy, and its ability to induce molecular responses applies across a range of different hematological malignancies,¹⁰⁷ including patients with CML and MPN patients with *JAK2V617F*^{105,106} and *CALR* mutations.¹⁰⁸ Although acknowledging that interferon treatment likely has multiple effects in MPN patients, potentially including immunological effects and an

impact on the bone marrow microenvironment, 1 possible mechanism by which it achieves molecular remissions in MPNs may relate to its ability to stimulate normally quiescent stem cell populations into cycle. As this cycling is more marked in JAK2V617F-positive HSC, interferon leads to preferential depletion of MPN stem cells in *Jak2V617F* mouse models,^{109,110} a finding that has also been corroborated in primary MPN samples.¹¹¹ Long-term molecular remissions after discontinuation of interferon therapy have been reported, suggesting that JAK2V617F-mutant HSC can be eradicated by interferon.¹¹² However, molecular relapses after cessation of interferon have also been observed,¹¹³ supporting the concept that in some patients, MPN stem cells persist and are able to mediate relapse. It should also be remembered that conventional chemotherapy agents such as busulfan have also been shown to induce molecular remissions in a small number of patients with PV,¹¹⁴ suggesting that the potential of conventional treatments to eradicate MPN stem cells in rare patients may not be restricted to interferon.

A number of agents in clinical development are showing promise with regards to their ability to target MPN stem cells. For example, in vitro treatment of JAK2V617F-positive CD34⁺ cells with MDM2 inhibitors (alone or in combination with interferon) has been found to reduce the degree of donor-derived chimerism and the JAK2V617F allele burden following injection of the treated CD34⁺ cells into xenografts, suggesting that the treatment targets MPN stem cells.¹¹⁵ Preclinical genetic mouse models have also identified selective sensitivity of MPN progenitors to estrogen-induced apoptosis,¹¹⁶ prompting initiation of a clinical trial (ISRCTN65011803) to test the efficacy of tamoxifen in MPNs, with a primary endpoint focused on molecular response. As research studies continue to advance the understanding of the mechanisms by which MPN stem cells become clonally dominant, the potential to exploit the biological insights gained in order to selectively target MPN stem cells in patients will become more feasible.

Future directions

Although tremendous progress has been made in the field of MPNs, particularly with regard to our understanding of the genetic basis of MPNs and to the development of molecularly targeted therapies with JAK2 inhibitors, significant gaps remain in our understanding of the mechanisms by which MPN stem cells become clonally dominant in the bone marrow. Furthermore, recent studies on CHIP have highlighted additional deficits in our understanding of the factors that constrain and promote the transition from JAK2V617F-driven clonal hematopoiesis to JAK2V617F-induced MPNs. This incomplete understanding of MPN stem cell biology underlies, at least in the part, the failure of current therapies to effectively target MPN stem cells, an essential step toward achieving cure. Progress in this area is further impeded by the failure of most clinical trials to incorporate appropriate biological sample banking that would allow future analysis of disease stem cells. Incorporation into clinical studies of biological secondary endpoints that focus on analysis of the stem cell compartment, such as assessment of molecular response in HSC, will provide an invaluable resource for future studies. Ultimately, effective eradication of MPN stem cells may require a multitargeted approach, tackling both the cell-intrinsic and the cell-extrinsic mechanisms that support MPN stem cell propagation.

Acknowledgment

The authors thank Julie-Aurore Losman for critically reviewing the manuscript.

Authorship

Contribution: A.J.M. and A.M. coauthored this review article.

References

- Nangalia J, Massie CE, Baxter EJ, et al. Somatic CALR mutations in myeloproliferative neoplasms with nonmutated JAK2. *N Engl J Med*. 2013; 369(25):2391-2405.
- Klampff T, Gisslinger H, Harutyunyan AS, et al. Somatic mutations of calreticulin in myeloproliferative neoplasms. *N Engl J Med*. 2013;369(25):2379-2390.
- Araki M, Yang Y, Masubuchi N, et al. Activation of the thrombopoietin receptor by mutant calreticulin in CALR-mutant myeloproliferative neoplasms. *Blood*. 2016;127(10):1307-1316.
- Chachoua I, Pecquet C, El-Khoury M, et al. Thrombopoietin receptor activation by myeloproliferative neoplasm associated calreticulin mutants. *Blood*. 2016;127(10): 1325-1335.
- Elf S, Abdelfattah NS, Chen E, et al. Mutant calreticulin requires both its mutant C-terminus and the thrombopoietin receptor for oncogenic transformation. *Cancer Discov*. 2016;6(4): 368-381.
- Marty C, Pecquet C, Nivarthi H, et al. Calreticulin mutants in mice induce an MPL-dependent thrombocytosis with frequent progression to myelofibrosis. *Blood*. 2016;127(10):1317-1324.
- Milosevic Feenstra JD, Nivarthi H, Gisslinger H, et al. Whole-exome sequencing identifies novel MPL and JAK2 mutations in triple-negative myeloproliferative neoplasms. *Blood*. 2016; 127(3):325-332.
- Oh ST, Simonds EF, Jones C, et al. Novel mutations in the inhibitory adaptor protein LNK drive JAK-STAT signaling in patients with myeloproliferative neoplasms. *Blood*. 2010; 116(6):988-992.
- Grand FH, Hidalgo-Curtis CE, Ernst T, et al. Frequent CBL mutations associated with 11q acquired uniparental disomy in myeloproliferative neoplasms. *Blood*. 2009;113(24):6182-6192.
- Sanada M, Suzuki T, Shih LY, et al. Gain-of-function of mutated C-CBL tumour suppressor in myeloid neoplasms. *Nature*. 2009;460(7257): 904-908.
- Vainchenker W, Delhommeau F, Constantinescu SN, Bernard OA. New mutations and pathogenesis of myeloproliferative neoplasms. *Blood*. 2011;118(7): 1723-1735.
- Mullally A, Lane SW, Brumme K, Ebert BL. Myeloproliferative neoplasm animal models. *Hematol Oncol Clin North Am*. 2012;26(5): 1065-1081.
- Lundberg P, Takizawa H, Kubovcakova L, et al. Myeloproliferative neoplasms can be initiated from a single hematopoietic stem cell expressing JAK2-V617F. *J Exp Med*. 2014;211(11): 2213-2230.
- Genovese G, Kähler AK, Handsaker RE, et al. Clonal hematopoiesis and blood-cancer risk inferred from blood DNA sequence. *N Engl J Med*. 2014;371(26):2477-2487.
- Jaiswal S, Fontanillas P, Flannick J, et al. Age-related clonal hematopoiesis associated with adverse outcomes. *N Engl J Med*. 2014;371(26): 2488-2498.
- Xie M, Lu C, Wang J, et al. Age-related mutations associated with clonal hematopoietic expansion and malignancies. *Nat Med*. 2014; 20(12):1472-1478.
- Steensma DP, Bejar R, Jaiswal S, et al. Clonal hematopoiesis of indeterminate potential and its distinction from myelodysplastic syndromes. *Blood*. 2015;126(1):9-16.
- Nielsen C, Bojesen SE, Nordestgaard BG, Kofoed KF, Birgens HS. JAK2V617F somatic mutation in the general population: myeloproliferative neoplasm development and progression rate. *Haematologica*. 2014;99(9): 1448-1455.
- Delhommeau F, Dupont S, Della Valle V, et al. Mutation in TET2 in myeloid cancers. *N Engl J Med*. 2009;360(22):2289-2301.
- Lundberg P, Karow A, Nienhold R, et al. Clonal evolution and clinical correlates of somatic mutations in myeloproliferative neoplasms. *Blood*. 2014;123(14):2220-2228.
- Ortmann CA, Kent DG, Nangalia J, et al. Effect of mutation order on myeloproliferative neoplasms. *N Engl J Med*. 2015;372(7):601-612.
- Zhang SJ, Rampal R, Manshoury T, et al. Genetic analysis of patients with leukemic transformation of myeloproliferative neoplasms shows recurrent SRSF2 mutations that are associated with adverse outcome. *Blood*. 2012; 119(19):4480-4485.
- Rampal R, Ahn J, Abdel-Wahab O, et al. Genomic and functional analysis of leukemic transformation of myeloproliferative neoplasms. *Proc Natl Acad Sci USA*. 2014;111(50): E5401-E5410.
- Campbell PJ, Baxter EJ, Beer PA, et al. Mutation of JAK2 in the myeloproliferative disorders: timing, clonality studies, cytogenetic associations, and role in leukemic transformation. *Blood*. 2006; 108(10):3548-3555.
- Theocharides A, Boissinot M, Girodon F, et al. Leukemic blasts in transformed JAK2-V617F-positive myeloproliferative disorders are frequently negative for the JAK2-V617F mutation. *Blood*. 2007;110(1):375-379.
- Jamieson CH, Gotlib J, Durocher JA, et al. The JAK2 V617F mutation occurs in hematopoietic stem cells in polycythemia vera and predisposes toward erythroid differentiation. *Proc Natl Acad Sci USA*. 2006;103(16):6224-6229.
- Ishii T, Bruno E, Hoffman R, Xu M. Involvement of various hematopoietic-cell lineages by the JAK2V617F mutation in polycythemia vera. *Blood*. 2006;108(9):3128-3134.
- Delhommeau F, Dupont S, Tonetti C, et al. Evidence that the JAK2 G1849T (V617F) mutation occurs in a lymphomyeloid progenitor in polycythemia vera and idiopathic myelofibrosis. *Blood*. 2007;109(1):71-77.
- Mullally A, Lane SW, Ball B, et al. Physiological Jak2V617F expression causes a lethal myeloproliferative neoplasm with differential effects on hematopoietic stem and progenitor cells. *Cancer Cell*. 2010;17(6):584-596.
- Mullally A, Poveromo L, Schneider RK, Al-Shahrour F, Lane SW, Ebert BL. Distinct roles for long-term hematopoietic stem cells and erythroid precursor cells in a murine model of Jak2V617F-mediated polycythemia vera. *Blood*. 2012;120(1):166-172.
- Krivtsov AV, Twomey D, Feng Z, et al. Transformation from committed progenitor to leukaemia stem cell initiated by MLL-AF9. *Nature*. 2006;442(7104):818-822.
- Goardon N, Marchi E, Atzberger A, et al. Coexistence of LMPP-like and GMP-like leukemia stem cells in acute myeloid leukemia. *Cancer Cell*. 2011;19(1):138-152.
- Reinisch A, Thomas D, Corces MR, et al. A humanized bone marrow ossicle xenotransplantation model enables improved engraftment of healthy and leukemic human hematopoietic cells. *Nat Med*. 2016;22(7): 812-821.
- Dupont S, Massé A, James C, et al. The JAK2 617V>F mutation triggers erythropoietin hypersensitivity and terminal erythroid amplification in primary cells from patients with polycythemia vera. *Blood*. 2007;110(3): 1013-1021.
- Gale RE, Allen AJ, Nash MJ, Linch DC. Long-term serial analysis of X-chromosome inactivation patterns and JAK2 V617F mutant levels in patients with essential thrombocythemia show that minor mutant-positive clones can remain stable for many years. *Blood*. 2007; 109(3):1241-1243.
- Godfrey AL, Chen E, Pagano F, et al. JAK2V617F homozygosity arises commonly and recurrently in PV and ET, but PV is characterized by expansion of a dominant homozygous subclone. *Blood*. 2012;120(13):2704-2707.
- Anand S, Stedham F, Beer P, et al. Effects of the JAK2 mutation on the hematopoietic stem and progenitor compartment in human myeloproliferative neoplasms. *Blood*. 2011;118(1):177-181.
- Stein BL, Williams DM, Rogers O, Isaacs MA, Spivak JL, Moliterno AR. Disease burden at the progenitor level is a feature of primary myelofibrosis: a multivariable analysis of 164 JAK2 V617F-positive myeloproliferative neoplasm patients. *Exp Hematol*. 2011;39(1): 95-101.
- Rumi E, Pietra D, Ferretti V, et al; Associazione Italiana per la Ricerca sul Cancro Gruppo Italiano Malattie Mieloproliferative Investigators. JAK2 or CALR mutation status defines subtypes of essential thrombocythemia with substantially

- different clinical course and outcomes. *Blood*. 2014;123(10):1544-1551.
40. Rumi E, Pietra D, Pascutto C, et al; Associazione Italiana per la Ricerca sul Cancro Gruppo Italiano Malattie Mieloproliferative Investigators. Clinical effect of driver mutations of JAK2, CALR, or MPL in primary myelofibrosis. *Blood*. 2014;124(7):1062-1069.
 41. Li J, Kent DG, Chen E, Green AR. Mouse models of myeloproliferative neoplasms: JAK of all grades. *Dis Model Mech*. 2011;4(3):311-317.
 42. Chen E, Schneider RK, Breyfogle LJ, et al. Distinct effects of concomitant Jak2V617F expression and Tet2 loss in mice promote disease progression in myeloproliferative neoplasms. *Blood*. 2015;125(2):327-335.
 43. Kameda T, Shide K, Yamaji T, et al. Loss of TET2 has dual roles in murine myeloproliferative neoplasms: disease sustainer and disease accelerator. *Blood*. 2015;125(2):304-315.
 44. Sashida G, Wang C, Tomioka T, et al. The loss of Ezh2 drives the pathogenesis of myelofibrosis and sensitizes tumor-initiating cells to bromodomain inhibition. *J Exp Med*. 2016;213(8):1459-1477.
 45. Shimizu T, Kubovcakova L, Nienhold R, et al. Loss of Ezh2 synergizes with JAK2-V617F in initiating myeloproliferative neoplasms and promoting myelofibrosis. *J Exp Med*. 2016;213(8):1479-1496.
 46. Yang Y, Akada H, Nath D, Hutchison RE, Mohi G. Loss of Ezh2 cooperates with Jak2V617F in the development of myelofibrosis in a mouse model of myeloproliferative neoplasm. *Blood*. 2016;127(26):3410-3423.
 47. Ishii T, Zhao Y, Sozer S, et al. Behavior of CD34+ cells isolated from patients with polycythemia vera in NOD/SCID mice. *Exp Hematol*. 2007;35(11):1633-1640.
 48. Xu M, Bruno E, Chao J, et al. The constitutive mobilization of bone marrow-repopulating cells into the peripheral blood in idiopathic myelofibrosis. *Blood*. 2005;105(4):1699-1705.
 49. Wang X, Prakash S, Lu M, et al. Spleens of myelofibrosis patients contain malignant hematopoietic stem cells. *J Clin Invest*. 2012;122(11):3888-3899.
 50. James C, Mazurier F, Dupont S, et al. The hematopoietic stem cell compartment of JAK2V617F-positive myeloproliferative disorders is a reflection of disease heterogeneity. *Blood*. 2008;112(6):2429-2438.
 51. Qian H, Buza-Vidas N, Hyland CD, et al. Critical role of thrombopoietin in maintaining adult quiescent hematopoietic stem cells. *Cell Stem Cell*. 2007;1(6):671-684.
 52. Sangkhae V, Etheridge SL, Kaushansky K, Hitchcock IS. The thrombopoietin receptor, MPL, is critical for development of a JAK2V617F-induced myeloproliferative neoplasm. *Blood*. 2014;124(26):3956-3963.
 53. Vu T, Austin R, Kuhn CP, et al. Jak2V617F driven myeloproliferative neoplasm occurs independently of interleukin-3 receptor beta common signaling. *Haematologica*. 2016;101(3):e77-e80.
 54. Pikman Y, Lee BH, Mercher T, et al. MPLW515L is a novel somatic activating mutation in myelofibrosis with myeloid metaplasia. *PLoS Med*. 2006;3(7):e270.
 55. Scott LM, Tong W, Levine RL, et al. JAK2 exon 12 mutations in polycythemia vera and idiopathic erythrocytosis. *N Engl J Med*. 2007;356(5):459-468.
 56. Grisouard J, Li S, Kubovcakova L, et al. JAK2 exon 12 mutant mice display isolated erythrocytosis and changes in iron metabolism favoring increased erythropoiesis. *Blood*. 2016;128(6):839-851.
 57. Grover A, Mancini E, Moore S, et al. Erythropoietin guides multipotent hematopoietic progenitor cells toward an erythroid fate. *J Exp Med*. 2014;211(2):181-188.
 58. Enver T, Jacobsen SE. Developmental biology: Instructions writ in blood. *Nature*. 2009;461(7261):183-184.
 59. Tiedt R, Hao-Shen H, Sobas MA, et al. Ratio of mutant JAK2-V617F to wild-type Jak2 determines the MPD phenotypes in transgenic mice. *Blood*. 2008;111(8):3931-3940.
 60. Chen E, Beer PA, Godfrey AL, et al. Distinct clinical phenotypes associated with JAK2V617F reflect differential STAT1 signaling. *Cancer Cell*. 2010;18(5):524-535.
 61. Tapper W, Jones AV, Kralovics R, et al. Genetic variation at MECOM, TERT, JAK2 and HBS1L-MYB predisposes to myeloproliferative neoplasms. *Nat Commun*. 2015;6:6691-6702.
 62. Hinds DA, Barnholt KE, Mesa RA, et al. Germ line variants predispose to both JAK2 V617F clonal hematopoiesis and myeloproliferative neoplasms. *Blood*. 2016;128(8):1121-1128.
 63. Eaves CJ. Hematopoietic stem cells: concepts, definitions, and the new reality. *Blood*. 2015;125(17):2605-2613.
 64. Sanjuan-Pla A, Macaulay IC, Jensen CT, et al. Platelet-biased stem cells reside at the apex of the haematopoietic stem-cell hierarchy. *Nature*. 2013;502(7470):232-236.
 65. Yamamoto R, Morita Y, Oeohara J, et al. Clonal analysis unveils self-renewing lineage-restricted progenitors generated directly from hematopoietic stem cells. *Cell*. 2013;154(5):1112-1126.
 66. Sun J, Ramos A, Chapman B, et al. Clonal dynamics of native haematopoiesis. *Nature*. 2014;514(7522):322-327.
 67. Saint-Martin C, Leroy G, Delhommeau F, et al; French Group of Familial Myeloproliferative Disorders. Analysis of the ten-eleven translocation 2 (TET2) gene in familial myeloproliferative neoplasms. *Blood*. 2009;114(8):1628-1632.
 68. Dykstra B, Olthof S, Schreuder J, Ritsema M, de Haan G. Clonal analysis reveals multiple functional defects of aged murine hematopoietic stem cells. *J Exp Med*. 2011;208(13):2691-2703.
 69. Grover A, Sanjuan-Pla A, Thongjuea S, et al. Single-cell RNA sequencing reveals molecular and functional platelet bias of aged haematopoietic stem cells. *Nat Commun*. 2016;7:11075-11087.
 70. Morrison SJ, Scadden DT. The bone marrow niche for haematopoietic stem cells. *Nature*. 2014;505(7483):327-334.
 71. Sánchez-Aguilera A, Méndez-Ferrer S. The hematopoietic stem-cell niche in health and leukemia. *Cell Mol Life Sci*. 2017;74(4):579-590.
 72. Walkley CR, Olsen GH, Dworkin S, et al. A microenvironment-induced myeloproliferative syndrome caused by retinoic acid receptor gamma deficiency. *Cell*. 2007;129(6):1097-1110.
 73. Walkley CR, Shea JM, Sims NA, Purton LE, Orkin SH. Rb regulates interactions between hematopoietic stem cells and their bone marrow microenvironment. *Cell*. 2007;129(6):1081-1095.
 74. Kim YW, Koo BK, Jeong HW, et al. Defective Notch activation in microenvironment leads to myeloproliferative disease. *Blood*. 2008;112(12):4628-4638.
 75. Wang L, Zhang H, Rodriguez S, et al. Notch-dependent repression of miR-155 in the bone marrow niche regulates hematopoiesis in an NF- κ B-dependent manner. *Cell Stem Cell*. 2014;15(1):51-65.
 76. Schepers K, Pietras EM, Reynaud D, et al. Myeloproliferative neoplasia remodels the endosteal bone marrow niche into a self-reinforcing leukemic niche. *Cell Stem Cell*. 2013;13(3):285-299.
 77. Arranz L, Sánchez-Aguilera A, Martín-Pérez D, et al. Neuroathy of haematopoietic stem cell niche is essential for myeloproliferative neoplasms. *Nature*. 2014;512(7512):78-81.
 78. Wen QJ, Yang Q, Goldenson B, et al. Targeting megakaryocytic-induced fibrosis in myeloproliferative neoplasms by AURKA inhibition. *Nat Med*. 2015;21(12):1473-1480.
 79. Pronk CJ, Veiby OP, Bryder D, Jacobsen SE. Tumor necrosis factor restricts hematopoietic stem cell activity in mice: involvement of two distinct receptors. *J Exp Med*. 2011;208(8):1563-1570.
 80. Larsson J, Karlsson S. The role of Smad signaling in hematopoiesis. *Oncogene*. 2005;24(37):5676-5692.
 81. Fleischman AG, Aichberger KJ, Luty SB, et al. TNF α facilitates clonal expansion of JAK2V617F positive cells in myeloproliferative neoplasms. *Blood*. 2011;118(24):6392-6398.
 82. Boveri E, Passamonti F, Rumi E, et al. Bone marrow microvessel density in chronic myeloproliferative disorders: a study of 115 patients with clinicopathological and molecular correlations. *Br J Haematol*. 2008;140(2):162-168.
 83. Gianelli U, Vener C, Raviele PR, et al. VEGF expression correlates with microvessel density in Philadelphia chromosome-negative chronic myeloproliferative disorders. *Am J Clin Pathol*. 2007;128(6):966-973.
 84. Trivai I, Ziegler M, Bergholz U, et al. Endogenous retrovirus induces leukemia in a xenograft mouse model for primary myelofibrosis. *Proc Natl Acad Sci USA*. 2014;111(23):8595-8600.
 85. Kagoya Y, Yoshimi A, Tsuruta-Kishino T, et al. JAK2V617F+ myeloproliferative neoplasm clones evoke paracrine DNA damage to adjacent normal cells through secretion of lipocalin-2. *Blood*. 2014;124(19):2996-3006.
 86. Lu M, Xia L, Liu YC, et al. Lipocalin produced by myelofibrosis cells affects the fate of both hematopoietic and marrow microenvironmental cells. *Blood*. 2015;126(8):972-982.
 87. Tehranchi R, Woll PS, Anderson K, et al. Persistent malignant stem cells in del(5q) myelodysplasia in remission. *N Engl J Med*. 2010;363(11):1025-1037.
 88. Gallipoli P, Abraham SA, Holyoake TL. Hurdles toward a cure for CML: the CML stem cell. *Hematol Oncol Clin North Am*. 2011;25(5):951-966, v.
 89. Holyoake T, Jiang X, Eaves C, Eaves A. Isolation of a highly quiescent subpopulation of primitive leukemic cells in chronic myeloid leukemia. *Blood*. 1999;94(6):2056-2064.
 90. Graham SM, Jørgensen HG, Allan E, et al. Primitive, quiescent, Philadelphia-positive stem cells from patients with chronic myeloid leukemia are insensitive to STI571 in vitro. *Blood*. 2002;99(1):319-325.
 91. Kröger NM, Deeg JH, Olavarria E, et al. Indication and management of allogeneic stem cell transplantation in primary myelofibrosis: a consensus process by an EBMT/ELN international working group. *Leukemia*. 2015;29(11):2126-2133.
 92. McLornan DP, Mead AJ, Jackson G, Harrison CN. Allogeneic stem cell transplantation for

- myelofibrosis in 2012. *Br J Haematol*. 2012; 157(4):413-425.
93. DeWitt MA, Magis W, Bray NL, et al. Selection-free genome editing of the sickle mutation in human adult hematopoietic stem/progenitor cells. *Sci Transl Med*. 2016;8(360):360ra134.
 94. Deininger M, Radich J, Burn TC, Huber R, Paranagama D, Verstovsek S. The effect of long-term ruxolitinib treatment on JAK2p.V617F allele burden in patients with myelofibrosis. *Blood*. 2015;126(13):1551-1554.
 95. Guglielmelli P, Rotunno G, Bogani C, et al; COMFORT-II Investigators. Ruxolitinib is an effective treatment for CALR-positive patients with myelofibrosis. *Br J Haematol*. 2016;173(6):938-940.
 96. Wang X, Ye F, Tripodi J, et al. JAK2 inhibitors do not affect stem cells present in the spleens of patients with myelofibrosis. *Blood*. 2014;124(19):2987-2995.
 97. Bhagwat N, Koppikar P, Keller M, et al. Improved targeting of JAK2 leads to increased therapeutic efficacy in myeloproliferative neoplasms. *Blood*. 2014;123(13):2075-2083.
 98. Meyer SC, Keller MD, Chiu S, et al. CHZ868, a type II JAK2 inhibitor, reverses type I JAK inhibitor persistence and demonstrates efficacy in myeloproliferative neoplasms. *Cancer Cell*. 2015;28(1):15-28.
 99. Park SO, Wamsley HL, Bae K, et al. Conditional deletion of Jak2 reveals an essential role in hematopoiesis throughout mouse ontogeny: implications for Jak2 inhibition in humans. *PLoS One*. 2013;8(3):e59675.
 100. Akada H, Akada S, Hutchison RE, Sakamoto K, Wagner KU, Mohi G. Critical role of Jak2 in the maintenance and function of adult hematopoietic stem cells. *Stem Cells*. 2014;32(7):1878-1889.
 101. Grisouard J, Hao-Shen H, Dirnhofer S, Wagner KU, Skoda RC. Selective deletion of Jak2 in adult mouse hematopoietic cells leads to lethal anemia and thrombocytopenia. *Haematologica*. 2014;99(4):e52-e54.
 102. Bandaranayake RM, Ungureanu D, Shan Y, Shaw DE, Silvennoinen O, Hubbard SR. Crystal structures of the JAK2 pseudokinase domain and the pathogenic mutant V617F. *Nat Struct Mol Biol*. 2012;19(8):754-759.
 103. Leroy E, Dusa A, Colau D, et al. Uncoupling JAK2 V617F activation from cytokine-induced signalling by modulation of JH2 α C helix. *Biochem J*. 2016;473(11):1579-1591.
 104. Silver RT. Recombinant interferon-alpha for treatment of polycythaemia vera. *Lancet*. 1988; 2(8607):403.
 105. Kiladjian JJ, Cassinat B, Chevret S, et al. Pegylated interferon-alfa-2a induces complete hematologic and molecular responses with low toxicity in polycythemia vera. *Blood*. 2008; 112(8):3065-3072.
 106. Quintás-Cardama A, Kantarjian H, Manshouri T, et al. Pegylated interferon alfa-2a yields high rates of hematologic and molecular response in patients with advanced essential thrombocythemia and polycythemia vera. *J Clin Oncol*. 2009;27(32):5418-5424.
 107. Kiladjian JJ, Mesa RA, Hoffman R. The renaissance of interferon therapy for the treatment of myeloid malignancies. *Blood*. 2011; 117(18):4706-4715.
 108. Verger E, Cassinat B, Chauveau A, et al. Clinical and molecular response to interferon- α therapy in essential thrombocythemia patients with CALR mutations. *Blood*. 2015;126(24): 2585-2591.
 109. Mullally A, Bruedigam C, Poveromo L, et al. Depletion of Jak2V617F myeloproliferative neoplasm-propagating stem cells by interferon- α in a murine model of polycythemia vera. *Blood*. 2013;121(18):3692-3702.
 110. Hasan S, Lacout C, Marty C, et al. JAK2V617F expression in mice amplifies early hematopoietic cells and gives them a competitive advantage that is hampered by IFN α . *Blood*. 2013;122(8): 1464-1477.
 111. King KY, Matatali KA, Shen CC, Goodell MA, Swierczek SI, Prchal JT. Comparative long-term effects of interferon alpha and hydroxyurea on human hematopoietic progenitor cells. *Exp Hematol*. 2015;43(10):912-918.e2.
 112. Larsen TS, Møller MB, de Stricker K, et al. Minimal residual disease and normalization of the bone marrow after long-term treatment with alpha-interferon2b in polycythemia vera. A report on molecular response patterns in seven patients in sustained complete hematological remission. *Hematology*. 2009;14(6):331-334.
 113. Ishii T, Xu M, Zhao Y, et al. Recurrence of clonal hematopoiesis after discontinuing pegylated recombinant interferon-alpha 2a in a patient with polycythemia vera. *Leukemia*. 2007;21(2): 373-374.
 114. Kuriakose ET, Gjoni S, Wang YL, et al. JAK2V617F allele burden is reduced by busulfan therapy: a new observation using an old drug. *Haematologica*. 2013;98(11):e135-e137.
 115. Lu M, Xia L, Li Y, Wang X, Hoffman R. The orally bioavailable MDM2 antagonist RG7112 and pegylated interferon α 2a target JAK2V617F-positive progenitor and stem cells. *Blood*. 2014; 124(5):771-779.
 116. Sánchez-Aguilera A, Arranz L, Martín-Pérez D, et al. Estrogen signaling selectively induces apoptosis of hematopoietic progenitors and myeloid neoplasms without harming steady-state hematopoiesis. *Cell Stem Cell*. 2014;15(6): 791-804.