

## LYMPHOID NEOPLASIA

Activation of the *LMO2* oncogene through a somatically acquired neomorphic promoter in T-cell acute lymphoblastic leukemia

Sunniyat Rahman,<sup>1</sup> Michael Magnussen,<sup>1</sup> Theresa E. León,<sup>1</sup> Nadine Farah,<sup>1</sup> Zhaodong Li,<sup>2</sup> Brian J. Abraham,<sup>3</sup> Krisztina Z. Alapi,<sup>1</sup> Rachel J. Mitchell,<sup>1</sup> Tom Naughton,<sup>1</sup> Adele K. Fielding,<sup>1</sup> Arnold Pizzey,<sup>1</sup> Sophia Bustraan,<sup>1</sup> Christopher Allen,<sup>1</sup> Teodora Popa,<sup>1</sup> Karin Pike-Overzet,<sup>4</sup> Laura Garcia-Perez,<sup>4</sup> Rosemary E. Gale,<sup>1</sup> David C. Linch,<sup>1</sup> Frank J. T. Staal,<sup>4</sup> Richard A. Young,<sup>3,5</sup> A. Thomas Look,<sup>2,6</sup> and Marc R. Mansour<sup>1</sup>

<sup>1</sup>Department of Haematology, University College London Cancer Institute, London, United Kingdom; <sup>2</sup>Department of Pediatric Oncology, Dana-Farber Cancer Institute, Harvard Medical School, Boston, MA; <sup>3</sup>Whitehead Institute for Biomedical Research, Cambridge, MA; <sup>4</sup>Department of Immunohematology, Leiden University Medical Center, Leiden, The Netherlands; <sup>5</sup>Department of Biology, Massachusetts Institute of Technology, Cambridge, MA; and <sup>6</sup>Division of Hematology/Oncology, Children's Hospital, Boston, MA

## Key Points

- Recurrent intronic mutations that create probable MYB, ETS1, and RUNX1 binding sites occur at the *LMO2* promoter in some T-ALL patients.
- CRISPR/Cas9-mediated disruption of the mutant MYB site in PF-382 cells markedly downregulates *LMO2* expression.

Somatic mutations within noncoding genomic regions that aberrantly activate oncogenes have remained poorly characterized. Here we describe recurrent activating intronic mutations of *LMO2*, a prominent oncogene in T-cell acute lymphoblastic leukemia (T-ALL). Heterozygous mutations were identified in PF-382 and DU.528 T-ALL cell lines in addition to 3.7% of pediatric (6 of 160) and 5.5% of adult (9 of 163) T-ALL patient samples. The majority of indels harbor putative de novo MYB, ETS1, or RUNX1 consensus binding sites. Analysis of 5'-capped RNA transcripts in mutant cell lines identified the usage of an intermediate promoter site, with consequential monoallelic *LMO2* overexpression. CRISPR/Cas9-mediated disruption of the mutant allele in PF-382 cells markedly downregulated *LMO2* expression, establishing clear causality between the mutation and oncogene dysregulation. Furthermore, the spectrum of CRISPR/Cas9-derived mutations provides important insights into the interconnected contributions of functional transcription factor binding. Finally, these mutations occur in the same intron as retroviral integration sites in gene therapy-induced T-ALL, suggesting that such events occur at preferential sites in the noncoding genome. (*Blood*. 2017;129(24):3221-3226)

## Introduction

LIM-domain-only protein 2 (*LMO2*) plays a crucial bridging role in the formation of a large multimeric transcriptional complex that includes TAL1, LDB1, GATA, RUNX1, ETS1, and MYB.<sup>1</sup> In mice, *Lmo2* is progressively silenced after the early T-cell progenitor (ETP) stage of thymic development and leads to T-cell acute lymphoblastic leukemia (T-ALL) when overexpressed in transgenic models.<sup>2-4</sup> In human thymi, *LMO2* is similarly downregulated after commitment to the T-cell lineage as indicated by DNA microarray analyses.<sup>5</sup> Overexpression of *LMO2* in human hematopoietic stem cells also leads exclusively to preleukemic alterations in thymocytes and T cells but not in other lineages.<sup>6</sup> Reported mechanisms of aberrant *LMO2* expression in human T-ALL include recurrent chromosomal translocations, such as t(11;14)(p13;q11) and t(7;11)(q35;p13); cryptic deletions of an upstream negative regulatory region, as in del(11)(p12p13); and retroviral insertional mutagenesis at the *LMO2* locus during gene therapy.<sup>7-11</sup> Although ~50% of T-ALL patients overexpress *LMO2*, only about 10% of patients have a detectable cytogenetic lesion.<sup>12</sup> Notably, many of these patients will overexpress *LMO2* from a single allele, a feature reminiscent of *TAL1* overexpressing T-ALL cases driven by small somatic indel mutations that create binding

sites for MYB, thus generating a neomorphic enhancer.<sup>13,14</sup> We hypothesized that *cis*-acting mechanisms may account for T-ALL cases with monoallelic *LMO2* expression that lack abnormalities of the *LMO2* locus.<sup>15,16</sup>

## Study design

Detailed methods are described in the supplemental Data available on the *Blood* Web site. Chromatin immunoprecipitation sequencing (ChIP-seq) was performed on T-ALL cell lines after immunoprecipitation with antibodies against MYB and acetylated H3K27 (H3K27ac). Analysis of motif enrichment was used to confirm enrichment of MYB motifs in the MYB ChIP-seq data (supplemental Tables 1 and 2). *LMO2* messenger RNA levels were quantified by quantitative reverse transcription polymerase chain reaction (qRT-PCR). Mutation screening of primary T-ALL samples was achieved by denaturing high-performance liquid chromatography of *LMO2* intron 1 PCR products. Luciferase reporter constructs consisting of 469 bp PCR products inserted upstream of an SV40 promoter and firefly luciferase gene were electroporated into Jurkat cells. CRISPR/Cas9 genome editing was used to target the *LMO2* intron 1 mutations in the PF-382 T-ALL cell line.

Submitted 27 September 2016; accepted 22 February 2017. Prepublished online as *Blood* First Edition paper, 7 March 2017; DOI 10.1182/blood-2016-09-742148.

The online version of this article contains a data supplement.

There is an Inside *Blood* Commentary on this article in this issue.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2017 by The American Society of Hematology




Figure 1.

## Results and discussion

To test this hypothesis, we first assessed *LMO2* expression by qRT-PCR in several T-ALL cell lines arrested at different stages of thymic differentiation. The ETP-like T-ALL cell line Loucy expressed *LMO2* at levels significantly higher than the more mature T-ALL cell lines (DND-41, ALL-SIL, Jurkat), reflecting physiological expression of *LMO2* at the ETP stage of thymic development (Figure 1A). The *TALI*-positive cell lines DU.528 and PF-382 both exhibited upregulated *LMO2* expression, yet crucially have no reported chromosomal lesions affecting this locus (Figure 1A).<sup>17,18</sup> In contrast to Loucy cells, aberrant H3K27ac marks (indicative of active chromatin) were identified before and encompassing the noncoding exon 2 of the *LMO2* gene by ChIP-seq in PF-382 and DU.528 T-ALL cell lines (Figure 1B; supplemental Figure 1). Sequencing across these peaks revealed a heterozygous 20 bp duplication in PF-382 cells and a heterozygous 1 bp deletion in DU.528 cells located close to a region recently described as an intermediate promoter for reasons that were not then apparent (Figure 1B).<sup>19</sup> Notably, the mutations were not described as normal germ line variants in the Single Nucleotide Polymorphism Database. In silico analysis of the reference sequence identified a high-confidence primary MYB binding motif (AACCGTT) that was duplicated in the PF-382 cell line, whereas the single bp deletion in DU.528 cells creates a CAACCGC sequence that closely resembles a secondary MYB binding motif (Figure 1B; supplemental Tables 3 and 4).

To assess whether the mutations form aberrant sites of MYB binding, we performed ChIP-seq for MYB and analyzed peaks of MYB enrichment at the *LMO2* locus. There was a complete absence of MYB binding at the intermediate promoter in cells that were wild-type at this locus, suggesting that the presence of the single native MYB motif in itself is insufficient to recruit MYB. In contrast, both PF-382 and DU.528 cells that harbor dual MYB motifs displayed precisely aligned MYB binding at the mutation site (Figure 1B). To determine whether the mutations affected promoter usage, we performed rapid amplification of 5' complementary DNA ends in *LMO2* mutant and wild-type cell lines by using a common primer in exon 6 capable of capturing the transcription start site of all *LMO2* isoforms. Whereas the majority (73%) of 5' capped transcripts in Loucy cells originated from the proximal promoter, both PF-382 and DU.528 cells demonstrated preferential usage of the recently described intermediate promoter (75% and 67% of transcripts, respectively; Figure 1C).


Our observations were not limited to T-ALL cell lines because heterozygous mutations at *LMO2* intron 1 were detected in diagnostic samples from 3.7% of pediatric (6 of 160) and 5.5% of adult (9 of 163) T-ALL patients (Figure 1D). Absence of the mutations in 7 available patient-matched remission samples confirmed that they were somatic (supplemental Figure 2). Notably, the mutations were densely distributed around highly conserved native ETS1, MYB, and GATA motifs (supplemental Figure 3). Including the cell lines, 7 mutations

introduced an additional MYB site, resulting in 2 MYB motifs spaced 10 or 20 bp apart, equivalent to 1 or 2 helical coils of DNA, respectively (Figure 1E). Three mutations created potential binding sites for both MYB and ETS1, 3 formed potential ETS1 sites, and 3 produced potential new RUNX1 binding sites (Figure 1E; supplemental Tables 3 and 4). Given that *NOTCH* and *TALI* have been shown to collaborate with *LMO2* to promote leukemogenesis in murine models of T-ALL, it is noteworthy that of the 15 patients with *LMO2* promoter mutations, 7 had *NOTCH-1* mutations and 8 had *TALI* activating lesions, including 2 with *TALI*-enhancer mutations (both creating new MYB motifs; supplemental Table 5).<sup>20,21</sup> Such collaboration between *TALI*, *LMO2*, and *NOTCH-1* has also been described in gene therapy-induced T-ALL, including 1 patient who harbored both a retroviral integration upstream of *LMO2* and an episomal reintegration at the *TALI* locus.<sup>9,13,22</sup>

To ascertain whether *LMO2* promoter mutations in T-ALL led to aberrant expression compared with its matched thymic counterpart, we assessed *LMO2* expression by qRT-PCR in thymic subsets sorted for different levels of thymic differentiation.<sup>5</sup> Validating earlier reports that used microarrays, *LMO2* expression was highest in the most immature, precommitment stages of T-cell development and was expressed at low levels from the double-negative stage onward, when thymocytes had undergone biallelic TCR- $\gamma$  rearrangement (Figure 2A).<sup>5</sup> To determine the level of differentiation arrest of the 15 mutant patient samples, we analyzed the TCR- $\gamma$  locus by qPCR (supplemental Figure 4); 12 of the 15 samples (including 5 of the 6 patients with available RNA) had biallelic TCR- $\gamma$  deletion (position weight matrices; supplemental Table 5), indicating that maturation arrest occurred after the pro-T-cell stage of differentiation, and that the majority of patients did not have the ETP ALL phenotype. Thus, compared with their physiological counterparts, those patients with RNA available for *LMO2* qRT-PCR exhibited aberrant *LMO2* overexpression ( $P < .002$  vs double-negative and double-positive subsets; Figure 2A). Although we were unable to confirm *LMO2* overexpression in all mutant samples because of the unavailability of RNA, all classes of mutation (additional MYB, ETS1, RUNX1, or MYB+ETS1 sites) were represented in the 6 patients with *LMO2* overexpression. Exploiting a heterozygous germ line single nucleotide polymorphism (rs3740617), DU.528 cells and 3 of 4 informative patient samples displayed skewed allelic expression of *LMO2* (Figure 2B). The observation of biallelic expression in sample A1 suggests a potential lesion on the second allele that remains undefined. Consistent with their *cis*-activating potential,  $\geq 96\%$  of reads from MYB ChIP-seq performed in DU.528 and PF-382 cells aligned to the mutant rather than the wild-type allele (Figure 2C; supplemental Figure 5). Furthermore, the gain-of-function nature of the mutations was confirmed by luciferase reporter assays conducted in Jurkat cells where all mutations markedly activated luciferase activity compared with the wild-type sequence (Figure 2D; supplemental Figure 6A).

To assess causality between the mutations and *LMO2* dysregulation, we used CRISPR/Cas9 genome editing with a guide

**Figure 1. *LMO2* intron 1 mutations in pediatric and adult human T-ALL.** (A) *LMO2* expression as determined by qRT-PCR in *LMO2* translocated T-ALL cell lines (KOPT-K1 and P12-Ichikawa) and nontranslocated T-ALL cell lines (DU.528, PF-382, Loucy, DND41, Jurkat, and ALL-SIL). (B) ChIP-Seq tracks at the *LMO2* locus for MYB and H3K27ac in PF-382, DU.528, Loucy, and Jurkat T-ALL cell lines. Y-axis values are reads per bin per million mapped reads. Mutations are shown below as identified by Sanger sequencing of PF-382 and DU.528 DNA, with inserted sequences shown in red and MYB motifs underlined. The position weight matrices for the primary and secondary MYB binding sites are from UniPROBE.<sup>27</sup> (C) Pie chart summarizing the percentage of *LMO2* transcripts identified by rapid amplification of 5' complementary DNA ends that start from the distal, intermediate, and proximal promoters for the PF-382, DU.528, and Loucy T-ALL cell lines. A total of 20, 21, and 22 *LMO2* transcripts was examined for PF-382, DU.528, and Loucy T-ALL cell lines, respectively. (D) Pie chart summarizing mutation recurrence within pediatric and adult human T-ALL cohorts. (E) Indels mapped to the *LMO2* intron 1 mutation hotspot labeled with the associated de novo consensus site as aligned to the UniPROBE or HOCOMOCO position weight matrices, in which MYB, ETS1, and RUNX1 sites are marked as a triangle, square, and diamond, respectively. Below, motif analysis of the region shows the native binding sites for members of the TAL1 complex, including RUNX1, E-box (for TAL1 binding), ETS1, MYB, and GATA. TSS, transcription start site.



**Figure 2. *LMO2* intron 1 indels are predominantly monoallelically activating, and CRISPR/Cas9-mediated knockout of the PF-382 mutant allele downregulates *LMO2* expression.** (A) *LMO2* expression as determined by qRT-PCR in human sorted thymic subsets, primary patient samples with *LMO2* intron 1 indels, and the wild-type Jurkat cell line.  $P < .002$  for samples A1, A2, A3, A9, and P6 vs double negative and double positive by two-tailed  $t$  test. Primary patient samples were assessed for the absence of biallelic TCR- $\gamma$  deletion (ABD), of which patient sample A4 (orange bar) exhibited ABD, whereas all other patients were non-ABD. (B) The informative single nucleotide polymorphism (SNP) (rs3740617) was amplified in 4 patient samples and the DU.528 cell line from both genomic DNA (gDNA) and complementary DNA (cDNA) templates to infer monoallelic expression. To do this, if one chromatogram peak is detected at a heterozygous SNP within the cDNA, the expression can be interpreted as coming from one allele. (C) Quantification of the number of reads mapped to the wild-type (WT) or mutant (MUT) allele where 54 of 56 reads and 85 of 85 reads mapped to the mutant alleles for DU.528 and PF-382, respectively. (D) Firefly luciferase activity after renilla and no-insert vector normalization for patient-derived indels. Data shown are from  $\geq 3$  independent experiments performed in triplicate. Values are mean  $\pm$  standard deviation and  $P$  values were calculated by a two-tailed Student  $t$  test. (E) The yellow highlighted sequence is the target region for the CRISPR/Cas9 guide RNA. Aligned sequences are from CRISPR/Cas9-edited PF-382 single-cell clones showing the associated genomic edits generated. Red sequences are inserted sequences, blue are altered, and dashes represent deleted bases. Underlined region shows the presence of the native and mutant MYB binding sites. (F) Gene expression of *LMO2* for each PF-382 clone, as determined by qRT-PCR. Data are expressed as fold change relative to the mean expression of the unedited clones in arbitrary units (AU). Clones are labeled as "unedited" when CRISPR/Cas9 did not edit the region targeted by the guide RNA and "edited" when successful targeting led to the formation of an indel. \* $P \leq .05$ , \*\* $P \leq .01$ , \*\*\* $P \leq .001$ , and \*\*\*\* $P \leq .0001$ . NOI, no-insert control; T/C, T to C single nucleotide polymorphism.

RNA designed to target the duplicated MYB site in PF-382 cells (supplemental Figure 6B). Crucially, clone 4F11, which had a single T>C substitution disrupting the MYB binding site, and clone 1A8, in which the mutant allele had been reverted to wild-type, resulted in the most dramatic downregulation of *LMO2* (Figure 2E-F; supplemental Figure 7). Interestingly, 2 clones (4H12 and 6D4) that increased the distance between the native and the mutant MYB sites resulted in a marked reduction in *LMO2* expression, supporting the hypothesis that MYB binding is augmented when additional motifs are orientated on the same side

of the DNA helix.<sup>23</sup> This was further validated by the lack of reduction in *LMO2* expression in a clone (5F10) in which the sequence between the 2 MYB sites was altered but the spacing distance was unchanged.

In conclusion, we identified and functionally validated a novel recurrent mutation hotspot occurring in a noncoding site that drives *LMO2* overexpression from a neomorphic promoter in a substantial proportion of both adult and pediatric T-ALL patients. Remarkably, the mutations create potential binding sites for MYB, ETS1, or RUNX1, all of which are members of a highly oncogenic TAL1-*LMO2* complex in




Figure 2. (Continued).

T-ALL; this indicates that *LMO2* is a component of an autoregulatory self-sustaining positive feedback loop in these cells, analogous to the autoregulation of *TALI* we recently described in Jurkat cells.<sup>14,24</sup> To prove that the newly formed ETS1 and RUNX1 sites are sufficient to drive *LMO2* expression, we attempted to but ultimately were unable to knock in these mutations in vitro. Thus, the oncogenic potential of these particular mutations remain an area of ongoing study. There is still a question regarding exactly how various members of the TAL1 complex position themselves on DNA with regard to spacing, orientation, and order of motifs, so-called syntax.<sup>25</sup> Thus, identifying gain-of-function noncoding mutations that have been selected for during tumorigenesis in vivo offers important insights into the optimal DNA syntax required for nucleation of such multiprotein transcription factor complexes. For instance, it may become apparent why a single MYB binding site is sufficient to drive expression from certain loci, such as at the *TALI* enhancer, whereas others require dual MYB motifs. Finally, we note that these mutations occur within the same intron as retroviral integration sites described in 2 cases of gene therapy-induced T-ALL (supplemental Figure 8).<sup>22,26</sup> This raises the possibility that formation of aberrant promoters and enhancers, either by mutation or retroviral insertion, occur at preferred rather than random sites in the noncoding genome.

### Acknowledgments

The authors thank the patients, families, and clinical teams who have been involved in both trials. Primary childhood leukemia samples used in this study were provided by the Bloodwise Childhood Leukemia Cell Bank, working with the laboratory teams in the Bristol Genetics Laboratory, Southmead Hospital, Bristol, United Kingdom; Molecular Biology Laboratory, Royal Hospital for Sick Children, Glasgow, United Kingdom; Molecular Haematology Laboratory,

Royal London Hospital, London, United Kingdom; and Molecular Genetics Service and Sheffield Children’s Hospital, Sheffield, United Kingdom.

This work was supported by National Institutes of Health, National Cancer Institute grants 1R01CA176746-01, 5P01CA109901-08, and 5P01CA68484 (A.T.L.), by Bloodwise and Gabrielle’s Angels Foundation (M.R.M. and S.R.), by the Freemason’s Grand Charity (M.M.), by Alex’s Lemonade Stand Foundation for Childhood Cancer (Z.L.), by grants from Bloodwise (formerly known as Leukaemia and Lymphoma Research, UK) and the Medical Research Council (UK) for the UKALL2003 trial (ISRCTN07355119), and by Cancer Research UK for the UKALL14 trial (ISRCTN66541317). This work was undertaken at University College London, which receives funding from the Department of Health’s National Institute of Health Research Biomedical Research Centre. B.J.A. is a Hope Funds for Cancer Research Grillo-Marxuach Family Fellow.

### Authorship

Contribution: S.R., M.M., T.E.L., N.F., A.P., Z.L., S.B., C.A., T.P., K.P.-O., L.G.-P., and B.J.A. performed experimental work; K.Z.A., R.J.M., T.N., A.K.F., R.E.G., K.P.-O., L.G.-P., and F.J.T.S. provided primary samples; S.R., M.M., T.E.L., N.F., D.C.L., R.A.Y., F.J.T.S., and A.T.L. analyzed data; S.R., R.E.G., F.J.T.S., D.C.L., and M.R.M. wrote the manuscript; M.R.M. designed the study; and all authors approved the final manuscript.

Conflict-of-interest disclosure: R.A.Y. is a founder and member of the Board of Directors of Syros Pharmaceuticals, which develops therapies that target gene regulatory elements. The remaining authors declare no competing financial interests.

Correspondence: Marc R. Mansour, University College London Cancer Institute, Department of Haematology, 72 Huntley St, London WC1E 6BT, United Kingdom; e-mail: m.mansour@ucl.ac.uk.

## References

- Chambers J, Rabbitts TH. LMO2 at 25 years: a paradigm of chromosomal translocation proteins. *Open Biol*. 2015;5(6):150062.
- Boehm T, Foroni L, Kaneko Y, Perutz MF, Rabbitts TH. The rhombotin family of cysteine-rich LIM-domain oncogenes: distinct members are involved in T-cell translocations to human chromosomes 11p15 and 11p13. *Proc Natl Acad Sci USA*. 1991;88(10):4367-4371.
- Fisch P, Boehm T, Lavenir I, et al. T-cell acute lymphoblastic lymphoma induced in transgenic mice by the RBTN1 and RBTN2 LIM-domain genes. *Oncogene*. 1992;7(12):2389-2397.
- Herblot S, Steff AM, Hugo P, Aplan PD, Hoang T. SCL and LMO1 alter thymocyte differentiation: inhibition of E2A-HEB function and pre-T alpha chain expression. *Nat Immunol*. 2000;1(2):138-144.
- Dik WA, Pike-Overzet K, Weerkamp F, et al. New insights on human T cell development by quantitative T cell receptor gene rearrangement studies and gene expression profiling. *J Exp Med*. 2005;201(11):1715-1723.
- Wiekmeijer AS, Pike-Overzet K, Brugman MH, et al. Overexpression of LMO2 causes aberrant human T-Cell development in vivo by three potentially distinct cellular mechanisms. *Exp Hematol*. 2016;44(9):838-849.e9.
- Hacein-Bey-Abina S, von Kalle C, Schmidt M, et al. A serious adverse event after successful gene therapy for X-linked severe combined immunodeficiency. *N Engl J Med*. 2003;348(3):255-256.
- Hacein-Bey-Abina S, Von Kalle C, Schmidt M, et al. LMO2-associated clonal T cell proliferation in two patients after gene therapy for SCID-X1. *Science*. 2003;302(5644):415-419.
- Howe SJ, Mansour MR, Schwarzwaelder K, et al. Insertional mutagenesis combined with acquired somatic mutations causes leukemogenesis following gene therapy of SCID-X1 patients. *J Clin Invest*. 2008;118(9):3143-3150.
- Van Vlierberghe P, Ferrando A. The molecular basis of T cell acute lymphoblastic leukemia. *J Clin Invest*. 2012;122(10):3398-3406.
- Van Vlierberghe P, van Grotel M, Beverloo HB, et al. The cryptic chromosomal deletion del(11)(p12p13) as a new activation mechanism of LMO2 in pediatric T-cell acute lymphoblastic leukemia. *Blood*. 2006;108(10):3520-3529.
- Ferrando AA, Look AT. Gene expression profiling in T-cell acute lymphoblastic leukemia. *Semin Hematol*. 2003;40(4):274-280.
- Navarro JM, Touzart A, Pradel LC, et al. Site- and allele-specific polycomb dysregulation in T-cell leukaemia. *Nat Commun*. 2015;6:6094.
- Mansour MR, Abraham BJ, Anders L, et al. Oncogene regulation. An oncogenic super-enhancer formed through somatic mutation of a noncoding intergenic element. *Science*. 2014;346(6215):1373-1377.
- Ferrando AA, Herblot S, Palomero T, et al. Biallelic transcriptional activation of oncogenic transcription factors in T-cell acute lymphoblastic leukemia. *Blood*. 2004;103(5):1909-1911.
- Van Vlierberghe P, Beverloo HB, Buijs-Gladdines J, et al. Monoallelic or biallelic LMO2 expression in relation to the LMO2 rearrangement status in pediatric T-cell acute lymphoblastic leukemia. *Leukemia*. 2008;22(7):1434-1437.
- Chen S, Nagel S, Schneider B, et al. Novel non-TCR chromosome translocations t(3;11)(q25;p13) and t(X;11)(q25;p13) activating LMO2 by juxtaposition with MBNL1 and STAG2. *Leukemia*. 2011;25(10):1632-1635.
- Dong WF, Xu Y, Hu QL, et al. Molecular characterization of a chromosome translocation breakpoint t(11;14)(p13;q11) from the cell line KOPT-K1. *Leukemia*. 1995;9(11):1812-1817.
- Oram SH, Thoms JAI, Pridans C, et al. A previously unrecognized promoter of LMO2 forms part of a transcriptional regulatory circuit mediating LMO2 expression in a subset of T-acute lymphoblastic leukaemia patients. *Oncogene*. 2010;29(43):5796-5808.
- Larson RC, Lavenir I, Larson TA, et al. Protein dimerization between Lmo2 (Rbtn2) and Tal1 alters thymocyte development and potentiates T cell tumorigenesis in transgenic mice. *EMBO J*. 1996;15(5):1021-1027.
- O'Neil J, Calvo J, McKenna K, et al. Activating Notch1 mutations in mouse models of T-ALL. *Blood*. 2006;107(2):781-785.
- Hacein-Bey-Abina S, Garrigue A, Wang GP, et al. Insertional oncogenesis in 4 patients after retrovirus-mediated gene therapy of SCID-X1. *J Clin Invest*. 2008;118(9):3132-3142.
- Molvaersmyr AK, Saether T, Gilfillan S, et al. A SUMO-regulated activation function controls synergy of c-Myb through a repressor-activator switch leading to differential p300 recruitment. *Nucleic Acids Res*. 2010;38(15):4970-4984.
- Sanda T, Lawton LN, Barrasa MI, et al. Core transcriptional regulatory circuit controlled by the TAL1 complex in human T cell acute lymphoblastic leukemia. *Cancer Cell*. 2012;22(2):209-221.
- Farley EK, Olson KM, Zhang W, Rokhsar DS, Levine MS. Syntax compensates for poor binding sites to encode tissue specificity of developmental enhancers. *Proc Natl Acad Sci USA*. 2016;113(23):6508-6513.
- Braun CJ, Boztug K, Paruzynski A, et al. Gene therapy for Wiskott-Aldrich syndrome—long-term efficacy and genotoxicity. *Sci Transl Med*. 2014;6(227):227ra33.
- Hume MA, Barrera LA, Gisselbrecht SS, Butyk ML. UniPROBE, update 2015: new tools and content for the online database of protein-binding microarray data on protein-DNA interactions. *Nucleic Acids Res*. 2015;43(Database issue):D117-D122.