



TO THE EDITOR:

Treatment with 5-azacytidine induces a sustained response in patients with angioimmunoblastic T-cell lymphoma

François Lemonnier,¹⁻³ Jehan Dupuis,¹ Pierre Sujobert,^{4,5} Olivier Toumillhac,⁶ Morgane Cheminant,⁷⁻⁹ Clémentine Sarkozy,^{5,10} Laura Pelletier,³ Ambroise Marçais,⁷⁻⁹ Cyrielle Robe,³ Virginie Fataccioli,^{3,11} Corinne Haioun,¹⁻³ Olivier Hermine,⁷⁻⁹ Philippe Gaulard,^{2,3,11,*} and Richard Delarue^{7-9,*}

¹Unité Hémopathies Lymphoïdes, Hôpitaux Universitaires Henri Mondor, Assistance Publique des Hôpitaux de Paris, Créteil, France; ²Université Paris-Est Créteil, Créteil, France; ³Institut Mondor de Recherche Biomédicale, INSERM U955, Créteil, France; ⁴Hospices Civils de Lyon, Service d'Hématologie Biologique, Pierre Bénite, France; ⁵INSERM 1052, CNRS 5286, Université Claude Bernard, Faculté de Médecine Lyon-Sud Charles Mérieux, Université de Lyon, Pierre Bénite, France; ⁶Service d'Hématologie Clinique Adulte et de Thérapie Cellulaire, CHU Clermont-Ferrand, Clermont-Ferrand, France; ⁷Service d'Hématologie Adultes, Hôpital Necker, Assistance Publique des Hôpitaux de Paris, Paris, France; ⁸Université Paris Descartes, Paris, France; ⁹INSERM UMR 1163, CNRS ERL 8254, Laboratory of Cellular and Molecular Mechanisms of Hematological Disorders and Therapeutical Implications, Imagine Institute, Paris, France; ¹⁰Service d'Hématologie Clinique, Hospices Civils de Lyon, Hôpital Lyon Sud, Pierre Bénite, France; and ¹¹Département de Pathologie, Hôpitaux Universitaires Henri Mondor, Assistance Publique des Hôpitaux de Paris, Créteil, France

Peripheral T-cell lymphomas (PTCLs) are heterogeneous diseases resulting from the malignant transformation of mature T or natural killer cells. Their epidemiology varies widely, but PTCLs that derive from T follicular helper (TFH) cells, which include angioimmunoblastic T-cell lymphoma (AITL), follicular PTCL, and other nodal PTCLs with a TFH phenotype,¹ appear to be frequent among PTCLs.^{2,3} Patients have a poor prognosis, especially after relapse, with a median overall survival (OS) of ~6 months.⁴ The US Food and Drug Administration has approved pralatrexate, romidepsin, and belinostat for relapsing/refractory PTCL, but these 3 drugs still show limited activity in PTCL, with an overall response rate of ~30%.⁵⁻⁷ Thus, PTCL therapy is still an unmet medical need.

Recurrent mutations in *TET2*,⁸⁻¹⁰ *DNMT3A*,⁹⁻¹¹ and/or *IDH2*^{9,10,12} are detected in ~80%, 25%, and 25% of patients with TFH-derived PTCL, respectively. These 3 genes directly or indirectly regulate cytosine methylation and hydroxymethylation, and mutations of these genes result in changes in DNA methylation levels.¹³ *TET2*, *DNMT3A*, and *IDH1/2* are also mutated in myeloid neoplasms, especially acute myeloid leukemia and myelodysplastic syndromes.¹³ Treatment with the hypomethylating agents (HMAs) 5-azacytidine and decitabine shows efficacy in these diseases, and the response rate to HMAs appears to correlate with *TET2*, *IDH1/2*, and/or *DNMT3A* mutations.¹⁴⁻¹⁶ This suggests that HMAs could have activity against TFH-derived PTCL. We previously reported 2 patients with AITL and chronic myelomonocytic leukemia (CMML) who experienced sustained complete remission of both diseases after treatment with 5-azacytidine.^{17,18} Here, we expand these observations and describe the response and outcome of a retrospective series of 12 AITL patients who received 5-azacytidine for concomitant myeloid neoplasm or used as compassionate therapy in relapsing/refractory AITL patients in the absence of available therapy or when such therapy was contraindicated, at the discretion of the physician.

Patients received 5-azacytidine 75 mg/m² daily, subcutaneously, for 7 consecutive days, every 28 days, until progression or unacceptable toxicity. Evaluations were performed by computed

tomography scan, and responses were assessed by investigators following the 1999 Cheson criteria.¹⁹

Two patients (PTCL1¹⁷ and PTCL7¹⁸) were previously reported. AITL diagnoses were all confirmed by expert pathologists in the framework of the national program "lymphopath,"³ based on the criteria of the World Health Organization 2008 classification, and tumor samples were collected in the frame of the Tenomic network from the Lymphoma Study Association. *TET2*, *IDH2*, *DNMT3A*, and *RHOA* status was centrally determined by targeted deep sequencing of DNA extracted from formalin-fixed, paraffin-embedded samples using PGM technology (ThermoFisher) (see supplemental Methods, available at the *Blood* Web site). The median depth of sequencing was 1432×.

Patient characteristics are given in Table 1. The median age was 70.5 years (interquartile range [IQR], 67.5-73.5 years). All but one had relapsed-refractory disease, with a median of 2 lines of therapy (IQR, 1.75-3), including cyclophosphamide, doxorubicin, vincristine, and prednisone (CHOP)-like therapy in 10 cases and mini-CHOP in one. PTCL1 was the only treatment-naïve patient. She received 5-azacytidine as first-line therapy for concomitant CMML with excess blasts and thrombocytopenia. Three other patients had asymptomatic CMML, and 1 patient had refractory cytopenia with multilineage dysplasia. In total, 5 out of 12 patients (41%) had an associated myeloid neoplasm. All patients had disseminated (stage III-IV) disease, and 4 out of 12 patients (33%) had a poor performance status (Eastern Cooperative Oncology Group [ECOG] >2) at 5-azacytidine initiation.

Treatment was introduced between January 2013 and July 2016. Patients received a median of 5.5 cycles (IQR, 3.75-17 cycles). In addition to 5-azacytidine, 6 out of 12 patients (50%) received rituximab because of the presence of Epstein-Barr virus replication or numerous Epstein-Barr virus B-blasts in the lymph node biopsy.

Treatment was well tolerated. Three patients required transfusion, and none developed febrile neutropenia. One patient experienced grade 2 neuropathy that was considered to be

Table 1. Patient characteristics and treatment

ID	Sex	Age at diagnosis (y)	Associated myeloid disorder	Mutations (VAF%)	IPI at diagnosis	Number of previous therapies	Previous auto-HSCT	Stage	LDH	ECOG	Rituximab (no. of cycles)	5-azacytidine (no. of cycles)	Best response
PTCL1	F	79	CMML	TET2 p.Q891fs (34.6)	4	0	0	3	1	2	4	61	CR
				TET2 c.3955-2A>G (38.4)									
PTCL2	M	69	MDS	TET2 p.L249fs (30)	3	1	0	4	0	2	4	25	CR
PTCL3	F	70	No	TET2 p.C1271W (82)	2	6	0	3	0	1	0	7	CR
				RHOA p.K18N (11.4)									
PTCL4	M	75	CMML	TET2 p.E1879A (23)	2	4	0	3	1	3	0	6	CR
PTCL5	M	63	No	TET2 p.H1904R (42.2)	4	2	1	4	1	1	6	5	CR
				TET2 p.E1207K (44)									
				DNMT3A p.G707fs (40.20%)									
PTCL6	F	73	No	TET2 p.R1465X (28.3)	4	2	0	4	1	4	8	16	PR
				DNMT3A p.R882H (30.6)									
				RHOA p.G17V (24.3)									
PTCL7	F	85	CMML	TET2 p.S835X (27.6)	3	1	0	4	0	4	0	20	CR
				TET2 c.3594+1G>C (24.7)									
				DNMT3A p.R882H (27)									
				RHOA p.G17V (2.8)									
PTCL8	M	39	No	TET2 p.T938fs (30.3)	4	2	0	4	1	1	6	4	SD
				RHOA p.G17V (29.10)									
PTCL9	F	81	CMML	TET2 p.R1404X (30)	3	3	0	4	1	1	0	4	PR
				TET2 p.C1298Y (31)									
				RHOA p.G17V (12.10)									
PTCL10	M	51	No	TET2 p.R1216X (11.3)	2	3	1	3	1	1	0	3	PR
				TET2 p.C1271W (8.9)									
				IDH2 p.R172K (2.2)									

F, female; HSCT, hematopoietic stem cell transplant; IPI, International Prognostic Index; M, male; MDS, myelodysplastic syndrome; SD, stable disease; VAF, variant allele frequency.

Table 1. (continued)

ID	Sex	Age at diagnosis (y)	Associated myeloid disorder	Mutations (VAF%)	IPI at diagnosis	Number of previous therapies	Previous auto-HSCT	Stage	LDH	ECOG	Rituximab (no. of cycles)	5-azacytidine (no. of cycles)	Best response
PTCL11	M	71	No	TET2 p.H1904L (24.8)	5	2	0	4	1	2	6	2	SD
				DNMT3A p.R882H (45.3)									
PTCL12	M	69	No	TET2 p.L1244fs (28.3)	5	2	0	4	1	4	0	2	SD
				TET2 p.H937fs (10.2)									

F, female; HSCT, hematopoietic stem cell transplant; IPI, International Prognostic Index; M, male; MDS, myelodysplastic syndrome; SD, stable disease; VAF, variant allele frequency.

paraneoplastic syndrome unrelated to treatment, and another experienced unexpected digestive toxicity (grade 3 diarrhea). There were no treatment-related deaths.

Nine patients showed a response, including 6 complete responses (CRs) and 3 partial responses (PRs), leading to an overall response rate of 75%, including 50% CRs and 25% PRs (Figure 1A). After a median follow-up of 27 months, the median progression-free survival was 15 months and median OS of 21 months (Figure 1 B-C). No patient died of myeloid neoplasm during the follow-up period. It is noteworthy that some elderly patients with poor performance status (ECOG 3-4) had a sustained AITL response after 5-azacytidine treatment, with an acceptable tolerance.

Three patients stopped treatment after 5, 7, and 6 cycles: 1 because of digestive toxicity, 1 at his request, and 1 at the request of the physician. The first 2 patients remained in complete remission >18 months after discontinuing 5-azacytidine, whereas the last (PTCL4) relapsed 9 months after. Treatment was resumed, rapidly resulting in a sustained CR. He is currently still on therapy.

Five patients have shown a sustained response, as they are still in complete remission >23 months after treatment initiation. Three are still receiving treatment, whereas it was discontinued in the other 2 patients, raising the question of the optimal duration of HMA therapy and the possibility of discontinuing treatment.

Molecular studies were centrally performed using targeted deep sequencing. We detected *TET2* mutations in 12 out of 12 patients, and 7 out of 12 patients (58%) had 2 mutations. In addition, 4 out of 12 patients (33%) had *DNMT3A* mutations, 5 out of 12 patients (41%) had *RHOA* mutations, 4 out of 5 patients had a p.G17V substitution, and only 1 patient had an *IDH2*^{R172} mutation (supplemental Table 1). We were unable to assess the impact of *TET2* mutation on treatment response because the samples from all patients were *TET2* mutated. Within the limit of this cohort, we observed no relationship among number of *TET2* mutations; mutations in *DNMT3A*, *IDH2*, or *RHOA*; and response to treatment. An association with rituximab was not associated with a higher response rate (66% vs 83%; $P = 1$), likely reflecting the limited efficacy of this drug in AITL.²⁰ Finally, the 5 patients with an associated myeloid neoplasm responded (4 CRs and 1 PR), while 4 out of 7 patients without associated myeloid anomalies responded (2 CRs and 2 PRs; $P = .2$), indicating that the effect of 5-azacytidine on AITL is not restricted to patients with associated myeloid disease.

We highlight here an association of AITL with CMML in 4 patients. This association has rarely been reported^{17,18,21} but could be more frequent than previously thought. Indeed, these 2 neoplasms share common oncogenic events, such as *TET2* or *DNMT3A* mutations, which usually occur in hematopoietic progenitor cells and could lead to the development of both diseases. Clinical and molecular studies are required for a better understanding of this association. Furthermore, the mechanism of action of 5-azacytidine in AITL has not been elucidated yet. Indeed, it is unknown whether 5-azacytidine has a direct effect on neoplastic T cells or whether it acts on abnormal *TET2*-mutated myeloid cells. Indeed, it could be hypothesized that abnormal *TET2*-mutated myeloid cells could




Figure 1. Patient outcomes after 5-azacytidine treatment. (A) Best response and duration of response are shown on the swimmer plot. Blue indicates a complete response, green a partial response, violet stable disease (SD), and pink progressive disease (PD). Stars indicate deceased patients, arrows ongoing treatment, and vertical lines the cessation of treatment. (B) Kaplan-Meier curve showing overall survival. (C) Kaplan-Meier curve showing progression-free survival (PFS).

provide signals promoting survival and expansion of neoplastic T cells, which would be reversed by 5-azacytidine.

Several other questions remain unanswered, especially whether TFH-derived PTCLs other than AITL, which share similar mutational anomalies with AITL (especially recurrent *TET2*, *DNMT3A*, and *RHOA* mutations),²² respond similarly to 5-azacytidine or whether genetic or epigenetic response markers could be determined to predict which patients could benefit from this treatment. These questions warrant a prospective study, which is planned to start this year (EudraCT #2017-003909-17).

Acknowledgments

This work was supported by the Leukemia Lymphoma Society (grant LLS SCOR 7013-17).

Authorship

Contribution: F.L., O.H., P.G., and R.D. designed the study and wrote the manuscript; F.L., J.D., P.S., O.T., M.C., C.S., A.M., C.H., O.H., and R.D. treated the patients and approved the manuscript; V.F. collected cases and material and approved the manuscript; and F.L., P.S., C.R., L.P., and P.G. performed the histological and molecular studies and approved the study.

Conflict-of-interest disclosure: R.D., J.D., and C.H. received honoraria from Celgene. The remaining authors declare no competing financial interests.

Correspondence: Richard Delarue, Service d'Hématologie Adultes, Hôpital Necker, 149 rue de Sèvres, 75015 Paris, France; e-mail: richard.delarue@gmail.com; and Philippe Gaulard, Département de Pathologie, Hôpitaux Universitaires Henri Mondor, 51 ave de Lattre de Tassigny, 94010 Créteil, France; e-mail: philippe.gaulard@aphp.fr.

Footnotes

*P.G. and R.D. contributed equally to this study.

The online version of this article contains a data supplement.

REFERENCES

- Swerdlow S, Campo E, Harris NL, Jaffe S, and Pileri SA. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues. Lyon, France: IARC; 2017.
- de Leval L, Parrens M, Le Bras F, et al. Angioimmunoblastic T-cell lymphoma is the most common T-cell lymphoma in two distinct French information data sets. *Haematologica*. 2015;100(9):e361-e364.
- Laurent C, Baron M, Amara N, et al. Impact of expert pathologic review of lymphoma diagnosis: study of patients from the French Lymphopath Network. *J Clin Oncol*. 2017;35(18):2008-2017.
- Mak V, Hamm J, Chhanabhai M, et al. Survival of patients with peripheral T-cell lymphoma after first relapse or progression: spectrum of disease and rare long-term survivors. *J Clin Oncol*. 2013;31(16):1970-1976.
- Coiffier B, Pro B, Prince HM, et al. Results from a pivotal, open-label, phase II study of romidepsin in relapsed or refractory peripheral T-cell lymphoma after prior systemic therapy. *J Clin Oncol*. 2012;30(6):631-636.

6. O'Connor OA, Pro B, Pinter-Brown L, et al. Pralatrexate in patients with relapsed or refractory peripheral T-cell lymphoma: results from the pivotal PROPEL study. *J Clin Oncol*. 2011;29(9):1182-1189.
7. O'Connor OA, Horwitz S, Masszi T, et al. Belinostat in patients with relapsed or refractory peripheral T-cell lymphoma: results of the pivotal phase II BELIEF (CLN-19) study. *J Clin Oncol*. 2015;33(23):2492-2499.
8. Lemonnier F, Couronné L, Parrens M, et al. Recurrent TET2 mutations in peripheral T-cell lymphomas correlate with TFH-like features and adverse clinical parameters. *Blood*. 2012;120(7):1466-1469.
9. Odejide O, Weigert O, Lane AA, et al. A targeted mutational landscape of angioimmunoblastic T-cell lymphoma. *Blood*. 2014;123(9):1293-1296.
10. Sakata-Yanagimoto M, Enami T, Yoshida K, et al. Somatic RHOA mutation in angioimmunoblastic T cell lymphoma. *Nat Genet*. 2014;46(2):171-175.
11. Couronné L, Bastard C, Bernard OA. TET2 and DNMT3A mutations in human T-cell lymphoma. *N Engl J Med*. 2012;366(1):95-96.
12. Cairns RA, Iqbal J, Lemonnier F, et al. IDH2 mutations are frequent in angioimmunoblastic T-cell lymphoma. *Blood*. 2012;119(8):1901-1903.
13. Shih AH, Abdel-Wahab O, Patel JP, Levine RL. The role of mutations in epigenetic regulators in myeloid malignancies. *Nat Rev Cancer*. 2012;12(9):599-612.
14. Bejar R, Lord A, Stevenson K, et al. TET2 mutations predict response to hypomethylating agents in myelodysplastic syndrome patients. *Blood*. 2014;124(17):2705-2712.
15. Emadi A, Faramand R, Carter-Cooper B, et al. Presence of isocitrate dehydrogenase mutations may predict clinical response to hypomethylating agents in patients with acute myeloid leukemia. *Am J Hematol*. 2015;90(5):E77-E79.
16. Metzeler KH, Walker A, Geyer S, et al. DNMT3A mutations and response to the hypomethylating agent decitabine in acute myeloid leukemia. *Leukemia*. 2012;26(5):1106-1107.
17. Cheminant M, Bruneau J, Kosmider O, et al. Efficacy of 5-azacytidine in a TET2 mutated angioimmunoblastic T cell lymphoma. *Br J Haematol*. 2015;168(6):913-916.
18. Saillard C, Guermouche H, Derriex C, et al. Response to 5-azacytidine in a patient with TET2-mutated angioimmunoblastic T-cell lymphoma and chronic myelomonocytic leukaemia preceded by an EBV-positive large B-cell lymphoma. *Hematol Oncol*. 2017;35(4):864-868.
19. Cheson BD, Pfistner B, Juweid ME, et al; International Harmonization Project on Lymphoma. Revised response criteria for malignant lymphoma. *J Clin Oncol*. 2007;25(5):579-586.
20. Delfau-Larue M-H, de Leval L, Joly B, et al. Targeting intratumoral B cells with rituximab in addition to CHOP in angioimmunoblastic T-cell lymphoma. A clinicobiological study of the GELA. *Haematologica*. 2012;97(10):1594-1602.
21. Gaulier A, Jary-Bourguignat L, Serna R, Pulik M, Davi F, Raphaël M. Occurrence of angioimmunoblastic T cell lymphoma in a patient with chronic myelomonocytic leukemia features. *Leuk Lymphoma*. 2000;40(1-2):197-204.
22. Dobay MP, Lemonnier F, Missiaglia E, et al. Integrative clinicopathological and molecular analyses of angioimmunoblastic T-cell lymphoma and other nodal lymphomas of follicular helper T-cell origin. *Haematologica*. 2017;102(4):e148-e151.

DOI 10.1182/blood-2018-04-840538

© 2018 by The American Society of Hematology

TO THE EDITOR:

Germline loss-of-function *SAMD9* and *SAMD9L* alterations in adult myelodysplastic syndromes

Yasunobu Nagata,^{1,*} Satoshi Narumi,^{2,*} Yihong Guan,¹ Bartłomiej P. Przychodzen,¹ Cassandra M. Hirsch,¹ Hideki Makishima,³ Hirohito Shima,² Mai Aly,^{1,4} Victor Pastor,⁵ Teodora Kuzmanovic,¹ Tomas Radivoyevitch,^{1,6} Vera Adema,¹ Hassan Awada,¹ Kenichi Yoshida,³ Samuel Li,⁷ Francesc Sole,⁸ Rabi Hanna,⁹ Babal K. Jha,¹ Thomas LaFramboise,⁷ Seishi Ogawa,³ Mikkael A. Sekeres,¹⁰ Marcin W. Wlodarski,⁵ Jörg Cammenga,¹¹ and Jaroslaw P. Maciejewski¹

¹Department of Translational Hematology and Oncology Research, Taussig Cancer Institute, Cleveland Clinic, Cleveland, OH; ²Department of Molecular Endocrinology, National Research Institute for Child Health and Development, Tokyo, Japan; ³Department of Pathology and Tumor Biology, Graduate School of Medicine, Kyoto University, Kyoto, Japan; ⁴Clinical Hematology Unit, Faculty of Medicine, Assiut University, Assiut, Egypt; ⁵Division of Pediatric Hematology and Oncology, Department of Pediatrics and Adolescent Medicine, University of Freiburg, Freiburg, Germany; ⁶Department of Quantitative Health Sciences, Lerner Research Institute, Cleveland Clinic, Cleveland, OH; ⁷Department of Genetics and Genome Sciences, Case Western Reserve University, Cleveland, OH; ⁸Myelodysplastic Syndrome Research Group, Josep Carreras Leukaemia Research Institute, Institut Català d'Oncologia-Hospital Germans Trias i Pujol, Universitat Autònoma de Barcelona, Barcelona, Spain; ⁹Pediatric Hematology Oncology and Blood and Marrow Transplantation and ¹⁰Leukemia Program, Department of Hematology and Medical Oncology, Cleveland Clinic Taussig Cancer Institute, Cleveland Clinic, Cleveland, OH; and ¹¹Department of Hematology, Linköping University, Linköping, Sweden

Familial bone marrow failure (BMF) syndromes present typically in children and younger adults.¹⁻³ A number of germline (GL) mutations in genes such as *DDX41*,⁴ *RUNX1*,⁵ *ETV6*,⁶ *GATA2*,⁷ and *ANKRD26*⁸ have been implicated in the pathogenesis of familial myelodysplastic syndromes (MDSs) and define a disease class of myeloid neoplasms with GL predisposition.⁹ GL *SAMD9* mutations arise in MIRAGE (myelodysplasia, infection, restriction of growth, adrenal hypoplasia, genital phenotypes, and enteropathy) syndrome patients,^{10,11} whereas GL *SAMD9L* mutations occur in pediatric MDS and BMF patients.^{7,12,13} *SAMD9* and *SAMD9L* are proximal on 7q21.2. GL variants in these genes enhance their physiologic growth-inhibitory function and are thus gains of function (GOF).^{12,14,15} Supporting this notion,

somatically acquired $-7/\text{del}(7q)$, or loss-of-function (LOF) missense and truncating mutations affecting the same *SAMD9*/*SAMD9L* mutant allele, revert their GL mutation to escape its inhibitory effects.^{10-12,15-17} We report a 9-month-old infant with familial thrombocytopenia with a *SAMD9L* variant, marrow normocellularity, and the absence of megakaryocytes, which resolved following 4 months of transfusion support (supplemental Figure 1; available on the *Blood* Web site). Whole exome sequencing (WES) identified a novel heterozygous GL variant in a conserved amino acid region of *SAMD9L* (Trp517Arg) with likely deleterious consequences predicted by in silico analysis (supplemental Table 1). Sequencing of the patients' families confirmed that the GL variant was of paternal origin. Although *SAMD9* and *SAMD9L* GL