

BIOSCIENCE REPORTS

ACCEPTED MANUSCRIPT

Comparison of interleukin-10 and interleukin-13 in cord blood of infants born by vaginal delivery and caesarean

Nastaran Khosravi, Nasrin Khalesi, Hamid Sheykhholeslami, Mohammad Nabavi, and Alireza Karimi

Background: Modification of the immunological balance based on the mode of delivery has been recently suggested. The present study assessed the levels of IL-13 and IL-10 in umbilical cord blood of infants born through normal vaginal delivery and infants born with cesarean section.

Methods: This pilot study was performed on 42 neonates born at Rasool-e-Akram hospital between May 2013 and May 2014 categorized into two groups born by vaginal delivery ($n = 21$) and those who born by cesarean section ($n = 21$). The cord blood levels of IL-13 and IL-10 were measured by ELISA technique.

Results: No difference was observed between the two groups with normal vaginal delivery and cesarean delivery in the level of IL-13 in umbilical cord blood (1.42 ± 0.23 versus 1.40 ± 0.22 , respectively, $p = 0.785$). The mean level of IL-10 in umbilical cord blood in the group with vaginal delivery was 6.35 ± 2.54 and in another group with cesarean section was 5.69 ± 2.42 with no significant difference ($p = 0.393$). According to the multivariate linear regression analyses, no difference was found between the two groups of the mode of delivery in the level of IL-10 (beta = -0.454, SE = 0.802, $p = 0.575$) and also in the level of IL-13 (beta = 0.012, SE = 0.076, $p = 0.877$). None of the indicators including gestational age, mother's age, sex of neonate, number of live births, history of abortion, and number of parity could predict increased level of the interleukins in umbilical cord blood.

Conclusion: Mode of delivery may not be an indicator for altering cord blood levels of IL-13 and IL-10.

Cite as Bioscience Reports (2016) DOI: 10.1042/BSR20160147

Copyright 2016 The Author(s).

This is an Accepted Manuscript; not the final Version of Record. You are encouraged to use the final Version of Record that, when published, will replace this manuscript and be freely available under a Creative Commons licence. All other rights reserved.

1 **Background:** Modification of the immunological balance based on the mode of delivery has
2 been recently suggested. The present study assessed the levels of IL-13 and IL-10 in umbilical
3 cord blood of infants born through normal vaginal delivery and infants born with cesarean
4 section.

5 **Methods:** This pilot study was performed on 42 neonates born at Rasool-e-Akram hospital
6 between May 2013 and May 2014 categorized into two groups born by vaginal delivery (n = 21)
7 and those who born by cesarean section (n = 21). The cord blood levels of IL-13 and IL-10 were
8 measured by ELISA technique.

9 **Results:** No difference was observed between the two groups with normal vaginal delivery and
10 cesarean delivery in the level of IL-13 in umbilical cord blood (1.42 ± 0.23 versus 1.40 ± 0.22 ,
11 respectively, $p = 0.785$). The mean level of IL-10 in umbilical cord blood in the group with
12 vaginal delivery was 6.35 ± 2.54 and in another group with cesarean section was 5.69 ± 2.42
13 with no significant difference ($p = 0.393$). According to the multivariate linear regression
14 analyses, no difference was found between the two groups of the mode of delivery in the level of
15 IL-10 (beta = -0.454, SE = 0.802, $p = 0.575$) and also in the level of IL-13 (beta = 0.012, SE =
16 0.076, $p = 0.877$). None of the indicators including gestational age, mother's age, sex of neonate,
17 number of live births, history of abortion, and number of parity could predict increased level of
18 the interleukins in umbilical cord blood.

19 **Conclusion:** Mode of delivery may not be an indicator for altering cord blood levels of IL-13
20 and IL-10.

21 **Keyword:** Delivery; Interleukin-10, Interleukin-13; Cord blood.

22

1 **Introduction**

2 In recent decades, the rate of cesarean section is increasing in most parts of the world [1] and Iran
3 is no exception of this upward trend [2]. The number of live births by caesarean section in
4 developed countries and in developing countries is rising [3]. In contrast, the results from
5 different studies show that the risk of maternal death during cesarean delivery is 3 times higher
6 than that of the vaginal delivery [4]. In addition, cesarean delivery method can be a dangerous
7 procedure not only for the mother but also for the neonates even for many years after birth [5].
8 Recently, it has been found the incidence of asthma and atopy in childhood may be increased by
9 cesarean section delivery [5]. Asthma and atopy are associated directly with the child's immune
10 system. The role of immunoglobulin, such as immunoglobulin E and some other cytokines such
11 as interferon-gamma ($INF\gamma$) in the etiology of asthma has been established [6]. The main
12 pathophysiological basis of allergic asthma is producing some cytokines secreted by T helper
13 cells [7]. A large number of cytokines have been identified as triggers of allergic asthma [8-10]
14 that among them the critical role of interleukins 13 and 4 secreted by T helper cells II is more
15 taken into consideration [5]. Interleukin-13 appears to have the ability to induce and begin all
16 processes of allergic asthma and plays its role independent of immunoglobulin E and eosinophil
17 [5]. Besides, interleukin-10 (IL-10) belongs to the group of type 2 cytokines that its effect is
18 mainly based on activation of the T helper cells type I and II. These cells are the main cells in
19 cellular immunity that particularly regulate IgE production in asthma condition. It seems that IL-
20 10 is a main suppressor of the production of other inflammatory cytokines [4].

21 Recent studies have shown different levels of serum immunoglobulin in both asthmatic and
22 atopy children compared to non-affected ones [3]. Recent evidences have also suggested that
23 cytokines play an important role in delivery process and thus affects the immune system of the

1 newborn [4]. In addition, the effect of mode of delivery on production of cytokines has been also
2 revealed [7]. Due to the increasing trend of cesarean section in our country and also the
3 relationship between the incidence of asthma and atopy and the mode of delivery, we aimed to
4 assess the level of IL-13 and IL-10 in umbilical cord blood of infants born through normal
5 vaginal delivery and infants born with cesarean section.

6

7 **Methods**

8 This pilot study was performed on 42 neonates born at Rasool-e-Akram hospital between May
9 2013 and May 2014. After explaining a detailed description of the study to parents or legal
10 guardians of infants, the written consent was taken from them. The neonates were categorized as
11 the two groups including neonates born by vaginal delivery (n = 21) and those who born by
12 cesarean section (n = 21). We included term and near term pregnancies that were terminated with
13 a normal vaginal delivery or planned cesarean section. The exclusion criteria were the presence
14 of any abnormal symptoms during the first examination of children, preterm delivery, parental
15 immunodeficiency diseases, starting of active phase of labor in cesarean group, chorioamnionitis
16 or any maternal infections and maternal history of asthma and atopy. After coordinating with the
17 laboratory of the hospital, 10cc of umbilical cord blood was extracted from immediately after the
18 completion of the third stage of labor. After centrifugation, the serum was stored in a refrigerator
19 at -20 ° C. Finally, the samples were transferred to the laboratory and the cord blood levels of IL-
20 13 and IL-10 were measured by ELISA kits. Baseline information including gender, family
21 history of asthma or atopy, mode of delivery, number of parity or gravidity, history of still birth

1 or abortion, mother's age, and history of any maternal disorders within pregnancy were asked
2 and recorded in study checklists.

3 Data were analyzed using IBM SPSS statistical software version 21.0 (Armonk, NY: IBM
4 Corp.). Quantitative variables were presented as mean \pm standard deviation, and categorical
5 variables were presented by absolute frequencies and percentages. Continuous variables were
6 compared using t test. Whenever the data did not appear to have normal distribution, Mann-
7 Whitney U test was used. Categorical variables were compared using chi-square test. Fisher
8 exact test was used when more than 20% of cells with expected count of less than 5 had been
9 observed. The multivariate logistic regression model was employed to determine difference in
10 the level of interleukin between the groups with the presence of confounders. P values of ≤ 0.05
11 were considered statistically significant.

12

13 **Results**

14 Baseline characteristics and clinical data are shown in table 1. The two groups with vaginal
15 delivery and cesarean section were similar in mean gestation age, age of mother, sex of neonate,
16 number of live births, history of abortion, number of parity, and also number of gravidity. None
17 of the groups had previous history of asthma or atopy. The mean level of IL-10 in umbilical cord
18 blood in the group with vaginal delivery was 6.35 ± 2.54 and in another group with cesarean
19 section was 5.69 ± 2.42 with no significant difference ($p = 0.393$). Also, no difference was
20 observed between the two groups with normal vaginal delivery and cesarean delivery in the level
21 of IL-13 in umbilical cord blood (1.42 ± 0.23 versus 1.40 ± 0.22 , respectively, $p = 0.785$). In the
22 group with vaginal delivery, no significant correlation was found between the level of IL-10 and
23 IL-13 ($r = -0.252$, $p = 0.271$). Similarly, in another group with cesarean section, the level of IL10
24 was not correlated with IL-13 ($r = -0.123$, $p = 0.595$). Regarding association between the level of

1 interleukins in umbilical cord blood and sex of neonate, it was shown no difference between
2 male and female neonates in the levels of IL-10 (6.45 ± 3.45 versus 5.84 ± 2.01 , $p = 0.577$) and
3 IL-13 (1.34 ± 0.26 versus 1.44 ± 0.20 , $p = 0.238$). Also, gestational age was not correlated with
4 the levels of IL-10 ($r = 0.241$, $p = 0.124$) and also the level of IL-13 ($r = -0.149$, $p = 0.349$).
5 According to the multivariate linear regression analyses (Tables 2 and 3), no difference was
6 found between the two groups of the mode of delivery in the level of IL-10 ($\beta = -0.454$, $SE =$
7 0.802 , $p = 0.575$) and also in the level of IL-13 ($\beta = 0.012$, $SE = 0.076$, $p = 0.877$). In this
8 regard, none of the indicators including gestational age, mother's age, sex of neonate, number of
9 live births, history of abortion, and number of parity could predict increased level of the
10 interleukins in umbilical cord blood.

11 12 **Discussion**

13 According to our analysis, the level of IL-10 and IL-13 is not influenced by the mode of delivery.
14 The effects of the method of delivery on the cord blood level of these cytokines led to
15 contradictory results in previous observations. Similar to our finding, Tutdibi showed that the
16 concentration of IL-10 receptor antagonist was not different between elective cesarean section
17 and vaginal delivery [9]. In another study by Blanco-Quirós *et al.* [10], the type of delivery
18 (vaginal versus caesarean) did not influence cord blood IL-10 results. According to the
19 observations by Bakheit *et al.* [8], concentration of IL-10 in the peripheral and placental sera was
20 higher in vaginal delivery, while cord IL-10 was not significantly different in the two groups.
21 Regarding association between mode of delivery and cord blood level of IL-13, Ly *et al.* [11]
22 showed that cesarean section was associated with increased levels of IL-13. Also, in multivariate

1 analyses, cesarean section was associated with an increment of 79.4 pg/ml in secretion of IL-13.
2 The paradoxical findings in the studies could be affected by various confounding factors. First,
3 the presence of the history of asthma or allergic disorders in the mother could potentially affect
4 the present increased levels of IL-10 and IL-13 in cord blood. On the other hand, lack of
5 significant changes in the levels of these cytokines in our study might be affected by the absence
6 of this history in our studied mothers. Second, it has been well shown that the frequency of
7 cesarean section is frequently higher in preterm neonates who are more susceptible to various
8 allergic and atopic disorders leading increased level of interleukins. In fact, because at present it
9 suggests that cesarean is performed due to individual wishes not clinical judgment, cesarean
10 section is not specified to preterm situation.

11 A few studies have focused an association between cesarean section and increased neonatal
12 secretion of IL-13 and also IL-10. In this regard, our findings can provide a potential
13 immunologic basis for previous reports of an association between cesarean section and clinical
14 conditions related to increase in cytokines such as atopy or asthma [12-19]. The observed
15 association between mode of delivery and neonatal immune responses may be explained by
16 absent or reduced labor in children delivered by cesarean section [5, 20]. The process of labor
17 may directly influence neonatal immune responses, thereby influencing cytokine secretion at
18 birth. Although a relationship between labor and neonatal secretion of IL-13 and IL-10 has not
19 been shown, the stress of labor has been associated with decrease of some immunological cell
20 types such as T lymphocytes and CD4+ T helper cells [21], and also increased some other cells
21 such as neutrophils [22, 23], and natural killer (NK) cells [24, 25] in cord blood. This evidence
22 can also explain some observed relationships between cesarean section and increased susceptible
23 to increase in cytokines and other immunological factors in cord blood. Mohammad Nabavi et al,

1 were found a significantly higher level of IgE in cord blood of neonates delivered via cesarean
2 section [26, 27]. However, immunological status may differ at older age not due a difference in
3 these cytokines, but due to some other factors such as IgE at birth.
4

5 **Conflicts of interest**

6 None declared.
7

8 **Acknowledgement**

9 Thanks to Dr Alireza Karimi for his great help and time to read the manuscript and give us their
10 comments to improve the quality of the manuscript.
11

12 **Funding**

13 This research received no specific grant from any funding agency in the public, commercial, or
14 not-for-profit sectors.
15

16 **Ethical issues**

17 The use of experimental on the human was approved by the committee of the Iran University of
18 Medical Sciences. This study was also entirely adhered to the declaration of the Helsinki in 2008.
19
20

1

2 **References**

- 3 [1] Rehan RSN, N. Prevalence and determinants of caesarean section in a teaching hospital of
4 Pakistan. *J Obstet Gynaecol* 2000;20:479-83.
- 5 [2] Ahmad Nia S, Delavar B, Eini Zinab H, Kazemipour S, Mehryar A, Naghavi M. Caesarean
6 section in the Islamic Republic of Iran: prevalence and some sociodemographic correlates. 2009.
- 7 [3] Miri Farahani L, Abbasi Shavazi MJ. Caesarean section change trends in iran and some
8 demographic factors associated with them in the past three decades. *J Fasa Univ Med Sci*
9 2012;2:127-34.
- 10 [4] Dosa L. Caesarean section delivery, an increasingly popular option. *Bull World Health Organ*
11 2001;79:1173-81.
- 12 [5] Rowe J, Heaton T, Kusel M, Suriyaarachchi D, Serralha M, Holt BJ, et al. High IFN- γ
13 production by CD8+ T cells and early sensitization among infants at high risk of atopy. *J Allergy*
14 *Clin Immunol* 2004;113:710-6.
- 15 [6] Asadullah K, Sterry W, Volk H. Interleukin-10 therapy—review of a new approach.
16 *Pharmacological Rev* 2003;55:241-69.
- 17 [7] Wills-Karp M, Luyimbazi J, Xu X, Schofield B, Neben TY, Karp CL, et al. Interleukin-13:
18 central mediator of allergic asthma. *Science* 1998;282:2258-61.
- 19 [8] Bakheit KH, Bayoumi NK, Adam I. Peripheral, placental and cord cytokines profile in
20 spontaneous labor and elective caesarean section. *Iranian J Immunology* 2008;5:185.
- 21 [9] Tutdibi E, Hunecke A, Lindner U, Monz D, Gortner L. Levels of cytokines in umbilical cord
22 blood in relation to spontaneous term labor. *J Perinat Med.* 2012 Sep;40(5):527-32. doi:
23 10.1515/jpm-2011-0204.

- 1 [10] Blanco-Quirós A, Arranz E, Solis G, Villar A, Ramos A, Coto D. Cord blood interleukin-10
2 levels are increased in preterm newborns. *Eur J Pediatr* 2000;159:420-3.
- 3 [11] Ly NP, Ruiz-Pérez B, Onderdonk AB, Tzianabos AO, Litonjua AA, Liang C, et al. Mode of
4 delivery and cord blood cytokines: a birth cohort study. *Clin Mol Allergy* 2006;4:1.
- 5 [12] Bager P, Melbye M, Rostgaard K, Benn CS, Westergaard T. Mode of delivery and risk of
6 allergic rhinitis and asthma. *J Allergy Clin Immunol* 2003;111:51-6.
- 7 [13] Eggesbø M, Botten G, Stigum H, Nafstad P, Magnus P. Is delivery by cesarean section a
8 risk factor for food allergy? *J Allergy Clin Immunol* 2003;112:420-6.
- 9 [14] Håkansson S, Källén K. Caesarean section increases the risk of hospital care in childhood
10 for asthma and gastroenteritis. *Clin Exp Allergy* 2003;33:757-64.
- 11 [15] Kero J, Gissler M, Grönlund M-M, Kero P, Koskinen P, Hemminki E, et al. Mode of
12 delivery and asthma—is there a connection? *Pediatric Res* 2002;52:6-11.
- 13 [16] Lange J, Ngoumou G, Berkenheide S, Moseler M, Mattes J, Kuehr J, et al. High interleukin-
14 13 production by phytohaemagglutinin-and Der p 1-stimulated cord blood mononuclear cells is
15 associated with the subsequent development of atopic dermatitis at the age of 3 years. *Clin Exp*
16 *Allergy*. 2003;33:1537-43.
- 17 [17] Renz-Polster H, David M, Buist AS, Vollmer W, O'Connor E, Frazier E, et al. Caesarean
18 section delivery and the risk of allergic disorders in childhood. *Clin Exp Allergy*. 2005;35:1466-
19 72.
- 20 [18] Salam MT, Margolis HG, McConnell R, McGregor JA, Avol EL, Gilliland FD. Mode of
21 delivery is associated with asthma and allergy occurrences in children. *Ann Epidemiol*
22 2006;16:341-6.

- 1 [19] Xu B, Pekkanen J, Hartikainen A-L, Järvelin M-R. Caesarean section and risk of asthma
2 and allergy in adulthood. *J Allergy Clin Immunol* 2001;107:732-3.
- 3 [20] Neville WA, Tisler C, Bhattacharya A, Anklam K, Gilbertson-White S, Hamilton R, et al.
4 Developmental cytokine response profiles and the clinical and immunologic expression of atopy
5 during the first year of life. *J Allergy Clin Immunol* 2003;112:740-6.
- 6 [21] Pittard WB, Schleich DM, Geddes KM, Sorensen RU. Newborn lymphocyte
7 subpopulations: the influence of labor. *Am J Obstet Gynecol* 1989;160:151-4.
- 8 [22] Herson VC, Block C, Eisenfeld LI, Maderazo E, Krause PJ. Effect of labor and delivery on
9 neonatal polymorphonuclear leukocyte number and function. *Am J Perinatol* 1992;9:285-8.
- 10 [23] Samelson R, Larkey DM, Amankwah K, Mcconnachie P. Effect of Labor on Lymphocyte
11 Subsets in Full-Term Neonates. *Am J Reprod Immunol* 1992;28:71-3.
- 12 [24] Malamitsi-Puchner A, Protonotariou E, Boutsikou T, Makrakis E, Sarandakou A, Creatsas
13 G. The influence of the mode of delivery on circulating cytokine concentrations in the perinatal
14 period. *Early Human Development* 2005;81:387-92.
- 15 [25] Thilaganathan B, Meher-Homji N, Nicolaidis KH. Labor: an immunologically beneficial
16 process for the neonate. *Am J Obstet Gynecol* 1994;171:1271-2.
- 17 [26] Nabavi M, Ghorbani R, Asadi AM, Faranoush M. Factors associated with cord blood IgE
18 levels. *Asian Pac J Allergy Immunol* 2013;31:157.
- 19 [27] Scirica CV, Gold DR, Ryan L, Abulkerim H, Celedón JC, Platts-Mills TA, et al. Predictors
20 of cord blood IgE levels in children at risk for asthma and atopy. *J Allergy Clin Immunol*
21 2007;119:81-8.
- 22
23

Table 1: Baseline characteristics between the two groups with vaginal delivery and cesarean.

Variable	Vaginal delivery	Cesarean section	P-value
Gestational age	38.52 ± 0.98	38.43 ± 0.60	0.706
Mother's age	30.76 ± 4.99	31.81 ± 4.65	0.486
Male gender	6 (28.6)	6 (28.6)	1.000
Number of live birth			0.100
1	13 (61.9)	7 (33.3)	
2	8 (38.1)	12 (57.1)	
3	0 (0.0)	2 (9.5)	
History of abortion	2 (9.5)	3 (14.3)	0.999
Number of parity			0.268
1	12 (57.1)	7 (33.3)	
2	7 (33.3)	12 (57.1)	
3	2 (9.5)	2 (9.5)	
Number of gravidity			0.624
1	9 (42.9)	6 (28.6)	
2	9 (42.9)	11 (52.4)	
3	3 (14.3)	4 (19.0)	

1
2
3

Table 2: Multivariate linear regression model to assess difference in IL-10 level between the groups with vaginal delivery and cesarean section.

Variable	Unstandardized Coefficients		Standardized Coefficients	t	Sig.
	B	Std. Error	Beta		
Mode of delivery	-0.454	0.802	-0.093	-0.566	0.575
Gestation age	0.379	0.557	0.123	0.680	0.501
Mother's age	-0.108	0.082	-0.209	-1.309	0.199
Sex of neonate	-0.403	0.835	-0.074	-0.482	0.633
Live births	0.094	1.075	0.022	0.087	0.931
Abortion	1.586	1.176	0.210	1.348	0.186
Parity	-0.716	0.990	-0.190	-0.723	0.474

1
2
3
4
5
6
7
8
9
10
11
12

Table 3: Multivariate linear regression model to assess difference in IL-13 level between the groups with vaginal delivery and cesarean section.

Variable	Unstandardized Coefficients		Standardized Coefficients	t	Sig.
	B	Std. Error	Beta		
Mode of delivery	0.012	0.0760	.027	0.155	0.877
Gestation age	-0.013	0.053	-0.048	-0.250	0.804
Mother's age	0.001	0.008	-0.017	-0.099	0.922
Sex of neonate	0.091	0.079	0.186	1.141	0.262
Live births	-0.137	0.102	-0.365	-1.346	0.187
Abortion	0.008	0.112	0.011	0.068	0.946
Parity	0.087	0.094	0.256	0.923	0.362

1
2
3
4
5