

Research Article

RAGE-mediated functional DNA methylated modification contributes to cigarette smoke-induced airway inflammation in mice

Ping Li^{1,*}, Tao Wang^{1,*}, Mei Chen^{2,*}, Jun Chen¹,  Yongchun Shen¹ and  Lei Chen¹

¹Laboratory of Pulmonary Diseases and Department of Respiratory and Critical Care Medicine, West China Hospital, West China School of Medicine, Sichuan University, Chengdu, Sichuan 610041, P.R. China; ²Department of Respiratory and Critical Care Medicine, Chengdu Fifth People's Hospital, Chengdu, Sichuan 611130, P.R. China

Correspondence: Lei Chen (lchens@126.com) or Yongchun Shen (shen_yongchun@126.com)



Our previous study indicated knockout of receptor for advanced glycation end-products (RAGE) significantly attenuated cigarette smoke (CS)-induced airway inflammation in mice. In the present study, we aim to further detect the mediatory effects of RAGE in DNA methylated modification in CS-induced airway inflammation. Lung tissues from the CS-exposed mouse model of airway inflammation were collected for profiling of DNA methylation by liquid hybridization capture-based bisulfite sequencing, which were used for conjoint analysis with our previous data of gene expression by cDNA microarray to identify functional methylated genes, as well as hub genes selected by protein–protein interaction (PPI) network analysis, and functional enrichment analyses were then performed. After RAGE knockout, 90 genes were identified by intersection of the differentially methylated genes and differentially expressed genes. According to the reversed effects of methylation in promoters on gene transcription, 14 genes with functional methylated modification were further identified, among which chemokine (C–X–C motif) ligand 1 (CXCL1), Toll-like receptor 6 (TLR6) and oncostatin M (OSM) with hypomethylation in promoters, were selected as the hub genes by PPI network analysis. Moreover, functional enrichment analyses showed the 14 functional methylated genes, including the 3 hub genes, were mainly enriched in immune-inflammatory responses, especially mitogen-activated protein kinase, tumor necrosis factor, TLRs, interleukin (IL)-6 and IL-17 pathways. The present study suggests that RAGE mediates functional DNA methylated modification in a cluster of 14 targeted genes, particularly hypomethylation in promoters of CXCL1, TLR6 and OSM, which might significantly contribute to CS-induced airway inflammation via a network of signaling pathways.

Introduction

Chronic obstructive pulmonary disease (COPD) is characterized by persistent respiratory symptoms and airflow limitation that are associated with persistent airway inflammation induced by cigarette smoke (CS), a major causative factor for COPD [1].

Receptor for advanced glycation end-products (RAGE), a membrane protein from the immunoglobulin superfamily, has been implicated in the pathogenesis of COPD [2]. Overexpression of RAGE contributed to airway inflammation in CS-associated COPD [3]. Our previous study further indicated knockout (KO) of RAGE gene significantly attenuated CS-induced airway inflammation in a mouse model [4]. However, the mechanisms regarding the effects of RAGE on airway inflammation in COPD remain not clear. Recently, some evidences implied DNA methylation, an epigenetic regulation, played important

*These authors contributed equally to this work.

Received: 06 February 2021
Revised: 14 May 2021
Accepted: 17 May 2021

Accepted Manuscript online:
21 May 2021
Version of Record published:
29 June 2021




Figure 1. Venn diagram of the intersection model

The colorful oval shapes represent CS-associated DEGs (yellow), CS-associated DMGs (blue), RAGE-associated DEGs (green) and RAGE-associated DMGs (purple).

roles in RAGE-mediated inflammatory responses in various diseases [5–7], although the role in DNA methylated modification mediated by RAGE in airway inflammation in COPD was not reported.

In consequence, we did the present study, using the established mouse model of CS-induced airway inflammation, to explore the underlying mechanisms of RAGE-mediated DNA methylated modification in airway inflammation in COPD.

Materials and methods

Animal model

The animal model of CS-induced airway inflammation has been established in our former study [4]. C57BL/6 mice (7–9 weeks old, 20–22 g weight) were used to generate RAGE KO mice through CRISPR/Cas9 gene targeting technology by Bioray Biotechnology (Shanghai, China). The four experimental groups ($n=3$ mice per group) were included in the present study, as follows: (i) wildtype (WT) group, (ii) CS+WT group, (iii) CS+KO group, (iv) WT+KO group. All mice were specific pathogen-free and kept on a 12-h light/12-h dark cycle, at a room temperature of $22 \pm 2^\circ\text{C}$, with free access to food and water. WT and RAGE KO mice were exposed to mainstream CS or room air for 2 h twice daily, 6 days per week for consecutive 4 weeks. After 4-week CS exposure, the mice were anesthetized intraperitoneally with pentobarbital sodium and killed by femoral artery transection. The animal study was approved by the Panel on Laboratory Animal Care of West China Hospital of Sichuan University and took place at the Experimental Animal Center of West China Hospital of Sichuan University.

DNA extraction

The total DNA of lung tissues was extracted and purified by DNeasy Blood Tissue Kit (Qiagen), according to the manufacturer's instructions. Purified DNA was then quantified by NanoDrop2000 Spectrophotometer (Thermo) and agarose gel electrophoresis. Only DNA samples with $A_{260/280}$ ratio between 1.8 and 2.0 were used for further experiments.

Liquid hybridization capture-based bisulfite sequencing

Genomic DNA (1 μg per sample) was randomly fragmented into approximately 200–300 bp by sonication. After purification, the DNA fragments were repaired in the blunt and phosphorylated ends, which were subsequently 3'




Figure 2. PPI network analysis of the candidate genes

Edges indicate correlations between two genes. The bigger circles indicate more related genes and red circles indicate the hub genes.

adenylated and then ligated to the methylated adapter using the SureSelect^{XT} Mouse methyl-seq Library Prep Kit (Agilent). After that, the DNA hybridization was performed using the SureSelectTM Methyl-SeQ Hybridization Kit (Agilent), which covered 100 Mb of mouse genomic regions, including CpG islands, Gencode promoters, tissue-specific differential methylated regions (DMRs) and DNase I hypersensitive sites. The hybridized DNA was subsequently bisulfite-treated using the EZ DNA Methylation-GoldTM Kit (Zymo Research) to convert unmethylated cytosine into uracil according to the manufacturer's instructions. Finally, the treated DNA was amplified by polymerase chain reaction (PCR) and sequenced on Illumina Novaseq PE150.

DNA methylation data analyses

The Fastp software (version 1.2.1) was used to process methylated raw data and remove the low quality reads, including (i) contaminated sequences; (ii) the Q-value of 3' end is less than 20; (iii) reads with less than 15 bp; (iv) reads with 40% base Q-value less than 15; (v) reads containing N bases greater than 5 ($Q = -10 \cdot \log_{10}(P)$, P is the probability of error) [8]. The clean reads were aligned to the reference genome using Bismark software (version 0.19.0) with bowtie2 (version 2.3.4.2) to attain the methylated type, status and proportion [9]. DMRs were subsequently analyzed using the R package methylKit (version 1.6.1) [10] and eDMR [11], with an adjusted P -value < 0.05 and absolute differential

methylation levels (absolute meth. diff > 5%). Finally, related differential methylated genes (DMGs) were located and annotated in the DMRs by the ChIPseeker software.

cDNA microarray

The data of gene expression by cDNA microarray have been obtained in our former study [4]. In the present study, differentially expressed genes (DEGs) were identified by fold-change, and only genes that at least 1.2-fold up-regulated or down-regulated were analyzed.

Candidate genes selection

To identify the candidate genes with methylated modification mediated by RAGE, a novel intersection model was performed (Figure 1). Briefly, CS-associated (CS+WT vs WT) and RAGE-associated (CS+KO vs CS+WT plus WT+KO vs WT) DMGs intersected to get the overlapped DMGs, and so did CS-associated and RAGE-associated DEGs (overlapped DEGs). Then, the overlapped DMGs and DEGs intersected again to select the candidate genes.

Functional methylated genes and hub genes selection

The candidate genes with functional methylated modification, also called functional methylated genes, were identified according to the reversed effects of methylation in promoters on gene transcription. To further identify the hub genes that were most correlated with other genes, protein–protein interaction (PPI) network analysis among the functional methylated genes was performed using the online database STRING [12] and displayed using Cytoscape [13].

Functional enrichment analyses

Functional enrichment analyses on the functional methylated genes, as well as the hub genes, using Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) databases were performed with the clusterProfiler package using *P*-value <0.05 was set as the threshold.

Results

Data production

Based on the liquid hybridization capture-based bisulfite sequencing (LHC-BS) method, 15 Gbp raw sequence data were generated on average for each sample. More than 79% were mapped to at less one genomic position, covering ~77% of the target regions, with an average of 57× sequencing depth per CpG and 12.5% duplication data, and the duplicated sequence reads were filtered for the subsequent analyses.

RAGE-mediated functional DNA methylated modification in targeted genes in CS-induced airway inflammation

After RAGE KO, as reported in our previous study, CS-induced airway inflammation was significantly improved [4]. Meanwhile, 90 overlapping candidate genes were identified via the intersection of DMGs with DEGs (Supplementary Table S1). It is well-known that DNA methylation occurs almost in CpG islands that are primarily located in the regions of promoter and are negatively correlated with gene expression [14]. As a result, 14 genes from the 90 candidate genes, with functional methylated modification in promoters were identified (Table 1), among which, three genes, including chemokine (C–X–C motif) ligand 1 (CXCL1), Toll-like receptor 6 (TLR6) and oncostatin M (OSM) with hypomethylation in promoters were finally selected as the hub genes by PPI network analysis (Figure 2). Furthermore, GO and KEGG analyses indicated the 14 functional methylated genes, including the 3 hub genes, were significantly enriched in immune-inflammatory responses, especially mitogen-activated protein kinase (MAPK), tumor necrosis factor (TNF), TLRs, interleukin (IL)-6 and IL-17 signaling pathways (Supplementary Tables S2 and S3).

Discussion

DNA methylation is closely associated with COPD susceptibility, exacerbation and lung function decline [15,16]. In this process, RAGE may play a regulatory role in DNA methylated modification in COPD, whereas the mechanisms remain unexplained. Consequently, in the present study, the 90 DMGs, regarding RAGE-mediated airway inflammation induced by CS exposure, were initially selected using a novel intersection model, and the functional DNA methylated modification in 14 targeted genes were subsequently identified, especially hypomethylation in CXCL1, TLR6 and OSM promoters, which might significantly contribute to RAGE-mediated airway inflammation in COPD via a network of signaling pathways, such as MAPK, TNF, TLRs, ILs, etc.

Table 1 The 14 functional methylated genes

| Gene | Description | Genomic localization | Methylation change* | Fold change |
|-----------------|--|---|---|-------------|
| <i>Alox5ap</i> | Aachidonate 5-lipoxygenase activating protein | Promoter Distal intergenic 3' UTR | Hyper (15.6131412) Hypo (−12.032621795) Hypo (−13.27280036) | −1.2 |
| <i>Arhgap22</i> | Rho GTPase activating protein 22 | Promoter Promoter | Hyper (10.29366302) Hypo (−6.476953839) | −1.3 |
| <i>Cxcl1</i> | Chemokine (C–X–C motif) ligand 1 | Promoter | Hyper (10.2010746) | −1.2 |
| <i>Dmc1</i> | DNA meiotic recombinase 1 | Promoter | Hypo (9.450360901) | 1.2 |
| <i>Inf2</i> | Inverted formin, FH2 and WH2 domain containing | Promoter Distal intergenic | Hyper (13.93768033) Hyper (13.60386215) | −1.2 |
| <i>Mir1981</i> | MicroRNA 1981 | Promoter | Hypo (−6.417671883) | 1.2 |
| <i>Mmp25</i> | Matrix metalloproteinase 25 | Promoter Promoter | Hyper (15.40303433) Hypo (−13.06629416) | −1.4 |
| <i>Osm</i> | Oncostatin M | Promoter | Hyper (15.05770235) | −1.4 |
| <i>Psmb9</i> | Proteasome 20S subunit β 9 | Promoter | Hyper (10.74327874) | −1.3 |
| <i>Rnaseh2b</i> | Ribonuclease H2, subunit B | Promoter | Hyper (11.78927527) | −1.3 |
| <i>Slc16a6</i> | Solute carrier family 16 member 6 | Promoter Intron | Hyper (8.126762879) Hyper (8.360891276) | −1.2 |
| <i>Srgn</i> | Srglycin | Promoter | Hyper (9.932392771) | −1.3 |
| <i>Tlr6</i> | Toll-like receptor 6 | Promoter | Hyper (17.51844104) | −1.4 |
| <i>Tmod2</i> | Tropomodulin 2 | Promoter | Hypo (−7.597567585) | 1.3 |

*Methylation changes (%) in gene expression by CS+KO vs CS+WT. Abbreviations: Hyper, hypermethylation; Hypo, hypomethylated; Inc, increased variance.

According to the GO and KEGG analyses, the majority of the candidate genes, especially the 14 functional methylated genes widely participated in CS-associated inflammatory-immune responses. In particular, the three hub genes were documented to play important roles in the inflammatory process in COPD. CXCL1 is a member of chemokine subfamily of CXC [17] and the increased level of CXCL1 was detected in the lungs of COPD [18]. CXCL1 served as a chemoattractant for neutrophils migrating from circulation to respiratory tracts, which contributed to neutrophilic inflammation of COPD [19]. TLR6 belongs to the Toll-like receptor family [20], which initiates innate immune responses in airway epithelial cells and triggers inflammatory responses [21]. Furthermore, TLRs activation could increase CXCL1 production in human pulmonary macrophages [22]. OSM, a member of IL-6 subfamily [23], participates in a variety of inflammatory diseases with a high level [24–26]. In COPD, the inflammatory mechanisms for OSM may be stimulating IL-6 production and IL-6 related inflammation, which is positively correlated with pulmonary function decline [23,27]. Noticeably, as suggested in our study, CXCL1, TLR6 and OSM have been documented to be targeted genes of DNA methylated modification in other diseases, such as schizophrenia [28], type 1 diabetes [29] and Richter syndrome [30].

However, some limitations in the present study should be considered. First, the sample size of mice in each group was relatively small, although the minimum requirement for biological repeat was reached. Second, the novel intersection model might be theoretically imperfect. Third, validation was thus needed in future studies.

In summary, the present study performed intersections of DMGs with DEGs in a CS-exposed mouse model and indicated RAGE could mediate functional methylated modification in multiple targeted genes, especially CXCL1, TLR6 and OSM, which might significantly contribute to airway inflammation in COPD.

Data Availability

The data presented in this manuscript are available from the corresponding author (Lei Chen) on reasonable request.

Competing Interests

The authors declare that there are no competing interests associated with the manuscript.

Funding

This work was supported by the National Natural Science Foundation of China [grant number 81970040].

CRediT Author Contribution

Ping Li: Software, Investigation, Visualization, Methodology, Writing—original draft. **Tao Wang:** Software, Investigation, Visualization, Methodology. **Mei Chen:** Software, Investigation, Visualization, Methodology. **Jun Chen:** Software, Investigation. **Yongchun Shen:** Data curation, Supervision, Validation, Writing—review and editing. **Lei Chen:** Conceptualization, Formal Analysis, Writing—review and editing, Funding acquisition.

Abbreviations

COPD, chronic obstructive pulmonary disease; CS, cigarette smoke; CXCL1, chemokine (C–X–C motif) ligand 1; DEG, differentially expressed gene; DMG, differential methylated gene; DMR, differential methylated region; GO, Gene Ontology; IL, interleukin; KEGG, Kyoto Encyclopedia of Genes and Genomes; KO, knockout; MAPK, mitogen-activated protein kinase; OSM, oncostatin M; PPI, protein–protein interaction; RAGE, receptor for advanced glycation end-product; TLR, Toll-like receptor; TNF, tumor necrosis factor; WT, wildtype.

References

- Vogelmeier, C.F., Criner, G.J., Martinez, F.J., Anzueto, A., Barnes, P.J., Bourbeau, J. et al. (2017) Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Lung Disease 2017 Report. GOLD Executive Summary. *Am. J. Respir. Crit. Care Med.* **195**, 557–582, <https://doi.org/10.1164/rccm.201701-0218PP>
- Yonchuk, J.G., Silverman, E.K., Bowler, R.P., Agustí, A., Lomas, D.A., Miller, B.E. et al. (2015) Circulating soluble receptor for advanced glycation end products (sRAGE) as a biomarker of emphysema and the RAGE axis in the lung. *Am. J. Respir. Crit. Care Med.* **192**, 785–792, <https://doi.org/10.1164/rccm.201501-0137PP>
- Chen, L., Wang, T., Guo, L.L., Shen, Y.C., Yang, T., Wan, C. et al. (2014) Overexpression of RAGE contributes to cigarette smoke-induced nitric oxide generation in COPD. *Lung* **192**, 267–275, <https://doi.org/10.1007/s00408-014-9561-1>
- Chen, M., Wang, T., Shen, Y., Xu, D., Li, X., An, J. et al. (2017) Knockout of RAGE ameliorates mainstream cigarette smoke-induced airway inflammation in mice. *Int. Immunopharmacol.* **50**, 230–235, <https://doi.org/10.1016/j.intimp.2017.06.018>
- Kan, S., Wu, J., Sun, C., Hao, J. and Wu, Z. (2018) Correlation between RAGE gene promoter methylation and diabetic retinal inflammation. *Exp. Ther. Med.* **15**, 242–246
- Maslinska, D., Laure-Kamionowska, M. and Maslinski, S. (2014) Methyl-CpG binding protein 2, receptors of innate immunity and receptor for advanced glycation end-products in human viral meningoencephalitis. *Folia Neuropathol.* **52**, 428–435, <https://doi.org/10.5114/fn.2014.47844>
- Zhang, M., Li, Y., Rao, P., Huang, K., Luo, D., Cai, X. et al. (2018) Blockade of receptors of advanced glycation end products ameliorates diabetic osteogenesis of adipose-derived stem cells through DNA methylation and Wnt signalling pathway. *Cell Prolif.* **51**, e12471, <https://doi.org/10.1111/cpr.12471>
- Chen, S.F., Zhou, Y.Q., Chen, Y.R. and Gu, J. (2018) fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**, 884–890, <https://doi.org/10.1093/bioinformatics/bty560>
- Krueger, F. and Andrews, S.R. (2011) Bismark: a flexible aligner and methylation caller for Bisulfite-Seq applications. *Bioinformatics* **27**, 1571–1572, <https://doi.org/10.1093/bioinformatics/btr167>
- Akalin, A., Kormaksson, M., Li, S., Garrett-Bakelman, F.E., Figueroa, M.E., Melnick, A. et al. (2012) methylKit: a comprehensive R package for the analysis of genome-wide DNA methylation profiles. *Genome Biol.* **13**, R87, <https://doi.org/10.1186/gb-2012-13-10-r87>
- Li, S., Garrett-Bakelman, F.E., Akalin, A., Zumbo, P., Levine, R., To, B.L. et al. (2013) An optimized algorithm for detecting and annotating regional differential methylation. *BMC Bioinformatics* **14**, S10, <https://doi.org/10.1186/1471-2105-14-S5-S10>
- Szklarczyk, D., Morris, J.H., Cook, H., Kuhn, M., Wyder, S., Simonovic, M. et al. (2017) The STRING database in 2017: quality-controlled protein-protein association networks, made broadly accessible. *Nucleic Acids Res.* **45**, D362–D368, <https://doi.org/10.1093/nar/gkw937>
- Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D. et al. (2003) Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res.* **13**, 2498–2504, <https://doi.org/10.1101/gr.1239303>
- Bender, C.M., Gonzalgo, M.L., Gonzales, F.A., Nguyen, C.T., Robertson, K.D. and Jones, P.A. (1999) Roles of cell division and gene transcription in the methylation of CpG islands. *Mol. Cell. Biol.* **19**, 6690–6698, <https://doi.org/10.1128/MCB.19.10.6690>
- Vucic, E.A., Chari, R., Thu, K.L., Wilson, I.M., Cotton, A.M., Kennett, J.Y. et al. (2014) DNA methylation is globally disrupted and associated with expression changes in chronic obstructive pulmonary disease small airways. *Am. J. Respir. Cell Mol. Biol.* **50**, 912–922, <https://doi.org/10.1165/rcmb.2013-03040C>
- Busch, R., Qiu, W.L., Lasky-Su, J., Morrow, J., Criner, G. and DeMeo, D. (2016) Differential DNA methylation marks and gene comethylation of COPD in African-Americans with COPD exacerbations. *Respir. Res.* **17**, 15, <https://doi.org/10.1186/s12931-016-0459-8>
- Charo, I.F. and Ransohoff, R.M. (2006) Mechanisms of disease - The many roles of chemokines and chemokine receptors in inflammation. *N. Engl. J. Med.* **354**, 610–621, <https://doi.org/10.1056/NEJMra052723>
- Traves, S.L., Culpitt, S.V., Russell, R.E.K., Barnes, P.J. and Donnelly, L.E. (2002) Increased levels of the chemokines GRO α and MCP-1 in sputum samples from patients with COPD. *Thorax* **57**, 590–595, <https://doi.org/10.1136/thorax.57.7.590>
- Hikichi, M., Mizumura, K., Maruoka, S. and Gon, Y. (2019) Pathogenesis of chronic obstructive pulmonary disease (COPD) induced by cigarette smoke. *J. Thorac. Dis.* **11**, S2129–S2140, <https://doi.org/10.21037/jtd.2019.10.43>
- Takeuchi, O., Kawai, T., Sanjo, H., Copeland, N.G., Gilbert, D.J., Jenkins, N.A. et al. (1999) TLR6: a novel member of an expanding Toll-like receptor family. *Gene* **231**, 59–65, [https://doi.org/10.1016/S0378-1119\(99\)00098-0](https://doi.org/10.1016/S0378-1119(99)00098-0)

- 21 Koff, J.L., Shao, M.X.G., Ueki, I.F. and Nadel, J.A. (2008) Multiple TLRs activate EGFR via a signaling cascade to produce innate immune responses in airway epithelium. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **294**, L1068–L1075, <https://doi.org/10.1152/ajplung.00025.2008>
- 22 Grassin-Delyle, S., Abrial, C., Salvator, H., Brolo, M., Naline, E. and Devillier, P. (2020) The role of Toll-like receptors in the production of cytokines by human lung macrophages. *J. Innate Immun.* **12**, 63–73, <https://doi.org/10.1159/000494463>
- 23 Tanaka, M. and Miyajima, A. (2003) Oncostatin M, a multifunctional cytokine. *Rev. Physiol. Biochem. Pharmacol.* **149**, 39–52, <https://doi.org/10.1007/s10254-003-0013-1>
- 24 Hasegawa, M., Sato, S., Fujimoto, M., Ihn, H., Kikuchi, K. and Takehara, K. (1998) Serum levels of interleukin 6 (IL-6), oncostatin M, soluble IL-6 receptor, and soluble gp130 in patients with systemic sclerosis. *J. Rheumatol.* **25**, 308–313
- 25 Pradeep, A.R., S, T.M., Garima, G. and Raju, A. (2010) Serum levels of oncostatin M (a gp 130 cytokine): an inflammatory biomarker in periodontal disease. *Biomarkers* **15**, 277–282, <https://doi.org/10.3109/13547500903573209>
- 26 Liang, H., Block, T.M., Wang, M., Nefsky, B., Long, R., Hafner, J. et al. (2012) Interleukin-6 and oncostatin M are elevated in liver disease in conjunction with candidate hepatocellular carcinoma biomarker GP73. *Cancer Biomark.* **11**, 161–171, <https://doi.org/10.3233/CBM-2012-00276>
- 27 Rincon, M. and Irvin, C.G. (2012) Role of IL-6 in asthma and other inflammatory pulmonary diseases. *Int. J. Biol. Sci.* **8**, 1281–1290, <https://doi.org/10.7150/ijbs.4874>
- 28 Zhou, C., Chen, J., Tang, X., Feng, X., Yu, M., Sha, W. et al. (2018) DNA methylation and gene expression of the chemokine (C-X-C motif) ligand 1 in patients with deficit and non-deficit schizophrenia. *Psychiatry Res.* **268**, 82–86, <https://doi.org/10.1016/j.psychres.2018.06.067>
- 29 Xie, Z., Huang, G., Wang, Z., Luo, S., Zheng, P. and Zhou, Z. (2018) Epigenetic regulation of Toll-like receptors and its roles in type 1 diabetes. *J. Mol. Med.* **96**, 741–751, <https://doi.org/10.1007/s00109-018-1660-7>
- 30 Rinaldi, A., Mensah, A.A., Kwee, I., Forconi, F., Orlandi, E.M., Lucioni, M. et al. (2013) Promoter methylation patterns in Richter syndrome affect stem-cell maintenance and cell cycle regulation and differ from de novo diffuse large B-cell lymphoma. *Br. J. Haematol.* **163**, 194–204