

Identification of a mitochondrial-binding site on the N-terminal end of hexokinase II

Nadezda Bryan* and Kevin P. Raisch*¹

*Department of Otolaryngology, University of Florida, Gainesville, FL 32610, USA

Synopsis

Hexokinase II (HKII) is responsible for the first step in the glycolysis pathway by adding a phosphate on to the glucose molecule so it can proceed down the pathway to produce the energy for continuous cancer cell growth. Tumour cells overexpress the HKII enzyme. In fact, it is the overexpression of the HKII enzyme that makes the diagnosis of cancer possible when imaged by positron emission tomography (PET). HKII binds to the voltage-dependent anion channel (VDAC) located on the mitochondrial outer membrane (MOM). When bound to the MOM, HKII is blocking a major cell death pathway. Thus, HKII is responsible for two characteristics of cancer cells, rapid tumour growth and inability of cancer cells to undergo apoptosis. One method to identify novel compounds that may interfere with the HKII–VDAC-binding site is to create a molecular model using the crystal structure of HKII. However, the amino acid(s) responsible for HKII binding to VDAC are not known. Therefore, a series of truncations and point mutations were made to the N-terminal end of HKII to identify the binding site to VDAC. Deletions of the first 10 and 20 amino acids indicated that important amino acid(s) for binding were located within the first 10 amino acids. Next, a series of point mutations were made within the first 10 amino acids. It is clear from the immunofluorescence images and immunoblot results that mutating the fifth amino acid from histidine to proline completely abolished binding to the MOM.

Key words: cancer, glycolysis, hexokinase II, mitochondrial-binding site, mutations.

Cite this article as: Bioscience Reports (2015) 35, e00205, doi:10.1042/BSR20150047

INTRODUCTION

In 1931 Dr Otto Warburg won the Nobel Prize in Physiology and Medicine for his discoveries associated with respiratory enzymes. Dr Warburg's early studies, known as the Warburg effect, were the first to describe the high rate of glucose uptake by tumour cells with an increased release of lactic acid [1,2]. Dr Warburg hypothesized that tumour cell growth is caused by a defect in the mitochondria leading to an increased reliance on the breakdown of glucose (glycolysis) to generate ATP [3]. Subsequently, studies have shown that tumour cells overexpress hexokinase II (HKII) and the overexpression of HKII plays a critical role in maintaining the high rate of glycolysis in tumour cells. HKII has a direct relationship with two major hallmarks of tumour cells: (1) the increased rate of glycolysis and subsequent ATP production and (2) the decreased ability to undergo apoptosis. HKII binds to the voltage-dependent anion channel (VDAC) on the mitochondrial outer membrane (MOM), thereby blocking BAX

(BCL2-associated X protein) from activating the mitochondrial-related cell death pathway.

Subcellular studies have shown that HKII is catalytically active when bound to the MOM [4,5]. Agents that induce the translocation of HKII from the MOM to the cytoplasm have been shown to disrupt the glycolysis pathway and sensitize tumour cells to chemotherapy drugs [6–8]. Our laboratory has shown that the translocation of HKII from the MOM to the cytoplasm arrests the cells in the late G₁-cell cycle phase and thereby sensitizes tumour cells to radiation treatment [9].

The goal of the present study was to identify the amino acid(s) in HKII that are associated with binding to VDAC on the MOM. The binding site to VDAC has been reported to be within the first 15–23 amino acids of the N-terminal end [10–12]. Therefore, a series of truncations and point mutations were made to identify the amino acids critical for binding to VDAC. Deletions of the first 10 and 20 amino acids indicated that important amino acids for binding to the MOM were located within the first 10 amino acids. A series of single point mutations were created within the

Abbreviations: aa, amino acid; BAX, BCL2-associated X protein; CIAP, calf intestinal alkaline phosphatase; FLHKII, full-length HKII; GFP, green fluorescent protein; HKII, hexokinase II; MOM, mitochondrial outer membrane; PCR, polymerase chain reaction; PET, positron emission tomography; siRNA, small interfering RNA; VDAC, voltage-dependent anion channel.

¹ To whom correspondence should be addressed (email raisch@ufl.edu).

Table 1 Hexokinase II mutations were made in the putative VDAC-binding site located on the N-terminal end

Deletion mutations of 20 amino acids (pI2_Q20del) and 10 amino acids (pI2_F10del) by PCR and the clones analysed for binding to the mitochondrial membrane. Using PredictProtein [14,15], 3 amino acids were predicted to bind protein within the 10 amino acid deletion. Single point mutations at amino acids 4(S4L), 5(H5P) and 8(A8L) were made by PCR and the clones analysed for binding to the mitochondrial membrane. Mutations depicted in red.

Name	N-terminal end
wtHKII	MIASHLLAYFFTELNHDQVQKVDQYLYHMRLS.....
HKII (pI2_Q20del)	M.....VDQYLYHMRLS.....
HKII (pI2_F10del)	M.....TELNHDQVQKVDQYLYHMRLS.....
S4L-HKII	MIALHLLAYFFTELNHDQVQKVDQYLYHMRLS.....
H5P-HKII	MIAS ^P LLAYFFTELNHDQVQKVDQYLYHMRLS.....
A8L-HKII	MIASHLL ^L YFFTELNHDQVQKVDQYLYHMRLS.....

first 10 amino acids of the N-terminal end of HKII. All HKII proteins were tagged with green fluorescent protein (GFP) in order to follow their expression and localization by fluorescence microscopy. Immunoblot analyses were used on mitochondrial and cytoplasmic fractions to confirm the microscopic observations presented herein.

EXPERIMENTAL

Mutating hexokinase II and plasmid constructs

Deletions and point mutations were made using the HKII eukaryotic expression vector, pFLHKII-pGFP-N3, developed by Dr Ardehali and purchased from Addgene as the template for PCR-based mutations [13]. All PCR primers were synthesized by Eurofins Genomics. Since the VDAC-binding site was predicted to be within the first 20 amino acids of the N-terminal end of HKII, two deletion mutants of HKII were initially designed to narrow the region for point mutations (Table 1). Deletions of the first 10 (HKII pI2_F10del) and 20 (HKII pI2_Q20del) amino acids of the N-terminal end of HKII were cloned in-frame with GFP. The downstream PCR primers were designed to incorporate an EcoRI restriction enzyme site downstream of the start codon and an annealing region of 26 nucleotides upstream of the deleted region. The downstream primers were GCTAGCCTACGATTCTGAATTCATGTTACGGAGCTCAACCATGACCAAGT (pI2_F10del) and GCTAGCCTACGATTCTGAATTCATGAAGGTTGACCAGTACTCTACCACAT (pI2_Q20del).

Using the ISIS method in PredictProtein (<https://www.predictprotein.org>), 3 amino acids were predicted to bind proteins within the first 10 amino acids of HKII [14,15]. The following downstream primers were synthesized to incorporate point mutations in the fourth amino acid, serine to leucine (S4L), fifth amino acid histidine to proline (H5P) and the eighth amino acid alanine to leucine (A8L): GCTAGCCTACGATTCTGAATTCATGATTGCCCTTGACATCT-

GCTTGCCACTTCTTCACGGAGCTAACCATGACCAAGT (S4L); GCTAGCCTACGATTCTGAATTCATGATTGCCCTCGCTCTGCTTGCCTACTTCTTCACGGAGCTCAACCATGACCAAGT (H5P); GCTAGCCTACGATTCTGAATTCATGATTGCCCTCGCATCTGCTTCTTACTTCTTCACGGAGCTCAACCATGACCAAGT (A8L). The upstream PCR primer (ATGGTCCCAACTG-TGTCATT) incorporated an EcoRV restriction enzyme site and was used in all PCR amplifications. For each PCR, the reaction buffer consisted of 1.5 mM MgCl₂, 0.2 mM each dNTP, 10 pmol/50 µl each upstream and downstream primer, 2 ng plasmid DNA and 2.5 units GoTaq polymerase (Promega). The reaction conditions for all primer sets were 95 °C for 2 min followed by 35 cycles of 95 °C for 30 s, 55 °C for 1 min, 72 °C for 1 min with a final 5 min extension at 72 °C. All PCR amplicons were separated using 1% agarose gel electrophoresis, stained with ethidium bromide and detected by UV light using a VersaDoc 1000 camera system (Bio-Rad Laboratories).

Amplicons with an expected size of approximately 600 bp were excised from the agarose gel using X-tractor gel extractor tool (Promega) and eluted from the gel using Wizard SV Gel and PCR Clean-Up System (Promega). PCR amplicons and pFLHKII-pGFP-N3 were cut with the restriction enzyme, EcoRI (Promega), for 1 h at 37 °C, then heat inactivated and precipitated after adding one-tenth volume 5 M NaCl, 2 volumes ethanol and 1 µg glycogen as a carrier. The EcoRI cut amplicons and plasmid was suspended in water then cut with the restriction enzyme, EcoRV (Promega), for 1 h at 37 °C. Following heat-inactivation, the double-digested plasmid was separated by 1% agarose gel electrophoresis and eluted from the gel as described above. The double-digested amplicons and plasmid were precipitated as described above. Double-digested plasmid DNA was suspended in water and reacted with calf intestinal alkaline phosphatase (CIAP, Promega) as described by the manufacturer's protocol then heat inactivated and the DNA precipitated with glycogen as a carrier. Each amplicon and plasmid was suspended in water, then the amplicons were ligated to the plasmid DNA using T4 DNA ligase (Promega) for 18 h at 4 °C using a vector:insert ratio of 3:1. Competent JM109 cells (Promega) were transformed with the ligated plasmid DNA and colonies were selected on LB agar plates containing kanamycin (50 µg/ml). Following a 24 h incubation at 37 °C, individual colonies were selected and grown overnight in LB broth with kanamycin at 37 °C. Plasmid DNA was isolated from the overnight broth cultures using PureYield Plasmid miniprep system (Promega). Plasmid DNA was submitted for sequencing to Eurofins Genomics using primers upstream and downstream of the ligation sites. Sequencing results were analysed using MacVector Software v12.7.5 and plasmids containing the correct truncation or point mutation in-frame with HKII-GFP were used for transfection experiments.

Transfection of U-2OS cells

U-2OS cells (A.T.C.C.) were transfected with eukaryotic expression vectors containing truncations and point mutations of HKII using FuGene HD (Promega) as described in the manufacturer's protocol. Forty-eight hours after transfection, G418 (500 µg/ml)

was added to the culture medium and the cells were maintained in G418-containing culture medium for 4 weeks. To select for stably-transfected cell lines, the G418 was removed from the culture medium for 2–4 weeks. Then G418 was added back to the culture medium and the cells were passaged for an additional 4 weeks before removing G418 from the culture medium. The surviving cell lines were considered stably-transfected with the plasmid DNA and were used for analysis by fluorescent microscopy and immunoblot analysis.

Detection of HKII protein by immunoblot and fluorescence microscopy

Stably-transfected cells were grown in 4-chamber slides (Thermo Scientific Nunc) until 60–80% confluence. Living cell fluorescence images were captured using an AMG EVOS FL Cell Imaging System with a Plan Fluorite 40×/0.65 objective.

Immunoblots were used to confirm the redistribution of HKII and mutated HKII proteins into the cytoplasm from the mitochondria. The enriched-mitochondrial fractions and cytoplasmic fractions were collected and processed as described previously [9]. Equal concentrations of protein were separated by 10% SDS/PAGE and transferred to Immobilon-P membrane (Millipore Corp.) by electroblotting. HKII, mitochondrial Hsp70 (mtHsp70), and β -actin were identified by immunoblotting using the following antibodies purchased from Thermo Scientific; anti-HKII, anti-mtHsp70, anti- β -actin, anti-mouse-HRP, anti-rabbit-HRP. Antibodies targeting β -actin and mtHsp70 were used as loading controls for cytoplasmic and mitochondrial fractions, respectively. Protein was detected by chemi-luminescent signalling (Pierce ECL) and the immunoblots were scanned and analysed by densitometry using UN-SCAN-IT software (Silk Scientific, Inc.). The densitometry results were expressed as a ratio of the intensity of HKII to that of β -actin or Hsp70 from the same source.

RESULTS

Analysis of HKII deletion mutations

For the initial analysis of HKII binding to the MOM, deletion mutants were cloned into a vector and expressed in U-2OS cells. On the N-terminal end of HKII, the first 10 (pI2_F10del) and first 20 (pI2_Q20del) amino acids were deleted following the start codon, Met (Table 1). The wild-type FLHKII (full-length HKII) protein and the deletion mutants were cloned in-frame with the GFP-tag to allow for observation and localization by fluorescence microscopy. For the wild-type FLHKII enzyme (Figure 1A) the fluorescence was observed on the mitochondrial-like structures located around the nuclei. For the two deletion mutants (Figures 1B and 1C), a diffuse fluorescence was observed throughout the cytoplasm indicating that the first 10 and first 20 amino acids were important for binding to the MOM. To confirm the results of the fluorescence microscopy, cytoplasmic- and mitochondrial-

enriched fractions were collected from stably-transfected cell lines expressing wild-type HKII and truncated mutants of the HKII. Wild-type FLHKII was detected by immunoblot in the cytoplasmic- and mitochondrial-enriched fractions (Figure 1D). However, the HKII deletion mutant proteins, pI2_F10del and pI2_Q20del, were absent from the mitochondrial-enriched fractions and only detected in the cytoplasmic-enriched fraction. In the cytoplasmic fraction, a 1.7-fold increase in HKII pI2_F10del and a 3-fold decrease in HKII pI2_Q20del protein expression were found as compared with the levels of FLHKII.

Analysis of HKII point mutations

The results from the deletion mutations indicated that the amino acids required for binding to the MOM were located within the first 10 amino acids of the N-terminal end of HKII. Therefore the HKII amino acid sequence was submitted to PredictProtein and using the protein–protein interaction sites (ISIS) method, the fourth (serine), fifth (histidine), and eighth (alanine) amino acids within the first 10 amino acids were predicted to bind protein. Using PCR-based site-directed mutagenesis the serine and alanine were mutated to a leucine and the histidine was mutated to a proline (Table 1). The resultant transfected U-2OS cell lines were screened by fluorescence microscopy using the GFP-tag to visualize the location of the HKII proteins. The S4L-HKII and the A8L-HKII mutated proteins were found to maintain binding to mitochondrial-like membrane structures as observed with the wild-type FLHKII protein (Figures 2A–2D). Whereas, a diffuse fluorescence was observed in the cytoplasm of the cells expressing the H5P-HKII mutated protein (Figure 2C). To confirm the binding of HKII proteins to the MOM, mitochondrial-enriched and cytoplasmic-enriched fractions were isolated from stably-transfected cell lines. As shown in Figure 2E, the presence of HKII protein in the mitochondrial-enriched fractions was identified from cell lines expressing FLHKII, S4L-HKII and A8L-HKII protein but no HKII protein was detected from the H5P-HKII expressing cell line. A greater than 6-fold increase in S4L-HKII mutant protein was found in the mitochondrial fraction as compared with the wild-type FLHKII protein. In the cytoplasmic-enriched fractions, HKII protein was detected from cell lines expressing FLHKII, S4L-HKII, H5P-HKII and A8L-HKII. A greater than 4-fold decrease in H5P-HKII protein levels as compared with wild-type HKII protein levels was found in the cytoplasmic fraction. The results from the point mutations indicate that the basic, hydrophilic histidine residue at position 5 is critical for HKII binding to the MOM.

DISCUSSION

There are four isoenzymes of hexokinase in mammalian cell lines. Types I, II, III and IV with types I–III expressed as 100 kDa proteins and thought to have been derived from a gene duplication/fusion of an ancestral 50 kDa hexokinase [16]. Type IV




Figure 1 Fluorescent and immunoblot images of the HKII truncation mutants

Stably-transfected U-2OS cells were grown in chamber slides with a plasmid carrying FLHKII, a 10aa deletion or a 20aa deletion in the N-terminal end of HKII. The cells were imaged at 400 \times magnification using an EVOS FL microscope. (A) U-2OS cells transfected with GFP-tagged wild-type FLHKII; (B) U-2OS cells transfected with GFP-tagged HKII (pI2_F10del); (C) U-2OS cells transfected with GFP-tagged HKII (pI2_Q20del); (D) immunoblot analysis of mitochondrial (Mito) and cytoplasmic (Cyto) fractions of U-2OS cells stably-transfected with wild-type FLHKII-GFP, HKII-GFP (pI2_F10del), or HKII-GFP (pI2_Q20del). Immunoblots were probed with primary antibodies to detect: hexokinase II, β -actin and mtHsp70. Densitometry was used to quantify the expression levels of HKII in the cytoplasmic and mitochondrial cell fractions.

hexokinase is a 50 kDa protein commonly referred to as glucokinase. In addition, HKII has two catalytic sites as compared with one site on types I, III and IV [17,18]. Early studies have shown that hexokinase I and II bind to a MOM pore-forming protein and have referred to the protein as VDAC or mitochondrial porin

[19–23]. Subsequent studies have identified the N-terminal domain of hexokinase I and II as necessary for binding to VDAC [24–26].

Using a peptide corresponding to the N-terminal domain for HKII, Pastorino et al. [27] has shown that the first 15 amino acids




Figure 2 Fluorescent and immunoblot images of the HKII mutants with single point mutations

U-2OS cells were transfected with plasmids carrying FLHKII or a point mutation of amino acids in position 4, 5, or 8. Stably-transfected cells were imaged at 400 \times magnification using an EVOS FL microscope. (A) U-2OS cells transfected with GFP-tagged wild-type FLHKII; (B) U-2OS cells transfected with (S4L)-HKII-GFP; (C) U-2OS cells transfected with (H5P)-HKII-GFP; (D) U-2OS cells transfected with (A8L)-HKII-GFP; (E) mitochondrial (Mito) and cytoplasmic (Cyto) fractions were isolated from U-2OS cells stably-transfected with wild-type FLHKII-GFP and each mutant HKII protein, then analysed by immunoblot. Immunoblots were probed with primary antibodies to detect: hexokinase II, β -actin and mtHsp70. Densitometry was used to quantify the expression levels of HKII in the cytoplasmic and mitochondrial cell fractions.

were important for binding to the MOM. We have presented evidence that the fifth amino acid, histidine, on the N-terminal end of HKII is an important component of the binding site to VDAC. Converting the positively charged, hydrophilic histidine to an uncharged, hydrophobic proline completely prevented the binding of HKII to the MOM. Studies of hexokinase I by the Honzatko laboratory have shown that mutating the fifth amino acid, glutamine to proline (Q5P) resulted in a 10-fold decrease in binding to the mitochondrial membrane [28].

When hexokinase is bound to the MOM it is not a substrate for the ubiquitin system, however unbound hexokinase protein was shown to be a target for the ubiquitination pathway and rapidly degraded [29,30]. We hypothesize that the reduced levels of mutant hexokinase protein found in the cytoplasm are due to ubiquitination but further studies are necessary to determine the fate of the mutant proteins.

Mutating the fourth amino acid from the uncharged, polar, hydrophilic serine to an uncharged, hydrophobic leucine has shown an increased stability of the mutant protein by showing higher protein levels found in the cytoplasm and mitochondrial fractions. The S4L-HKII mutant protein had greater than 6-fold affinity for the mitochondrial membrane as compared with FLHKII. We have made double-mutants of the fourth and fifth amino acids of HKII protein (S4L-H5P-HKII) and have shown that the double-mutant HKII protein maintains the ability to bind to the MOM (unpublished data).

Since mutating of the fourth amino acid enhanced the binding of HKII to the MOM, the relevance to the development of cancers needs to be further investigated. Hexokinase type II was shown to be overexpressed in tumour cells, conferring a 100-fold increase in enzymatic activity relative to normal cells [31]. Subcellular studies have shown that HKII is catalytically active when bound to the MOM, giving HKII preferential access to mitochondrial generated ATP [4,5]. Studies have shown that when HKII expression is down-regulated by siRNA or HKII is prevented from binding to the VDAC on the MOM, the tumour cells are sensitized to chemotherapy agents [6–8,32] and radiation [9]. Alternately, the phosphorylation of VDAC by glycocin synthase kinase 3 β (GSK3 β) disrupted the ability of HKII to bind to VDAC, potentiating the doxorubicin- and paclitaxel-induced cell death [6].

Using a structure-based virtual screen, we propose to identify drug-like small molecules with a putative mechanism-of-action of translocating HKII from the MOM to the cytoplasm. The results from the present study have identified the histidine located at the fifth position of the N-terminal end of HKII to be critical for binding to the MOM. To pursue a molecular docking screen for the binding site, the complete crystal structure of HKII must be obtained. The Toronto Structural Genomics Consortium (TSGC) identified the crystal structure of HKII [33], however the crystal structure was missing the first 16 N-terminal amino acids. The crystal structures of human hexokinase I are missing the first 10 or 15 amino acids on the N-terminal end [34,35]. The high-resolution crystal structure of yeast HKII does maintain the N-terminal end and the secondary structure was described as an alpha-helix (residues 1–12) and a six-stranded mixed β -sheet (residues 13–76) [36]. The hydrophobic N-terminal region of



human HKII has been described as important for binding to the MOM [11,13,37]. We present evidence that the fifth amino acid, histidine, located in the N-terminal domain of HKII is critical for binding to the MOM. Upon crystallization of the complete HKII enzyme, the secondary structure surrounding the histidine can be analysed by structure-based virtual screen to identify putative drug-like compounds that may interfere with the binding of HKII to VDAC on the MOM.

AUTHOR CONTRIBUTION

Kevin Raisch and Nadezda Bryan established the experimental design and contributed to the experimental work. Kevin Raisch prepared the manuscript.

FUNDING

This work was supported by the University of Florida Health Cancer Center and the Department of Otolaryngology (to K.P.R.); and the UF Interdisciplinary Center for Biotechnology Research (ICBR).

REFERENCES

- 1 Warburg, O., Wind, F. and Negelein, E. (1927) The metabolism of tumors in the body. *J. Gen. Physiol.* **8**, 519–530 [CrossRef](#) [PubMed](#)
- 2 Vander Heiden, M.G., Cantley, L.C. and Thompson, C.B. (2009) Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science* **324**, 1029–1033 [CrossRef](#) [PubMed](#)
- 3 Warburg, O. (1956) On the origin of cancer cells. *Science* **123**, 309–314 [CrossRef](#) [PubMed](#)
- 4 Bustamante, E., Morris, H.P. and Pedersen, P.L. (1981) Energy metabolism of tumor cells. Requirement for a form of hexokinase with a propensity for mitochondrial binding. *J. Biol. Chem.* **256**, 8699–8704 [PubMed](#)
- 5 Nakashima, R.A., Scott, L.J. and Pedersen, P.L. (1986) The role of mitochondrial hexokinase binding in the abnormal energy metabolism of tumor cell lines. *Ann. N.Y. Acad. Sci.* **488**, 438–450 [CrossRef](#) [PubMed](#)
- 6 Pastorino, J.G., Hoek, J.B. and Shulga, N. (2005) Activation of glycogen synthase kinase 3 β disrupts the binding of hexokinase II to mitochondria by phosphorylating voltage-dependent anion channel and potentiates chemotherapy-induced cytotoxicity. *Cancer Res.* **65**, 10545–10554 [CrossRef](#) [PubMed](#)
- 7 Peng, Q.P., Zhou, J.M., Zhou, Q., Pan, F., Zhong, D.P. and Liang, H.J. (2008) Downregulation of the hexokinase II gene sensitizes human colon cancer cells to 5-fluorouracil. *Chemotherapy* **54**, 357–363 [CrossRef](#) [PubMed](#)
- 8 Shulga, N., Wilson-Smith, R. and Pastorino, J.G. (2009) Hexokinase II detachment from the mitochondria potentiates cisplatin induced cytotoxicity through a caspase-2 dependent mechanism. *Cell Cycle* **8**, 3355–3364 [CrossRef](#) [PubMed](#)
- 9 Liu, H., Li, Y. and Raisch, K.P. (2010) Clotrimazole induces a late G1 cell cycle arrest and sensitizes glioblastoma cells to radiation *in vitro*. *Anticancer Drugs* **21**, 841–849 [CrossRef](#) [PubMed](#)
- 10 Gelb, B.D., Adams, V., Jones, S.N., Griffin, L.D., MacGregor, G.R. and McCabe, E.R. (1992) Targeting of hexokinase 1 to liver and hepatoma mitochondria. *Proc. Natl. Acad. Sci. USA* **89**, 202–206 [CrossRef](#)
- 11 Sui, D. and Wilson, J.E. (1997) Structural determinants for the intracellular localization of the isozymes of mammalian hexokinase: intracellular localization of fusion constructs incorporating structural elements from the hexokinase isozymes and the green fluorescent protein. *Arch. Biochem. Biophys.* **345**, 111–125 [CrossRef](#) [PubMed](#)
- 12 Chiara, F., Castellaro, D., Marin, O., Petronilli, V., Brusilow, W.S., Juhaszova, M., Sollott, S.J., Forte, M., Bernardi, P. and Rasola, A. (2008) Hexokinase II detachment from mitochondria triggers apoptosis through the permeability transition pore independent of voltage-dependent anion channels. *PLoS One* **3**, e1852 [CrossRef](#) [PubMed](#)
- 13 Sun, L., Shukair, S., Naik, T.J., Moazed, F. and Ardehali, H. (2008) Glucose phosphorylation and mitochondrial binding are required for the protective effects of hexokinases I and II. *Mol. Cell Biol.* **28**, 1007–1017 [CrossRef](#) [PubMed](#)
- 14 Rost, B. and Liu, J. (2003) The PredictProtein server. *Nucleic Acids Res.* **31**, 3300–3304 [CrossRef](#) [PubMed](#)
- 15 Yachdav, G., Kloppmann, E., Kajan, L., Hecht, M., Goldberg, T., Hamp, T., Honigschmid, P., Schafferhans, A., Roos, M., Bernhofer, M. et al. (2014) PredictProtein—an open resource for online prediction of protein structural and functional features. *Nucleic Acids Res.* **42**, W337–W343 [CrossRef](#) [PubMed](#)
- 16 Wilson, J.E. (2003) Isozymes of mammalian hexokinase: structure, subcellular localization and metabolic function. *J. Exp. Biol.* **206**, 2049–2057 [CrossRef](#) [PubMed](#)
- 17 Ardehali, H., Yano, Y., Printz, R.L., Koch, S., Whitesell, R.R., May, J.M. and Granner, D.K. (1996) Functional organization of mammalian hexokinase II. Retention of catalytic and regulatory functions in both the NH₂- and COOH-terminal halves. *J. Biol. Chem.* **271**, 1849–1852 [CrossRef](#) [PubMed](#)
- 18 Tsai, H.J. and Wilson, J.E. (1996) Functional organization of mammalian hexokinases: both N- and C-terminal halves of the rat type II isozyme possess catalytic sites. *Arch. Biochem. Biophys.* **329**, 17–23 [CrossRef](#) [PubMed](#)
- 19 Felgner, P.L., Messer, J.L. and Wilson, J.E. (1979) Purification of a hexokinase-binding protein from the outer mitochondrial membrane. *J. Biol. Chem.* **254**, 4946–4949 [PubMed](#)
- 20 Colombini, M. (1979) A candidate for the permeability pathway of the outer mitochondrial membrane. *Nature* **279**, 643–645 [CrossRef](#) [PubMed](#)
- 21 Fiek, C., Benz, R., Roos, N. and Brdiczka, D. (1982) Evidence for identity between the hexokinase-binding protein and the mitochondrial porin in the outer membrane of rat liver mitochondria. *Biochim. Biophys. Acta* **688**, 429–440 [CrossRef](#) [PubMed](#)
- 22 Linden, M., Gellerfors, P. and Nelson, B.D. (1982) Pore protein and the hexokinase-binding protein from the outer membrane of rat liver mitochondria are identical. *FEBS Lett.* **141**, 189–192 [CrossRef](#) [PubMed](#)
- 23 Nakashima, R.A., Mangan, P.S., Colombini, M. and Pedersen, P.L. (1986) Hexokinase receptor complex in hepatoma mitochondria: evidence from N,N'-dicyclohexylcarbodiimide-labeling studies for the involvement of the pore-forming protein VDAC. *Biochemistry* **25**, 1015–1021 [CrossRef](#) [PubMed](#)
- 24 Finney, K.G., Messer, J.L., DeWitt, D.L. and Wilson, J.E. (1984) Monoclonal antibodies against rat brain hexokinase. Effects on catalytic function and binding to the outer mitochondrial membrane. *J. Biol. Chem.* **259**, 8232–8237 [PubMed](#)
- 25 Polakis, P.G. and Wilson, J.E. (1985) An intact hydrophobic N-terminal sequence is critical for binding of rat brain hexokinase to mitochondria. *Arch. Biochem. Biophys.* **236**, 328–337 [CrossRef](#) [PubMed](#)
- 26 Wilson, J.E. and Smith, A.D. (1985) Monoclonal antibodies against rat brain hexokinase. Utilization in epitope mapping studies and establishment of structure–function relationships. *J. Biol. Chem.* **260**, 12838–12843 [PubMed](#)

- 27 Pastorino, J.G., Shulga, N. and Hoek, J.B. (2002) Mitochondrial binding of hexokinase II inhibits Bax-induced cytochrome c release and apoptosis. *J. Biol. Chem.* **277**, 7610–7618 [CrossRef PubMed](#)
- 28 Mehvar, N., Shah, N., Shen, L., Yan, D. and Honzatko, R.B. (2011) Single-residue determinants in the binding of recombinant human brain hexokinase to the mitochondrion. *FASEB J.* **25**, 732–737
- 29 Magnani, M., Crinelli, R., Corsi, D. and Serafini, G. (1992) Intracellular distribution of protein as a determinant for ubiquitination and proteolytic degradation. *Ann. N.Y. Acad. Sci.* **673**, 103–109 [CrossRef PubMed](#)
- 30 Okatsu, K., Iemura, S., Koyano, F., Go, E., Kimura, M., Natsume, T., Tanaka, K. and Matsuda, N. (2012) Mitochondrial hexokinase HKI is a novel substrate of the Parkin ubiquitin ligase. *Biochem. Biophys. Res. Commun.* **428**, 197–202 [CrossRef PubMed](#)
- 31 Nakashima, R.A., Paggi, M.G., Scott, L.J. and Pedersen, R.L. (1988) Purification and characterization of a bindable form of mitochondrial bound hexokinase from the highly glycolytic AS-30D rat hepatoma cell line. *Cancer Res.* **48**, 913–919 [PubMed](#)
- 32 Khalid, M.H., Shibata, S. and Hiura, T. (1999) Effects of clotrimazole on the growth, morphological characteristics, and cisplatin sensitivity of human glioblastoma cells *in vitro*. *J. Neurosurg.* **90**, 918–927 [CrossRef PubMed](#)
- 33 Rabeih, W.M., Zhu, H., Nedyalkova, L., Tempel, W., Wasney, G., Landry, R., Vedadi, M., Arrowsmith, C.H., Edwards, A.M., Sundstrom, M. et al. (2009) Crystal structure of human hexokinase II. University of Toronto Structural Genomics Consortium., doi:10.2210/pdb2nzt/pdb
- 34 Aleshin, A.E., Zeng, C., Bartunik, H.D., Fromm, H.J. and Honzatko, R.B. (1998) Regulation of hexokinase I: crystal structure of recombinant human brain hexokinase complexed with glucose and phosphate. *J. Mol. Biol.* **282**, 345–357 [CrossRef PubMed](#)
- 35 Kuser, P.R., Golubev, A.M. and Polikarpov, I. (1999) Crystallization and preliminary crystal analysis of yeast hexokinase PI and PII. *Acta Crystallogr. D Biol. Crystallogr.* **55**, 2047–2048 [CrossRef](#)
- 36 Kuser, P.R., Krauchenco, S., Antunes, O.A. and Polikarpov, I. (2000) The high resolution crystal structure of yeast hexokinase PII with the correct primary sequence provides new insights into its mechanism of action. *J. Biol. Chem.* **275**, 20814–20821 [CrossRef PubMed](#)
- 37 Smeele, K.M., Southworth, R., Wu, R., Xie, C., Nederlof, R., Warley, A., Nelson, J.K., van Horssen, P., van den Wijngaard, J.P., Heikkinen, S. et al. (2011) Disruption of hexokinase II-mitochondrial binding blocks ischemic preconditioning and causes rapid cardiac necrosis. *Circ. Res.* **108**, 1165–1169 [CrossRef PubMed](#)

Received 18 February 2015/20 April 2015; accepted 23 April 2015

Published as Immediate Publication 24 April 2015, doi 10.1042/BSR20150047
