

Title: Trastuzumab Deruxtecan in HER2-Positive Metastatic Breast Cancer Patients with Brain Metastases: A DESTINY-Breast01 Subgroup Analysis

Authors: Guy Jerusalem¹, Yeon Hee Park², Sara A. Hurvitz³, Shanu Modi⁴, Fabrice Andre⁵, Ian E. Krop⁶, Xavier González Farré⁷, Benoit You⁸, Cristina Saura⁹, Sung-Bae Kim¹⁰, Cynthia R. Osborne^{11,12}, Rashmi K. Murthy¹³, Lorenzo Gianni¹⁴, Toshimi Takano¹⁵, Yali Liu¹⁶, Jillian Cathcart¹⁶, Caleb Lee¹⁶, Christophe Perrin¹⁷

Affiliations: ¹Centre Hospitalier Universitaire du Sart Tilman Liège and Liège University, Department of Medical Oncology, Breast Clinic, Liège, Belgium. ²Division of Hematology-Oncology, Department of Medicine, Samsung Medical Center, Seoul, Republic of Korea. ³UCLA, Division of Hematology-Oncology, UCLA Jonsson Comprehensive Cancer Center, Los Angeles, California. ⁴Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, New York. ⁵Gustave Roussy, Department of Immunology, Université Paris-Sud, Villejuif, France. ⁶Department of Medical Oncology, Yale Cancer Center, New Haven, Connecticut. ⁷Instituto Oncológico Dr Rosell, Hospital General De Catalunya, SOLTI, Institut Oncològic, Barcelona, Spain. ⁸Medical Oncology Department, Institut de Cancérologie des Hospices Civils de Lyon, CITOHL, Department of Medical Oncology, Université Claude Bernard Lyon 1, Lyon, France. ⁹Vall d'Hebron University Hospital, Breast Cancer Unit, Medical Oncology Service and Vall d'Hebron Institute of Oncology (VHIO), Barcelona, Spain. ¹⁰Department of Oncology, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Republic of Korea. ¹¹US Oncology Research, McKesson Specialty Health, The

Woodlands, Texas. ¹²Texas Oncology, Baylor-Sammons Cancer Center, Medical Services, Dallas, Texas. ¹³Breast Medical Oncology, The University of Texas MD Anderson Cancer Center, Houston, Texas. ¹⁴Department of Oncology, Infermi Hospital, AUSL della Romagna, Rimini, Italy. ¹⁵Breast Medical Oncology Department, The Cancer Institute Hospital of JFCR, Tokyo, Japan, Tokyo, Japan. ¹⁶Daiichi Sankyo, Basking Ridge, New Jersey. ¹⁷Department of Oncology, Centre Eugène Marquis, Rennes, France.

Corresponding author:

Guy Jerusalem, MD

Department of Medical Oncology

Centre Hospitalier Universitaire du Sart-Tilman

Avenue de l'Hôpital, 1

4000 Liège, Belgium

Belgium

g.jerusalem@chu.ulg.ac.be

Phone: +32 (4) 3668414

Fax: +32 43667688

Running title: T-DXd in HER2+ Breast Cancer with Brain Metastases

Keywords: trastuzumab deruxtecan; T-DXd1; breast cancer; HER2; brain metastasis

Additional information

Financial support: Daiichi Sankyo and AstraZeneca

Conflict of interest:

Guy Jerusalem reports institutional grants from Novartis, Roche, and Pfizer; consulting fees from Novartis, Amgen, Roche, Pfizer, Bristol Myers Squibb, Lilly, AstraZeneca, Daiichi Sankyo, and AbbVie; honoraria from Novartis, Roche, Amgen, Pfizer, Bristol Myers Squibb, Lilly, and AstraZeneca; travel support from Novartis, Roche, Pfizer, Lilly, Amgen, Bristol Myers Squibb, and AstraZeneca; advisory board fees from Novartis, Roche, Amgen, Pfizer, Bristol Myers Squibb, Lilly, AstraZeneca, Daiichi Sankyo, and AbbVie; and medical writing fees from Novartis, Roche, Lilly, Amgen, Bristol Myers Squibb, AstraZeneca, Medimmune, Merck, and Daiichi Sankyo.

Yeon Hee Park reports grants from AstraZeneca, MSD, Sanofi, DongA-ST, Roche, Pfizer, and Alteogen; consulting fees from AstraZeneca, MSD, Roche, Novartis, Pfizer, Daiichi Sankyo, Eli Lilly, and Eisai; honoraria from MSD, Novartis, Samsung BioEpi, Celltrion, Pfizer, Roche, Eli Lilly, and Eisai; payment testimony from AstraZeneca, Pfizer, Roche, and Novartis; advisory board fees from AstraZeneca, Daiichi Sankyo, Pfizer, and Roche; and receipt of equipment, materials, drugs, medical writing, gifts, or other services from Pfizer and MSD.

Sara A. Hurvitz reports contracted research grants from Ambrx, Amgen, Astra Zeneca, Arvinas, Bayer, Daiichi Sankyo, Dignitana, Genentech/Roche, Gilead, GSK, Immunomedics, Lilly, MacroGenics, Novartis, Pfizer, OBI Pharma, Phoenix Molecular Designs, Ltd. Pieris, PUMA, Radius, Samumed, Sanofi, Seattle Genetics, and

Zymeworks; stock options from ROMtech, Ideal Implant, and NKMax; and travel grants from Lilly.

Shanu Modi reports consulting fees from Daiichi Sankyo, AstraZeneca, Genentech, and Macrogenics; honoraria from Daiichi Sankyo, AstraZeneca, Genentech, and Seattle Genetics; and advisory board fees from Macrogenics.

Fabrice Andre reports institutional grants from Novartis, Pfizer, AstraZeneca, Eli Lilly, Daiichi Sankyo, and Roche.

Ian E. Krop reports institutional grants from Genentech/Roche, Pfizer, and Macrogenics; consulting fees from Bristol Myers Squibb, Daiichi Sankyo, Macrogenics, Context Therapeutics, Taiho Oncology, Genetech/Roche, Seagen, and AstraZeneca; and advisory board fees from Novartis and Merck.

Xavier González Farré reports honoraria from Pierre Fabre and Roche and travel support from Roche.

Benoit You has nothing to disclose.

Cristina Saura reports institutional grants from AstraZeneca, Daiichi Sankyo, Eli Lilly and Company, Genentech, Immunomedics, Macrogenics, Merck, Sharp and Dohme España S.A., Novartis, Pfizer, Piquor Therapeutics, Puma, Roche, Synthon, and Zenith Pharma; honoraria from AstraZeneca, Daiichi Sankyo, Eisai, Exact Sciences, Exeter Pharma, F. Hoffmann-La Roche Ltd, MediTech, Merck Sharp & Dohme, Novartis, Pfizer, Philips, Pierre Fabre, Puma, Roche Farma, Sanofi-Aventis, Seagen and Zymeworks; and travel support from AstraZeneca, Daiichi Sankyo, Eisai, Novartis, Pfizer, Pierre Fabre, Puma, Roche Farma, and Seagen.

Sung-Bae Kim reports institutional grants from Novartis, Sanofi-Aventis, DongKook Pharm Co; consulting fees from Novartis, AstraZeneca, Lilly, Dae Hwa Pharmaceutical Co, Ltd, IUS Abxis, Daiichi Sankyo, and Beigene; advisory board fees from Novartis, AstraZeneca, Lilly, Dae Hwa Pharmaceutical Co, Ltd, IUS Abxis, Daiich Sankyo, and Beigene; leadership or fiduciary role on ESMO Breast 2021-2022; and stock options from Genopeak and Neogene TC.

Cynthia R. Osborne reports advisory board fees from AstraZeneca.

Rashmi K. Murthy reports institutional grants from Seagen, Genentech, Pfizer, Daiichi Sankyo, AstraZeneca, and EMD-Serono and consulting fees from Seagen, Genentech, Novartis, Pfizer, Astra Zeneca, Puma, and Sanofi.

Lorenzo Gianni reports support for the present manuscript from Daiichi Sankyo; travel support from Daiichi Sankyo, Novartis, and Pfizer; and advisory board fees from AstraZeneca.

Toshimi Takano reports institutional grants from Daiichi Sankyo, Chugai, Ono, MSD, and Eisai and honoraria from Daiichi Sankyo, Chugai, Eisai, Kyowa Kirin, Pfizer, Eli Lilly, and Celltrion Healthcare.

Yali Lui, Jillian Cathcart, and Caleb Lee are employees of Daiichi Sankyo.

Journal: *Cancer Discovery*

Current Counts/Journal Guidelines:

Article type: Research Brief

Abstract: 148/150 words

Statement of significance: 38/50 words

Body of text: 3605/3500 words

Tables/figures: 2 tables and 2 figures/4 maximum

References: 28/25

Supplementary appendix: Yes

Abstract (149/150)

DESTINY-Breast01 (NCT03248492) evaluated trastuzumab deruxtecan (T-DXd; DS-8201) in patients with heavily pretreated HER2-positive metastatic breast cancer (mBC). We present a subgroup of 24 patients with a history of treated brain metastases (BMs), a population with limited treatment options. In patients with BMs, confirmed objective response rate (cORR) was 58.3% (95% CI, 36.6%-77.9%) and median progression-free survival (mPFS) was 18.1 months (95% CI, 6.7-18.1 months). In patients without BMs ($n = 160$), cORR was 61.3% and mPFS was 16.4 months. Eight patients (47.1%) experienced a best overall intracranial response of partial response or complete response. Seven patients (41.2%) had a best percentage change in brain lesion diameter from baseline consistent with stable disease. Two patients (8.3%) with BMs and 2 (1.3%) without BMs experienced progression in the brain. The safety profile of T-DXd was consistent with previous studies. The durable clinical activity of T-DXd in this population warrants further investigation.

Significance (38/50)

Advances in treating HER2-positive metastatic breast cancer have greatly improved patient outcomes, but intracranial progression remains an important risk for which few therapeutic options are currently available. T-DXd demonstrated durable efficacy in patients with stable, treated brain metastases.

Introduction

Recent advances in treatment have improved the overall survival (OS) of patients with HER2-positive metastatic breast cancer (mBC) (1-4). However, approximately 22% to 50% of patients with advanced disease will develop brain metastases (BMs) (1-4). This relatively high incidence may be a result of improved OS after the initial breast cancer diagnosis due to better control of extracranial disease by HER2-targeted treatments, a biological predilection for HER2-positive breast cancer to metastasize to the brain, and greater detection because of more advanced imaging techniques (5). Patients with mBC with BMs typically have a poorer prognosis and a decreased quality of life compared with patients without BMs (1). Initial treatment of BMs in these patients typically involves locally directed therapies, including resection, together with postoperative radiation, stereotactic radiosurgery (SRS), and/or whole brain radiotherapy (6). Despite the prevalence of central nervous system metastases in patients with HER2-positive mBC, clinical trials of HER2-directed therapies have often excluded patients with progressive BMs (7).

For the underlying systemic disease, HER2-directed treatment is administered according to the locally prescribed regimens for HER2-positive mBC (i.e., trastuzumab plus pertuzumab and chemotherapy in the first-line setting) (1). However, the limited ability of some systemic treatments to cross the blood-brain barrier (BBB) and the unique brain tumor microenvironment, characterized by distinct cell types and metabolic limitations, may contribute to poor control of such lesions (1, 8). Thus, additional treatment options are needed to target the BMs frequently observed in patients with HER2-positive mBC.

The tumor microenvironment may cause enhanced vascular permeability, allowing for the extravasation of large molecules, such as antibody-drug conjugates (ADCs), through the BBB. In a preclinical study of the HER2-targeting ADC trastuzumab emtansine (T-DM1), it was proposed the enhanced vascular permeability of the BBB may lead to adequate concentrations of T-DM1 in BMs to elicit a clinical benefit (9). Trastuzumab deruxtecan (T-DXd) is an ADC that comprises a humanized, monoclonal, anti-HER2 antibody with a topoisomerase I inhibitor payload attached via a plasma-stable, tumor-selective cleavable, tetrapeptide-based linker (10, 11). Based on the DESTINY-Breast01 (Trial Registration Number: NCT03248492) and DESTINY-Breast03 (NCT03529110) studies (12, 13), T-DXd was approved for the treatment of patients with unresectable or metastatic HER2-positive breast cancer after a previous (United States) or ≥ 2 previous (various countries worldwide) anti-HER2-based regimens (14, 15). In mature results from DESTINY-Breast01, T-DXd demonstrated a confirmed objective response rate (cORR) by independent central review (ICR) of 62.0% (114/184) and a median progression-free survival (mPFS) of 19.4 months in patients with HER2-positive unresectable or mBC previously treated with T-DM1 at a data cutoff of March 26, 2021 (12, 16). In the phase 3 DESTINY-Breast03 trial, which included patients with HER2-positive mBC previously treated with trastuzumab and a taxane, mPFS was not yet reached (95% confidence interval [CI], 18.5-nonevaluable [NE]) in the T-DXd arm versus 6.8 months (95% CI, 5.6-8.2) in the T-DM1 arm (hazard ratio [HR], 0.28 [95% CI, 0.22-0.37]) (13).

Here I describe a subgroup analysis from DESTINY-Breast01, reporting safety and efficacy of T-DXd 5.4 mg/kg in patients with a history of BMs and with stable BMs visible on baseline imaging.

Results

Patient Characteristics

A total of 184 patients received T-DXd 5.4 mg/kg (12), 24 of whom (13.0%) had a history of BMs. Demographic and clinical characteristics of patients in the BM subgroup were generally similar to those of patients in the non-BM subgroup (**Supplementary Table S1**), although a greater proportion of patients in the BM subgroup had an Eastern Cooperative Oncology Group performance status of 0 (62.5% vs. 54.4%) or had hormone receptor–negative (58.3% vs. 43.1%) breast cancer compared with patients in the non-BM subgroup. A higher proportion of patients in the BM subgroup previously received a HER2 tyrosine kinase inhibitor (TKI; 62.5% vs. 48.8%) or radiotherapy (any location; 83.3% vs. 68.8%) compared with patients in the non-BM subgroup. In the BM subgroup, 58.3% (14/24) of patients had radiotherapy of their central nervous system lesion (including SRS and whole-brain radiation), 12.5% (3/24) of patients had radiotherapy plus surgery, 4.2% (1/24) of patients had radiotherapy plus surgery and capecitabine, 4.2% (1/24) of patients had surgery only, defined as craniotomy, metastasectomy, or resection/removal of the brain lesion, and 20.8% (5/24) of patients had no reported prior CNS therapy. Patients in both subgroups were heavily pretreated, with a median of 6 prior therapies in the metastatic setting.

At the time of data cutoff (August 1, 2019), 45.8% (11/24) of patients in the BM subgroup and 42.5% (68/160) of patients in the non-BM subgroup were still receiving T-DXd. Patients in both subgroups primarily discontinued treatment because of progressive disease (PD; 25.0% in the BM subgroup; 29.4% in the non-BM subgroup) and adverse events (12.5%; 15.6%). Median treatment duration was 11.0 months (range, 0.7-20.2) for the BM subgroup and 9.9 months (range, 0.7-20.5) for the non-BM subgroup.

Efficacy

The systemic efficacy of T-DXd 5.4 mg/kg in the BM and non-BM subgroups was consistent with the efficacy observed in patients in the overall study population (**Table 1**). The primary endpoint, cORR by ICR, was 58.3% (95% CI, 36.6-77.9) in the BM subgroup, with 1 (4.2%) of these patients achieving a complete response (CR) and 13 (54.2%) a partial response (PR). Of the 10 remaining patients in the BM subgroup, 8 (33.3%) had stable disease (SD), resulting in a disease control rate (DCR; CR + PR + SD) of 91.7% (22/24). In the non-BM subgroup, cORR was 61.3% (95% CI, 53.2-68.8), with 10 patients (6.3%) achieving CR and 88 (55.0%) a PR. SD was achieved by 59 patients (36.9%) and DCR was 98.1% (157/160). The median follow-up was 11.0 months (range, 0.7–19.6) for patients with BMs and 11.1 months (range, 2.4-19.9) for patients in the non-BM subgroup. The median time to response with confirmation was 2.8 months (95% CI, 1.3-4.1) for the BM subgroup and 1.5 months (95% CI, 1.4-2.6) for the non-BM subgroup. mPFS was 18.1 months (95% CI, 6.7-18.1) for the BM subgroup and 16.4 months (95% CI, 12.7-NE) for the non-BM subgroup (**Fig. 1, Table 1**).

Of the 24 patients in the BM subgroup, 17 had reported tumor measurements for brain lesions at baseline (**Table 2**). Fifteen of 17 patients had data available to evaluate responses in the brain. Of the patients with available data, 14 had completed radiotherapy ≥ 60 days before treatment and 1 had not received any prior radiotherapy. Per investigator assessment, 7 of 17 patients (41.2%) had a best percentage change in brain lesion diameters consistent with a response of PR ($\geq 30\%$ decrease) in the brain, and 7 of 17 patients (41.2%) had a best percentage change in brain lesion diameters consistent with SD (**Fig. 2A**). The reductions in brain lesions were sustained over time (**Fig. 2B**).

In total, 48 patients experienced systemic PD, including 8 patients with a history of BMs. Sites of progression were generally similar between patients in the BM and non-BM subgroups (**Supplementary Table S2**), although the rate of progression in the brain was greater in patients with a history of BMs relative to those without (8.3% and 1.3%, respectively) at time of data cutoff. Overall, 4 patients, 2 of whom had a history of BMs, had disease progression occurring in the brain. These progression events occurred earlier in the 2 BM subgroup patients (days 75 and 85) than in the 2 non-BM patients (days 323 and 498).

Safety

All patients in the BM subgroup and 99.4% of non-BM patients experienced ≥ 1 treatment-emergent adverse event (TEAE) (**Supplementary Table S3**). As in the overall study population, any-grade TEAEs in the BM subgroup were predominantly gastrointestinal or hematologic. The percentage of grade ≥ 3 TEAEs was 54.2% in the

BM subgroup and 57.5% in the non-BM subgroup. In the BM subgroup, the most prevalent grade ≥ 3 TEAEs were decreased neutrophil count (25.0%), decreased lymphocyte count (16.7%), fatigue (12.5%), and diarrhea (8.3%) (**Supplementary Table S4**). Drug-related TEAEs associated with discontinuation occurred in 2 patients (8.3%) with a history of BMs (grade 1 and 2 pneumonitis) and 25 (15.6%) in those without. Pneumonitis ($n = 11$) and interstitial lung disease (ILD; $n = 5$) were the predominant causes for discontinuation among the non-BM patients. There were 9 TEAEs associated with deaths, 2 of which were in the BM subgroup and resulted from disease progression and respiratory failure (each, $n = 1$; adjudicated as drug-related ILD/pneumonitis per an independent ILD/pneumonitis adjudication committee). TEAEs in the non-BM subgroup associated with death included 1 case each of acute respiratory failure, general physical health deterioration, lymphangitis, pneumonia, pneumonitis, and shock hemorrhagic. One patient had 2 TEAEs associated with death (acute kidney injury and acute hepatic failure) (**Supplementary Table S3**).

ILD/pneumonitis, an important risk associated with T-DXd, adjudicated by an independent committee as related to study treatment, occurred in 4 patients (16.7%) in the BM subgroup (grade 1, $n = 3$ [12.5%]; grade 5, $n = 1$ [4.2%]) and 21 patients (13.1%) in the non-BM subgroup (grade 1, $n = 6$ [3.8%]; grade 2, $n = 10$ [6.3%]; grade 3, $n = 2$ [1.3%], grade 5, $n = 3$ [1.9%]). There were 25 adjudicated ILD/pneumonitis cases in the overall study population (13.6%; grade 1/2, $n = 19$ [10.3%]; grade 3, $n = 2$ [1.1%]; grade 4, $n = 0$; grade 5, $n = 4$ [2.2%]) (12).

Patient Case Study

One patient treated with T-DXd 5.4 mg/kg every 3 weeks in the DESTINY-Breast01 study who had baseline BMs was a 48-year-old patient with HER2-positive (immunohistochemistry [IHC] 3+)/hormone receptor–negative mBC. Before enrollment, the patient had received 16 lines of treatment, including T-DM1, pertuzumab, trastuzumab, and lapatinib. The patient also underwent whole-brain radiotherapy 5 years before study inclusion and SRS 3 years before study inclusion to treat brain lesions. Lesions were observed at the patient’s baseline scan in the brain, lymph nodes, retroperitoneum, lung, pancreas, bone, and axillary lymph node.

At baseline, the diameter of the brain lesion was 10.6 mm and was reduced by 35% at the 6-week scan. The patient’s overall status improved, allowing for reduction in pain medication. PR was achieved in all target lesions and non-CR/non-PD response in the nontarget lesions.

By the subsequent scan at 12 weeks, the brain lesion measured 4.6 mm—a 57% reduction from baseline. Although a PR continued to be observed for all target lesions and a non-CR/non-PD response was observed for all nontarget lesions, new lesions were observed in the chest wall. The overall response at 12 weeks was PD, leading to the patient’s discontinuation from study treatment per the protocol.

Discussion

Among DESTINY-Breast01 patients with HER2-positive mBC, T-DXd 5.4 mg/kg demonstrated durable activity in patients with and without a history of BMs, similar to that observed in the overall population. The cORR and mPFS in the BM and non-BM subgroups were similar to the overall population. The cORR was 58.3% for patients with

BMs and 61.3% for patients without BMs, and mPFS was 18.1 months and 16.4 months, respectively. Additionally, the safety profile of T-DXd in patients with and without a history of BMs was similar to the overall population. Because additional treatment options are needed specifically for patients with HER2-positive mBC and BMs, these data are encouraging and warrant further investigation.

An important concern in treating BMs has been that large, systemically administered molecules may not penetrate the BBB to achieve an effective concentration in the tumor bed. Because of their smaller size, chemical inhibitors, such as TKIs, have been considered to have better penetration of the BBB than larger antibody-based treatments (e.g., trastuzumab, T-DM1, and T-DXd) (4). However, clinical and preclinical studies have detected accumulation of trastuzumab and T-DM1 in brain lesions, possibly as the result of disruption of the BBB from prior local surgery and/or radiation, or because of a blood-tumor barrier with varying permeability, or an altered blood-tumor barrier after penetration of the BBB by BMs (5, 9). Such evidence suggests that the brain's status as a "sanctuary site" in patients with HER2-positive mBC is not completely explained by lack of treatment accessibility. The immune privilege status of the brain may also restrict trastuzumab-mediated antibody-dependent cell-mediated cytotoxicity. Consequently, ADCs such as T-DXd or T-DM1 may be better suited than monoclonal antibodies for treating BMs since they can cause tumor cell death via delivery of a cytotoxic drug. The efficacy of anti-HER2 agents in treating BMs has recently been reported. T-DM1 was examined in the second-line or later setting in a subgroup analysis of the single-arm KAMILLA trial in patients with HER2-positive mBC who had treated or asymptomatic, untreated BMs at baseline (17). For this subgroup, the best overall

response was 21.4% (27/126; [95% CI, 14.6-29.6]; CR, $n = 3$; PR, $n = 24$), and a $\geq 30\%$ reduction in the sum of the largest diameters of target brain lesions was observed in 42.9% of patients (54/126). The mPFS was 5.5 months, compared with 7.7 months in patients without baseline BMs (17). The HER2CLIMB trial was one of the first trials to enroll a large proportion of patients with BMs, including patients with previously untreated, treated stable, or treated and progressing BMs (18). Based on HER2CLIMB, tucatinib in combination with trastuzumab and capecitabine was recently approved for patients with HER2-positive mBC, including those with BMs, after ≥ 1 prior (United States) or ≥ 2 prior (Europe) lines of HER2-targeted therapy (18, 19). Patients with BMs at baseline or a history of BMs were included in this trial and achieved a mPFS of 7.6 months with tucatinib combination therapy (19). The ORR for all patients in HER2CLIMB ($N = 511$) was 40.6% (95% CI, 35.3-46.0), and the intracranial ORR for patients with active BMs and measurable intracranial disease at baseline ($n = 75$) was 47.3% (95% CI, 33.7-61.2) (18, 19). In the overall population, median OS was 24.7 months (95% CI, 21.6-28.9) for tucatinib plus trastuzumab and capecitabine versus 19.2 months (95% CI, 16.4-21.4) for placebo plus trastuzumab and capecitabine (HR, 0.73 [95% CI, 0.59-0.90]; $P = 0.004$) (20).

While cross-trial comparisons are difficult to interpret, the response rate and mPFS for patients with a history of BMs in DESTINY-Breast01 were numerically higher compared with patients with BMs at baseline from the KAMILLA and HER2CLIMB trials (cORR, 58.3%; mPFS, 18.1 months) but there were differences between the study populations. Patients in the BM subgroup in DESTINY-Breast01 had stable BMs and required a washout period of ≥ 60 days before randomization if they had any history of radiation,

surgery, or other therapy, including steroids or anticonvulsants to control symptoms from BMs. The number of patients with BMs was also relatively low (12). In the KAMILLA trial, patients with BMs were included if they had untreated, asymptomatic BMs or controlled brain disease treated with radiotherapy >14 days before enrollment (17). In the HER2CLIMB trial, patients with BMs were included unless they needed immediate local intervention. HER2CLIMB also enrolled 291 patients with treated and stable, treated and progressing, and untreated BMs. Additionally, patients with untreated BMs >2 cm in diameter could be enrolled in HER2CLIMB provided they received approval from the medical monitor (19).

Intracranial efficacy of T-DXd has also been reported in preclinical and clinical studies. Kabraji et al. reported prolonged survival with T-DXd in HER2-positive and HER2-low breast cancer BM-derived murine models compared with control (21). Additionally, a clinical benefit rate of 75% (12/16) was observed in a retrospective cohort of breast cancer patients with BMs (21). In the phase 2 TUXEDO-1 trial (NCT04752059), T-DXd demonstrated efficacy and safety in patients with HER2-positive breast cancer with newly diagnosed BMs or BMs with radiologic progression after prior local therapy. Intracranial response was observed in 11/15 patients (73.3%) (22). In the phase 3 DESTINY-Breast03, 23/36 (63.9%) patients with stable BMs at baseline experienced intracranial CR or PR (23). In the ongoing phase 2 DEBBRAH trial (NCT04420598), T-DXd demonstrated preliminary efficacy in patients with HER2-positive advanced breast cancer with stable, asymptomatic untreated, or progressing BMs after radiotherapy and/or surgery. Intracranial ORR was reported in 2 patients (50.0% [95% CI, 6.7-93.2]) with asymptomatic untreated BMs and 4 patients (44.4% [95% CI, 13.7-

78.8]) with treated and progressing BMs (24). The ongoing DESTINY-Breast07 (NCT04538742), DESTINY-Breast09 (NCT04784715), DESTINY-Breast12 (NCT04739761), and HER2CLIMB-04 (NCT04539938) trials will continue to assess the efficacy of T-DXd in patients with HER2-expressing breast cancer and BMs, including active BMs.

Patients with HER2-negative mBC comprise approximately 85% of the mBC population, and approximately 60% of tumors from patients with HER2-negative mBC express low levels of HER2 (25, 26). T-DXd was approved in the United States for the treatment of patients with unresectable or metastatic HER2-low (IHC 1+ or IHC 2+/in situ hybridization-) breast cancer after prior chemotherapy in the metastatic setting based on results from DESTINY-Breast04 (NCT03734029) (27). Therefore, the data reported here suggest that a large subset of patients with mBC and BMs may experience clinical benefit with T-DXd. The ongoing DESTINY-Breast06 (NCT04494425) and DESTINY-Breast08 (NCT04556773) studies, and a cohort from DEBBRAH are continuing to assess HER2-low patients with treated/stable BMs.

DESTINY-Breast01 did not include patients with active BMs, and 19 of 24 patients in the BM subgroup had received prior surgery or radiation to the brain. BMs were to be listed as nontarget lesions; although BMs were monitored by imaging every 6 weeks (Q6W), diameter measurements were not required. In addition, our subgroup analysis included only 24 patients, of whom 17 had brain lesion measurements available for analysis. Tumor measurements were obtained retrospectively by contacting the treating investigators, and the date of data cutoff could not be independently verified. However, the responses noted in the brain generally occurred within the first 2 on-treatment

scans, and so the impact of data cutoff on ORR was likely minimal. Additionally, per Response Evaluation Criteria in Solid Tumors criteria, tumor lesions located in a previously irradiated area are usually not considered target lesions (28). Thus, proper response assessments were not possible for the patients in the BMs subgroup with prior irradiation.

In conclusion, T-DXd demonstrated strong clinical activity in both the overall population of patients with HER2-positive mBC and the subgroup of those with a history of BMs in DESTINY-Breast01. These positive findings lay the groundwork for further investigation of T-DXd in this substantial patient subset, including those with active BMs, for which treatment options remain limited.

Methods

Study Design and Patients

DESTINY-Breast01 is a 2-part, multicenter, open-label (unblinded), phase 2 trial of T-DXd in adult patients with unresectable or metastatic HER2-positive breast cancer previously treated with T-DM1. The overall design, study protocol, and primary results of this study have been previously published (12).

Briefly, patients were adults (≥ 20 years of age in Japan and Korea; ≥ 18 years of age in all other study sites) with unresectable or metastatic, HER2-positive (IHC 3+ or in situ hybridization+ as centrally confirmed on archival tissue) breast cancer that progressed on T-DM1 or who had discontinued T-DM1 for reasons other than PD. Patients previously treated with HER2-targeted TKIs or capecitabine were eligible. Additionally, patients with baseline BMs that were treated, asymptomatic, or did not require therapy

to control symptoms were eligible. All treatment to control symptoms, including radiation, surgery, or other therapy (including steroids or anticonvulsants), had to be completed >60 days before randomization. Radiotherapy had to have been completed ≥60 days before treatment initiation to decrease the likelihood that prior radiotherapy contributed to tumor evaluation on study. The planned number of patients with BMs or a history of BMs to be included in the study was approximately 20% of the ≈150 patients expected to receive the recommended part 2 dose (T-DXd 5.4 mg/kg every 3 weeks); additional patients with any current or past history of BMs were excluded. Tumors were assessed by imaging Q6W (±7 days); a collection of magnetic resonance imaging scans for central analysis of the brain Q6W was only required for patients with BMs at baseline. Brain lesion measurements were assessed per investigator, but were considered nontarget lesions, and thus collection of diameter measurements was not required. Patients with BMs had to have a target lesion elsewhere, which was measured for the analysis of sum of the longest diameters. Patients were also excluded if they had current or suspected noninfectious ILD/pneumonitis or a history of ILD/pneumonitis that required treatment with glucocorticoids. Complete eligibility criteria can be found in the previously published analysis (12).

Study Oversight

Daiichi Sankyo and AstraZeneca collaborated on this study, which was sponsored and designed by Daiichi Sankyo and approved by the institutional review board at each participating site. All patients provided written informed consent. Data were analyzed and interpreted by the sponsor and authors. All authors reviewed the manuscript and

confirmed the accuracy of the data, its analysis, and adherence to the protocol, available at cancerdiscovery.aacrjournals.org.

Endpoints

The data cutoff for this analysis was August 1, 2019. The primary endpoint was cORR (CR + PR) per ICR using Response Evaluation Criteria in Solid Tumors, version 1.1. Secondary endpoints were investigator assessed and included duration of response, progression-free survival (PFS), OS, DCR (CR + PR + SD), safety, and best percentage change in the sum of the longest diameters of measurable tumors. The primary and secondary endpoints were determined for all patients who received the recommended dose of T-DXd 5.4 mg/kg (previously described (12)) and are presented here for patients with and without a history of BMs.

Safety

TEAEs were categorized in accordance with the Medical Dictionary for Regulatory Activities, version 20.1, and graded using the National Cancer Institute Common Terminology Criteria for Adverse Events, version 4.03.

An independent adjudication committee was established to evaluate all potential incidents of ILD/pneumonitis (12).

Statistical Analysis

The statistical analyses for evaluating the endpoints in all patients who received the recommended part 2 dose were previously described (12) and were also applied to the

BM subgroup. Briefly, the Clopper–Pearson method was used to calculate the 2-sided 95% CI for the response rate and the Kaplan–Meier method was used to estimate the distribution of time-to-event end points of duration of response and PFS. The corresponding 2-sided 95% CI for the median time-to-event endpoints were calculated with the Brookmeyer and Crowley methods.

Data Availability Statement

Anonymized individual participant data and applicable supporting clinical trial documents may be available upon request at (<https://vivli.org>). In cases where clinical trial data and supporting documents are provided pursuant to our company policies and procedures, Daiichi Sankyo Companies will continue to protect the privacy of the company and our clinical study subjects. Details on data-sharing criteria and the procedure for requesting access can be found at this web address:

<https://vivli.org/ourmember/daiichi-sankyo>. Data will be provided upon request.

Acknowledgements

Editorial assistance was provided by Jenna M. Gaska, PhD (ArticulateScience, LLC), Jill Seabrook, PhD, and Laura Halvorson, PhD (ApotheCom), and financially supported by Daiichi Sankyo.

References

1. Hurvitz SA, O'Shaughnessy J, Mason G, Yardley DA, Jahanzeb M, Brufsky A, et al. Central nervous system metastasis in patients with HER2-positive metastatic breast cancer: patient characteristics, treatment, and survival from SystHERs. *Clin Cancer Res* 2019;25(8):2433-41.
2. Olson EM, Najita JS, Sohl J, Arnaout A, Burstein HJ, Winer EP, et al. Clinical outcomes and treatment practice patterns of patients with HER2-positive metastatic breast cancer in the post-trastuzumab era. *Breast* 2013;22(4):525-31.
3. Bendell JC, Domchek SM, Burstein HJ, Harris L, Younger J, Kuter I, et al. Central nervous system metastases in women who receive trastuzumab-based therapy for metastatic breast carcinoma. *Cancer* 2003;97(12):2972-7.
4. Clayton AJ, Danson S, Jolly S, Ryder WD, Burt PA, Stewart AL, et al. Incidence of cerebral metastases in patients treated with trastuzumab for metastatic breast cancer. *Br J Cancer* 2004;91(4):639-43.
5. Venur VA, Leone JP. Targeted therapies for brain metastases from breast cancer. *Int J Mol Sci* 2016;17(9):1543.
6. Ramakrishna N, Anders CK, Lin NU, Morikawa A, Temin S, Chandarlapaty S, et al. Management of Advanced Human Epidermal Growth Factor Receptor 2-Positive Breast Cancer and Brain Metastases: ASCO Guideline Update. *J Clin Oncol* 2022;40(23):2636-55.
7. Costa R, Gill N, Rademaker AW, Carneiro BA, Chae YK, Kumthekar P, et al. Systematic analysis of early phase clinical studies for patients with breast cancer: inclusion of patients with brain metastasis. *Cancer Treat Rev* 2017;55:10-5.

8. Hosonaga M, Saya H, Arima Y. Molecular and cellular mechanisms underlying brain metastasis of breast cancer. *Cancer Metastasis Rev* 2020;39(3):711-20.
9. Askoxylakis V, Ferraro GB, Kodack DP, Badeaux M, Shankaraiah RC, Seano G, et al. Preclinical efficacy of ado-trastuzumab emtansine in the brain microenvironment. *J Natl Cancer Inst* 2016;108(2):djv313.
10. Ogitani Y, Aida T, Hagihara K, Yamaguchi J, Ishii C, Harada N, et al. DS-8201a, a novel HER2-targeting ADC with a novel DNA topoisomerase I inhibitor, demonstrates a promising antitumor efficacy with differentiation from T-DM1. *Clin Cancer Res* 2016;22(20):5097-108.
11. Lüftner D, Peipp M. New therapeutic strategies in advanced nonoperable or metastatic HER2-positive breast cancer. *Geburtshilfe Frauenheilkd* 2021;81(6):666-78.
12. Modi S, Saura C, Yamashita T, Park YH, Kim SB, Tamura K, et al. Trastuzumab deruxtecan in previously treated HER2-positive breast cancer. *N Engl J Med* 2020;382(7):610-21.
13. Cortés J, Kim SB, Chung WP, Im SA, Park YH, Hegg R, et al. Trastuzumab deruxtecan versus trastuzumab emtansine for breast cancer. *N Engl J Med* 2022;386(12):1143-54.
14. Enhertu 100 mg powder for concentrate for solution for infusion. Summary of product characteristics. Daiichi Sankyo Europe GmbH: Munich, Germany; 02/2021.

15. Daiichi Sankyo I. Enhertu[®] (fam-trastuzumab deruxtecan-nxki) for injection, for intravenous use. Prescribing information. Daiichi Sankyo, Inc.: Basking Ridge, NJ; 2022. p 14.
16. Saura Manich C, Modi S, Krop I, Park YH, Kim SB, Tamura K, et al. 279P Trastuzumab deruxtecan (T-DXd) in patients with HER2-positive metastatic breast cancer (MBC): Updated survival results from a phase II trial (DESTINY-Breast01). *Ann Oncol* 2021;32:S485-S6.
17. Montemurro F, Delaloge S, Barrios CH, Wuerstlein R, Anton A, Brain E, et al. Trastuzumab emtansine (T-DM1) in patients with HER2-positive metastatic breast cancer and brain metastases: exploratory final analysis of cohort 1 from KAMILLA, a single-arm phase IIIb clinical trial(☆). *Ann Oncol* 2020;31(10):1350-8.
18. Lin NU, Borges V, Anders C, Murthy RK, Paplomata E, Hamilton E, et al. Intracranial efficacy and survival with tucatinib plus trastuzumab and capecitabine for previously treated HER2-positive breast cancer with brain metastases in the HER2CLIMB Trial. *J Clin Oncol* 2020;38(23):2610-9.
19. Murthy RK, Loi S, Okines A, Paplomata E, Hamilton E, Hurvitz SA, et al. Tucatinib, trastuzumab, and capecitabine for HER2-positive metastatic breast cancer. *N Engl J Med* 2020;382(7):597-609.
20. Curigliano G, Mueller V, Borges V, Hamilton E, Hurvitz S, Loi S, et al. Tucatinib versus placebo added to trastuzumab and capecitabine for patients with pretreated HER2+ metastatic breast cancer with and without brain metastases (HER2CLIMB): final overall survival analysis. *Ann Oncol* 2022;33(3):321-9.

21. Kabraji S, Ni J, Sammons S, Van Swearingen AED, Wang Y, Pereslete AM, et al. Preclinical and clinical efficacy of trastuzumab deruxtecan in breast cancer brain metastases (BCBM). *Can Res* 2022;82(4_Supplement). Abstract PD4-05-PD4.
22. Bartsch R, Berghoff AS, Furtner J, Marhold M, Bergen ES, Roider-Schur S, et al. Trastuzumab deruxtecan in HER2-positive breast cancer with brain metastases: a single-arm, phase 2 trial. *Nat Med* 2022;28(9):1840-7.
23. Hurvitz S, Kim SB, Chung WP, Im SA, Park YH, Hegg R, et al. Trastuzumab deruxtecan (T-DXd; DS-8201a) vs. trastuzumab emtansine (T-DM1) in patients (pts) with HER2+ metastatic breast cancer (mBC): subgroup analyses from the randomized phase 3 study DESTINY-Breast03. *Cancer Res* 2022;82 (4_suppl). Abstract GS3-01..
24. Pérez-García JM, Batista MV, Cortez P, Ruiz-Borrego M, Cejalvo JM, de la Haba-Rodríguez J, et al. Trastuzumab deruxtecan in patients with central nervous system involvement from HER2-positive breast cancer: The DEBBRAH Trial. *Neuro Oncol* 2022;noac144. doi: 10.1093/neuonc/noac144.
25. Tarantino P, Hamilton E, Tolaney SM, Cortes J, Morganti S, Ferraro E, et al. HER2-low breast cancer: Pathological and clinical landscape. *J Clin Oncol* 2020;38(17):1951-62.
26. Schettini F, Chic N, Brasó-Maristany F, Paré L, Pascual T, Conte B, et al. Clinical, pathological, and PAM50 gene expression features of HER2-low breast cancer. *NPJ Breast Cancer* 2021;7(1):1.

27. Modi S, Jacot W, Yamashita T, Sohn J, Vidal M, Tokunaga E, et al. Trastuzumab deruxtecan in previously treated HER2-low advanced breast cancer. *N Engl J Med* 2022;387(1):9-20 .
28. Eisenhauer EA, Therasse P, Bogaerts J, Schwartz LH, Sargent D, Ford R, et al. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur J Cancer* 2009;45(2):228-47.

Tables

Table 1. Systemic efficacy of T-DXd 5.4 mg/kg

	BM subgroup (n = 24)	non-BM subgroup (n = 160)	All patients (N = 184)
Confirmed ORR (CR + PR) by ICR, n (%)^a	14 (58.3) 95% CI, 36.6-77.9	98 (61.3) 95% CI, 53.2-68.8	112 (60.9) 95% CI, 53.4-68.0
CR	1 (4.2)	10 (6.3)	11 (6.0)
PR	13 (54.2)	88 (55.0)	101 (54.9)
SD	8 (33.3)	59 (36.9)	67 (36.4)
PD	1 (4.2)	2 (1.3)	2 (1.1)
NE	1 (4.2)	1 (0.6)	3 (1.6)
BOR in the brain (CR + PR), n (%)^b	8 (47.1)		
DOR (CR or PR), median mo (95% CI)	16.9 (5.7-16.9)	14.8 (13.8-NE)	14.8 (13.8-16.9)
PFS, median mo (95% CI)	18.1 (6.7-18.1)	16.4 (12.7-NE)	16.4 (12.7-NE)

Abbreviations: BOR, best overall response; DOR, duration of response.

^aData are for all randomly assigned patients who received ≥ 1 dose of T-DXd 5.4 mg/kg and had measurable tumors based on ICR at baseline ($N = 184$).

^bBy investigator assessment. Percentage expressed out of number of patients with baseline BM measurements as provided by the investigator ($n = 17$).

Table 2. Prior treatment and brain lesion measurements over time for patients in the BM subgroup

Patient	Prior BM treatment ^a	Imaging method ^b	Longest diameter of lesion (mm) at baseline and time following treatment											BOR in brain lesion ^c	BOR
			Baseline	Week 6	Week 12	Week 18	Week 24	Week 30	Week 36	Week 42	Week 48	Week 54	Week 60		
1	RT	MRI	9	9	7	7	7	7	7	7	7	7	7	SD	PR
2	Surgery, RT, capecitabine	CT	43	25.1	15.9	12.6	12.0	10.6	10.9	8.0	7.9	10.2	10.0	PR	PR
3	Surgery, RT	MRI	15	15	15	15	15	15	15	15	15	15	15	SD	PR
4	NR	CT	46	38	10	10	10	10	10	10	10	10		SD	PR
5	RT	CT	12	NR	CR	CR	CR	CR	CR	CR	CR	PD		PR	PR
		CT	18	NR	12	8	7	7	7	7	7	12			
		CT	10	NR	CR	CR	CR	CR	CR	CR	CR	PD			
6	RT	MRI	13	8	8	9	8	8	8	8				PR	SD
7	NR	CT	8	8	8	7	7	7						SD	SD
8	NR	MRI	17.1	19	19	16	16	16						SD	PR
9	RT	MRI	5.2	2.2	3.0	3.0								PR	PR
10	Surgery, RT	MRI	4.7	3.2	3.9	3.5								PR	SD
11	RT	MRI	23	15	13	13								PR	PR
12	RT	MRI	25	5	5	5								PR	PR
13	RT	MRI	5	5	5	5								SD	SD
14	RT	MRI	8	8	7									SD	PR
15 ^d	RT	CT	10.8	6.9	4.8									PR	SD
16	RT	MRI	7	9	13									PD	SD
17	RT	MRI	5	5										SD	SD
18	Surgery, RT	None	N/A											N/A	PR
19	RT	None	N/A											N/A	PR
20	RT														SD
21	RT	None	N/A											N/A	PR

Patient	Prior BM treatment ^a	Imaging method ^b	Longest diameter of lesion (mm) at baseline and time following treatment										BOR in brain lesion ^c	BOR	
			Baseline	Week 6	Week 12	Week 18	Week 24	Week 30	Week 36	Week 42	Week 48	Week 54			Week 60
22	NR	None	N/A											N/A	NE
23	Surgery	None	N/A											N/A	PD
24	NR	MRI	Data not available for this patient											CR	

BM, brain metastasis; BOR, best overall response; CR, complete response; CT, computed tomography; MRI, magnetic resonance imaging; N/A, not applicable (no brain lesion present at baseline); NE, nonevaluable; NR, nothing reported; PD, progressive disease; Q6W, every 6 weeks; RT, radiation therapy (brain-specific); SD, stable disease.

“RT” includes whole brain RT, brain-directed stereotactic RT, and brain-directed radiosurgery. “Surgery” includes any brain-directed surgery (craniotomy, metastasectomy in brain, resection or removal of brain lesion).

^aIncludes BM-specific treatment for past BMs as well as any at baseline. This data was collected retrospectively.

^bImaging was performed per protocol Q6W for subjects with baseline BMs. Measurements were not required per protocol but were performed by some sites and collected retrospectively. For subjects without measurements, imaging was performed but measurements were either not taken or not provided by the study site.

^cProvided by investigator, retrospectively

Figures

Figure 1. Progression-free survival. Kaplan-Meier analyses in the BM and non-BM subgroups. The tick marks in each panel indicate patient data that were censored.


Figure 2. Central nervous system response. (A) Best response in brain lesions in patients with stable BMs in the BM subgroup. (B) Brain lesion measurements over time in patients with stable BMs in the BM subgroup.

Three patients with reported baseline measurements had no change over time.

Two patients with brain metastases at baseline did not have sufficient data to evaluate response in the brain and are not shown.

Figure 1.

No Brain Metastases at Baseline



Patients still at risk:

	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	
BM (n=24)	24	23	22	18	18	17	17	14	13	13	11	10	6	2	1	1	1	1	1	0	0	0	0	0	0	0	0	0
Non-BM (n=160)	160	159	152	137	135	118	104	93	90	81	58	44	32	15	10	9	8	3	2	1	0	0	0	0	0	0	0	0

Downloaded from <http://aacrjournals.org/cancerres> by guest on 02/08/2016. DOI: 10.1158/2159-8290.CCR-12-0892

