

A Hereditary Nonpolyposis Colorectal Carcinoma Case Associated with Hypermethylation of the *MLH1* Gene in Normal Tissue and Loss of Heterozygosity of the Unmethylated Allele in the Resulting Microsatellite Instability-High Tumor¹

Isabella Gazzoli,² Massimo Loda, Judy Garber, Sapna Syngal, and Richard D. Kolodner

Ludwig Institute for Cancer Research [I. G., R. D. K.], Department of Medicine [R. D. K.], and Cancer Center [R. D. K.], University of California San Diego School of Medicine, La Jolla, California 92093, and Dana Farber Cancer Institute [M. L., J. G., S. S.] and Brigham and Women's Hospital [M. L., S. S.], Harvard Medical School, Boston, Massachusetts 02115

Abstract

Fourteen suspected hereditary nonpolyposis colorectal carcinoma cases with microsatellite unstable (microsatellite instability-high; MSI-H) tumors but no germ-line *MSH2*, *MSH6*, or *MLH1* mutations were examined for hypermethylation of CpG sites in the critical promoter region of *MLH1*. The methylation patterns were determined using methylation-specific PCR and by sequence analysis of sodium bisulfite-treated genomic DNA. In one case, DNA hypermethylation of one allele was detected in DNA isolated from blood. In the MSI-H tumor from this case, the unmethylated *MLH1* allele was eliminated by loss of heterozygosity, and the methylated allele was retained. This biallelic inactivation resulted in loss of expression of *MLH1* in the tumor as confirmed by immunohistochemistry. These results suggest a novel mode of germ-line inactivation of a cancer susceptibility gene.

Introduction

Approximately 60–90% of CpG dinucleotides are methylated on cytosine in the human genome, although unmethylated GC-rich regions are frequently associated with transcriptionally active genes. Epigenetic alteration of the human genome can affect cytosine methylation and chromatin structure (1–3). A number of studies have demonstrated that DNA methylation can directly prevent the binding of transcription factors, such as E2F, cyclic AMP-responsive element binding protein, and USF (4–6), or act through the binding of methylated CpG-binding proteins to induce chromatin configurations that interfere with the transcription machinery (1, 7). As a consequence, changes in the methylation of CpG-rich regions can result in altered gene expression that is independent of genetic alteration of either coding or regulatory sequences. In the last decade, a number of studies have reported that aberrantly hypermethylated CpG islands are associated with transcriptional silencing of tumor suppressor genes in sporadic cancers (reviewed in Refs. 2, 3). Thus, methylation-associated transcriptional silencing is an alternative to mutational inactivation as a cause of loss of tumor suppressor gene function. Consequently, the analysis of DNA cytosine methylation patterns provides an alternative strategy for the study of tumor suppressor gene inactivation in cases where mutations are not observed. One

of many such examples is the MMR³ gene *MLH1*, where hypermethylation of the *MLH1* promoter region is associated with loss of expression and appears to underlie the majority of MMR-defective sporadic cancer cases (8–11).

In a previous study (12, 13), we analyzed 70 HNPCC cases (which met Amsterdam, Modified Amsterdam, HNPCC-like, or Bethesda criteria) for germ-line defects in *MSH2*, *MLH1*, and *MSH6* genes. For 48 of the cases, a tumor sample was available, and the MSI status of the tumor was determined; 14 cases were MSI-H and showed a germ-line mutation in *MSH2* or *MLH1*, and 14 cases were MSI-H but did not have a germ-line mutation in *MSH2*, *MSH6*, or *MLH1* (14). The observation of HNPCC cases that have MSI-H tumors but no germ-line mutation in a known MMR gene, as detected by DNA sequencing or other methods for identifying point mutations, has been reported by many studies. Some of these types of case have been attributed to germ-line deletion mutations in *MSH2* or *MLH1* as well as uncharacterized mutations that cause loss of expression of *MLH1* (15, 16). In the present study, the mutation-negative 14 HNPCC cases were examined for hypermethylation of CpG sites in the critical promoter region of *MLH1* (–290 to –180 bp relative to the ATG), which shows a strong correlation with the loss of *MLH1* expression (17). DNA from blood of one of these 14 cases was found to have one methylated *MLH1* allele, and analysis of the MSI-H tumor from this case demonstrated that LOH eliminated the unmethylated *MLH1* allele, resulting in loss of expression of *MLH1*.

Materials and Methods

Tissue and DNA Samples. The HNPCC cases used in the studies described here have been described in detail elsewhere (12, 13). DNA from blood was isolated as described previously and analyzed for the presence of mutations in *MSH2*, *MLH1*, and *MSH6* by DNA sequencing as reported previously. Tumor DNA from each case was analyzed for MSI using the National Cancer Institute recommended five microsatellite markers as described previously (14). The 14 cases studied here comprised all 14 of the MSI-H cases that did not have mutation in *MSH2*, *MLH1*, or *MSH6* described previously (14). Control genomic DNA in which *MLH1* was not methylated was from blood of the case DF000268 containing the germ-line *MSH2* IVS5 + 3A>T splice site mutation (12). Control genomic DNA in which *MLH1* was fully methylated was isolated from SW48 cells (American Type Culture Collection CCL-231) with a Puregene DNA isolation kit (Gentra Systems).

Analysis of *MLH1* Methylation Using MS-PCR. One μg of total genomic DNA was modified by sodium bisulfite treatment using a CpGenome Modification kit (Intergen) according to the manufacturer's instructions and suspended in 10 μl of 1 \times TE buffer [10 mM Tris-HCl (pH 7.6), 1 mM EDTA]. The methylation status of the *MLH1* gene was then determined by MS-PCR

Received 4/25/02; accepted 6/5/02.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ This work was supported of a grant from the Starr Foundation and Grant CA06516 from the National Cancer Institute to the Dana-Farber Cancer Institute and Grant CA23100 from the University of California, San Diego Medical School, both Cancer Center Support Grants.

² To whom requests for reprints should be addressed, at Ludwig Institute for Cancer Research, UCSD School of Medicine-CMME 3019, 9500 Gilman Drive, La Jolla, CA 92093-0669. Phone: (858) 822-1181; Fax: (858) 534-7750; E-mail: igazzoli@ucsd.edu.

³ The abbreviations used are: MMR, mismatch repair; HNPCC, hereditary nonpolyposis colorectal carcinoma; MSI, microsatellite instability; MSI-H, MSI-high; LOH, loss of heterozygosity; MS-PCR, methylation-specific PCR.




Fig. 1. Analysis of the methylation status of the *MLH1* gene in an HNPCC case using MS-PCR. *M* and *U*, the 91- and 103-bp PCR products amplified using primers specific for methylated and unmethylated alleles, respectively. *B*, blood DNA; *T*, tumor DNA; *Me+*, DNA from the SW48 cell line in which *MLH1* is fully methylated; *Me-*, DNA sample with a genetic alteration in *MSH2* gene, in which *MLH1* is unmethylated.

using primers specific for both methylated and unmethylated sequences as essentially described previously (18). PCR reactions (20 μ l) contained 1.5 μ l of modified DNA and were performed using an Advantage GC2 PCR kit (Clontech). The thermocycling conditions used were 1 cycle of 94°C for 2 min; 3 cycles of 94°C for 30 s, 60°C for 30 s, 68°C for 30 s; 36 cycles 94°C for 30 s, 53°C for 30 s, 68°C for 30 s, and 1 cycle of 68°C for 2 min. Three μ l of each PCR reaction were then analyzed by electrophoresis through nondenaturing 10% polyacrylamide gels, which were stained with ethidium bromide and visualized under UV illumination.

Analysis of *MLH1* by DNA Sequencing. To analyze the methylation status of *MLH1* by direct DNA sequencing, genomic DNA was modified by treatment with sodium bisulfite as described above. The promoter region of the *MLH1* gene was amplified using two rounds of PCR with nested primers using an Advantage GC2 PCR kit. The first round of amplification used the primers 5'-GTTTGAGAAGTGTTAAGTATTTTTT-3' and 5'-CAAATAACCCCT-ACCACAAAC-3', and the second round of amplification used the primers 5'-GTATTTTTTTGTTTTGTYGTTAG-3' and 5'-CTATTAATTAACAAC-TTAAATACCAATC-3'. The final PCR product then was treated with shrimp alkaline phosphatase and exonuclease I (Amersham) to remove excess PCR primers and nucleotides, and the resulting PCR product was sequenced using a PE/ABI 377 sequencer and dye terminator chemistry (PE Applied Biosystems).

To analyze the presence of a polymorphism in the 5'-untranslated region of the *MLH1* gene, genomic DNA from tumor and blood samples was used as template in PCR reactions using primers 5'-GAAAAGTAGAGCCTCGTC-GACTT-3' (sense) and 5'-TAGCATTAGCTGGCCGCTGGATAAC-3' (anti-sense) essentially as described previously. The resulting PCR product was then purified and sequenced using a PE/ABI 377 sequencer and dye terminator chemistry (PE Applied Biosystems), and the sequence data obtained was analyzed using Sequencher 3.1 software (GeneCodes, Inc.).

Immunohistochemistry (IHC) Analysis of *MLH1* Expression. Tissue sections were stained with anti-*MLH1* antibody (G168-728; PharMingen) as previously described (19). The results of immunohistochemistry analysis were kept blinded until the *MLH1* methylation analysis was completed.

Results

In a previous study, we identified 14 suspected HNPCC cases (which met Amsterdam, Modified Amsterdam, HNPCC-like, or Bethesda criteria) that had MSI-H tumors but did not have a germ-line mutation in *MSH2*, *MSH6*, or *MLH1*. To understand the genetic basis for the MMR defect in these 14 HNPCC cases, we screened them for hypermethylation of the *MLH1* gene. The *MLH1* DNA methylation pattern was determined by sequencing PCR products amplified from bisulfite-modified DNA isolated from blood of all 14 mutation-negative cases and by MS-PCR using bisulfite-modified DNA from selected cases as template (18). Cytosine methylation protects against bisulfite modification-induced conversion of cytosine to uracil. Consequently, during sequencing of modified DNA, unmethylated cytosines appear as thymines, methylated cytosines appear as cytosines, and mixtures appear heterozygous. Similarly, during MS-PCR, the

primers specific for the sequences that result after bisulfite modification of methylated and unmethylated DNA generate two fragments of different lengths containing the same region, 91 and 103 bp long, respectively. The size of the resulting fragments allows discrimination between methylated and unmethylated alleles by acrylamide gel electrophoresis. The region of *MLH1* selected for analysis, -290 to -180 bp relative to the ATG, contains nine CpG sites close to the CCAAT box and different transcription factors binding sites, and hypermethylation of this region is associated with loss of expression of *MLH1*.

Using MS-PCR, one case (which met Bethesda criteria 4 alone; diagnosis at 25 years of age) was found to have both methylated and unmethylated *MLH1* alleles when blood DNA was analyzed, and only methylated alleles when tumor DNA was analyzed (Fig. 1). Only unmethylated alleles were observed when unmethylated control DNA was analyzed, and conversely, only methylated alleles were observed when fully methylated DNA was analyzed from the SW48 cell line, which is MMR defective and does not express *MLH1*. To further analyze the hypermethylation status of this region, we PCR amplified the region from bisulfite-modified DNA and sequenced the resulting PCR products (Fig. 2). The chromatograms showed that the *MLH1* promoter region was partially methylated (a mixture of C and T is seen at critical Cs in the region analyzed) in blood DNA and totally methylated (only C is seen at the critical Cs in the region analyzed) in tumor DNA; analysis of an area containing two CpG sites (8 and 15 bp downstream from the CAAT box located at position -282) is




Fig. 2. Sequence analysis of bisulfite-modified DNA. *Control*, unmethylated DNA from a *MSH2* mutant HNPCC case showing all Ts at positions -268, -264, and -261 attributable to complete modification of the DNA; *Blood*, DNA isolated from blood from the case showing a mixture of C and T at positions -268, -264, and -261 attributable to partial modification of the DNA because of partial methylation of the DNA; *Tumor*, DNA isolated from tumor from the case showing an increase in the level of C at positions -268, -264, and -261 attributable to reduced modification of the DNA because of increased methylation of the DNA; *SW48*, DNA from the SW48 cell line in which *MLH1* is completely methylated showing a high level C at positions -268, -264, and -261 attributable to reduced modification because of complete methylation of the DNA.




Fig. 3. Detection of LOH by sequence analysis of the 5' untranslated region of the *MLH1* gene. DNA isolated from blood was heterozygous for a G→A polymorphism at position -93 bp, whereas DNA from the tumor only had a G at this position.

shown in Fig. 2. Sequencing of unmethylated and fully methylated control DNAs detected only unmethylated and methylated alleles, respectively (Fig. 2). Sequencing 20 cloned PCR products from each sample (data not shown) revealed a mixture of methylated and unmethylated clones from blood DNA and an increased proportion of methylated clones from tumor DNA, consistent with the results obtained by direct sequencing of PCR products shown in Fig. 2. The observation that ~50% of the DNA from blood from the mutation-negative HNPCC case described above showed *MLH1* methylation by direct bisulfite sequencing suggests that the observed *MLH1* methylation is unlikely to result from shedding of tumor cells into the blood (18), because this would likely yield lower levels on *MLH1* methylation than observed here.

The above analysis documented an HNPCC case in which normal tissue contained a mixture of methylated and unmethylated *MLH1* alleles, whereas tumor tissue contained only methylated *MLH1* alleles. There are two possible explanations for the observation that the blood DNA contained one methylated and one unmethylated *MLH1* allele while the tumor DNA only contained methylated alleles; the

unmethylated allele could have been deleted or methylated in the tumor. Sequencing of the promoter region from unmodified DNA from this case revealed that the blood DNA was heterozygous for a previously described single nucleotide G→A polymorphism (Fig. 3; Ref. 20). When the same region of unmodified DNA from the tumor sample was sequenced, only the G allele was observed. This indicates that LOH had occurred in the tumor, resulting in the loss of the unmethylated allele (Fig. 3). Consistent with loss of the unmethylated, expressed *MLH1* allele in the tumor, immunohistochemistry analysis showed that this case did not express MLH1 in the tumor (Fig. 4), whereas MSH2 was still expressed (data not shown). These observations are in contrast to the observation of biallelic methylation of *MLH1* in sporadic colon tumors leading to loss of MLH1 expression (21).

Discussion

A growing body of evidence has suggested that the majority of HNPCC families showing MSI can be accounted for by mutations in *MSH2* and *MLH1* (see Refs. 12, 13, 14, 16). Initial studies documented missense, nonsense, frameshift, splice site, and specific deletion mutations in *MSH2* and *MLH1*, and more recent studies have documented large *MSH2* and *MLH1* deletion mutations and uncharacterized mutations that eliminate expression of these genes (15, 16). In the present study, we have identified an HNPCC case in which one allele of *MLH1* was methylated in DNA isolated from normal tissue (blood) and somatic LOH of the unmethylated allele subsequently occurred, resulting in a tumor that did not express MLH1 and was MMR defective. These results raise the possibility that methylation and associated silencing of *MLH1* could represent a germ-line alteration that underlies some HNPCC cases; however, samples from the parents were not available. Therefore, it was not possible to determine whether the methylated allele was actually inherited. We have detected one such case in our sample set of 14 mutation-negative, MSI-H HNPCC cases, indicating that germ-line silencing of *MLH1* could underlie a small but significant proportion of HNPCC cases. Larger scale studies will be required to definitively identify other HNPCC cases similar to the one described here and to determine how frequently this unique mechanism of tumorigenesis occurs.




Fig. 4. Immunohistochemistry analysis of MLH1 expression. MLH1 staining is in brown. N, normal tissue; T, tumor tissue.

The region of *MLH1* found to be methylated in our studies contains CpG sites close to the CCAAT box at position -282, and methylation of this region is associated with inactivation of the *MLH1* gene. The CCAAT box is recognized by CBF, and binding of this factor effects *MLH1* promoter activity. Functional studies have shown that methylation of this region interferes with binding of CBF, resulting in decreased transcription of *MLH1* (22). In addition to decreasing the binding of critical transcription factors, methylation of this region of *MLH1* likely increases the affinity of methyl-binding proteins and histone deacetylases, leading to the assembly of inactive chromatin configurations. Consistent with this view, full reactivation in tumor cell lines requires treatment with both inhibitors of DNA methylation and histone deacetylation (23). These observations support the view that the methylation of the region of *MLH1* observed here is directly associated with or underlies the silencing of the gene observed in the HNPCC case reported here.

A number of examples of loss of function of different tumor suppressor genes attributable to somatic silencing of the genes associated with hypermethylation have been reported. The types of examples reported in sporadic cancer include biallelic methylation, somatic methylation associated with LOH, and mutation associated with methylation (reviewed in Ref. 2). These observations have led to the hypothesis that hypermethylation and associated silencing of tumor suppressor genes represents an alternate pathway of the two-hit tumor suppressor gene inactivation hypothesis (2, 3). In those cases where methylation associated with either mutation or LOH has been observed, methylation appears to have been a somatic event. In at least one case, methylation of p16^{INK4} was observed in both recurrent tumors and adjacent normal epithelium, suggesting that methylation in normal tissue might underlie subsequent tumor development (24). In addition, age-related methylation of *MLH1* has been observed in normal colonic mucosa associated with the development of colon tumors, with high-level methylation being associated with the development of MSI-H tumors, supporting the idea that the age-related development of *MLH1* methylation is a precursor lesion (9). Our results raise the possibility that methylation and associated silencing of a gene like *MLH1* could actually represent a germ-line change that can underlie development of cancer. Alternately, somatic methylation could occur much more widely in normal tissues than thought previously (9). This suggests a novel mechanism of tumorigenesis that nonetheless fits the two-hit tumor suppressor gene inactivation hypothesis, although larger scale studies are clearly required to verify our observations and determine how prevalent this mechanism is. In this regard, a recent study detected three cases of HNPCC where *MLH1* expression was absent in the tumors, but a germ-line *MLH1* mutation could not be found; however, *MLH1* methylation was not analyzed in these cases (16).

References

- Tate, P. H., and Bird, A. P. Effects of DNA methylation on DNA-binding proteins and gene expression. *Curr. Opin. Genet. Dev.*, 3: 226-231, 1993.
- Jones, P. A., and Laird, P. W. Cancer epigenetics comes of age. *Nat. Genet.*, 21: 163-167, 1999.
- Rountree, M. R., Bachman, K. E., Herman, J. G., and Baylin, S. B. DNA methylation, chromatin inheritance, and cancer. *Oncogene*, 20: 3156-3165, 2001.
- Kovesdi, I., Reichel, R., and Nevins, J. R. Role of an adenovirus E2 promoter binding factor in E1A-mediated coordinate gene control. *Proc. Natl. Acad. Sci. USA*, 84: 2180-2184, 1987.
- Iguchi-Ariga, S. M., and Schaffner, W. CpG methylation of the cAMP-responsive enhancer/promoter sequence TGACGTCA abolishes specific factor binding as well as transcriptional activation. *Genes Dev.*, 3: 612-619, 1989.
- Watt, F., and Molloy, P. L. Cytosine methylation prevents binding to DNA of a HeLa cell transcription factor required for optimal expression of the adenovirus major late promoter. *Genes Dev.*, 2: 1136-1143, 1988.
- Graessmann, A., and Graessmann, M. DNA methylation, chromatin structure and regulation of *Herpes simplex virus tk* gene expression. *Gene (Amst.)*, 74: 135-137, 1988.
- Herman, J. G., Umar, A., Polyak, K., Graff, J. R., Ahuja, N., Issa, J. P., Markowitz, S., Willson, J. K., Hamilton, S. R., Kinzler, K. W., Kane, M. F., Kolodner, R. D., Vogelstein, B., Kunkel, T. A., and Baylin, S. B. Incidence and functional consequences of hMLH1 promoter hypermethylation in colorectal carcinoma. *Proc. Natl. Acad. Sci. USA*, 95: 6870-6875, 1998.
- Nakagawa, H., Nuovo, G. J., Zervos, E. E., Martin, E. W., Jr., Salovaara, R., Aaltonen, L. A., and de la Chapelle, A. Age-related hypermethylation of the 5' region of *MLH1* in normal colonic mucosa is associated with microsatellite-unstable colorectal cancer development. *Cancer Res.*, 61: 6991-6995, 2001.
- Kuismanen, S. A., Holmberg, M. T., Salovaara, R., Schweizer, P., Aaltonen, L. A., de la Chapelle, A., Nystrom-Lahiti, M., and Peltomaki, P. Epigenetic phenotypes distinguish microsatellite-stable and -unstable colorectal cancers. *Proc. Natl. Acad. Sci. USA*, 96: 12661-12666, 1999.
- Wheeler, J. M., Beck, N. E., Kim, H. C., Tomlinson, I. P., Mortensen, N. J., and Bodmer, W. F. Mechanisms of inactivation of mismatch repair genes in human colorectal cancer cell lines: the predominant role of hMLH1. *Proc. Natl. Acad. Sci. USA*, 96: 10296-10301, 1999.
- Syngal, S., Fox, E., Li, C., Dovidio, M., Eng, C., Kolodner, R. D., and Garber, J. E. Interpretation of genetic test results for hereditary nonpolyposis colon cancer. *J. Am. Med. Assoc.*, 282: 247-253, 1999.
- Kolodner, R. D., Tytell, J. D., Schmeits, J., Kane, M. F., Das Gupta, R., Weger, J., Wahlberg, S., Fox, E. A., Peel, D. J., Ziogas, A., Garber, J. E., Syngal, S., Anton-Culver, H., and Li, F. P. Germline *MSH6* mutations in colorectal cancer families. *Cancer Res.*, 59: 5068-5074, 1999.
- Wahlberg, S. S., Schmeits, J., Thomas, G., Loda, M., Garber, J. E., Syngal, S., Kolodner, R. D., and Fox, E. A. Evaluation of microsatellite instability and immunohistochemistry for the prediction of germline *MSH2* and *MLH1* mutations in HNPCC families. *Cancer Res.*, 62: 3485-3492, 2002.
- Charbonnier, F., Olschwang, S., Wang, Q., Boisson, C., Martin, C., Buisine, M. P., Puisieux, A., and Frebourg, T. *MSH2* in contrast to *MLH1* and *MSH6* is frequently inactivated by exonic and promoter rearrangements in hereditary nonpolyposis colorectal cancer. *Cancer Res.*, 62: 848-853, 2002.
- Yan, H., Papadopoulos, N., Marra, G., Perra, C., Jiricny, J., Boland, C. R., Lynch, H. T., Chadwick, R. B., de la Chapelle, A., Berg, K., Eshleman, J. R., Yuan, W., Markowitz, S., Laken, S. J., Lengauer, C., Kinzler, K. W., and Vogelstein, B. Conversion of diploidy to haploidy. *Nature (Lond.)*, 403: 723-724, 2000.
- Deng, G., Chen, A., Hong, J., Chae, H. S., and Kim, Y. S. Methylation of CpG in a small region of the hMLH1 promoter invariably correlates with the absence of gene expression. *Cancer Res.*, 59: 2029-2033, 1999.
- Grady, W. M., Rajput, A., Lutterbaugh, J. D., and Markowitz, S. D. Detection of aberrantly methylated hMLH1 promoter DNA in the serum of patients with microsatellite unstable colon cancer. *Cancer Res.*, 61: 900-902, 2001.
- Kane, M. F., Loda, M., Gaida, G. M., Lipman, J., Mishra, R., Goldman, H., Jessup, J. M., and Kolodner, R. D. Methylation of the hMLH1 promoter correlates with lack of expression of hMLH1 in sporadic colon tumors and mismatch repair-defective human tumor cell lines. *Cancer Res.*, 57: 808-811, 1997.
- Ito, E., Yanagisawa, Y., Iwahashi, Y., Suzuki, Y., Nagasaki, H., Akiyama, Y., Sugano, S., Yuasa, Y., and Maruyama, K. A core promoter and a frequent single-nucleotide polymorphism of the mismatch repair gene *hMLH1*. *Biochem. Biophys. Res. Commun.*, 256: 488-494, 1999.
- Veigl, M. L., Kasturi, L., Olechnowicz, J., Ma, A. H., Lutterbaugh, J. D., Periyasamy, S., Li, G. M., Drummond, J., Modrich, P. L., Sedwick, W. D., and Markowitz, S. D. Biallelic inactivation of hMLH1 by epigenetic gene silencing, a novel mechanism causing human MSI cancers. *Proc. Natl. Acad. Sci. USA*, 95: 8698-8702, 1998.
- Deng, G., Chen, A., Pong, E., and Kim, Y. S. Methylation in hMLH1 promoter interferes with its binding to transcription factor CBF and inhibits gene expression. *Oncogene*, 20: 7120-7127, 2001.
- Cameron, E. E., Bachman, K. E., Myohanen, S., Herman, J. G., and Baylin, S. B. Synergy of demethylation and histone deacetylase inhibition in the re-expression of genes silenced in cancer. *Nat. Genet.*, 21: 103-107, 1999.
- Muto, S., Horie, S., Takahashi, S., Tomita, K., and Kitamura, T. Genetic and epigenetic alterations in normal bladder epithelium in patients with metachronous bladder cancer. *Cancer Res.*, 60: 4021-4022, 2000.