

Mutagenicity of Several Classes of Antitumor Agents to *Salmonella typhimurium* TA98, TA100, and TA92¹

Yuko Seino,² Minako Nagao, Takie Yahagi, Akio Hoshi, Takashi Kawachi, and Takashi Sugimura

Biochemistry [Y. S., M. N., T. Y., T. K., T. S.] and Pharmacology [A. H.] Divisions, National Cancer Center Research Institute, Tsukiji, Chuo-ku, Tokyo 104, Japan

ABSTRACT

The mutagenic activities of antitumor agents, including 5 antibiotics, 19 antimetabolites, 5 alkylating agents, 2 alkaloids, 1 enzyme, and 1 adrenal steroid hormone, were tested on *Salmonella typhimurium* TA100, TA98, and TA92.

Four of these, busulfan, carbazilquinone, 1-(4-amino-2-methylpyrimidine-5-yl)methyl-3-(2-chloroethyl)-3-nitrosourea hydrochloride, and pipobroman were shown for the first time to be mutagenic. Further, the known mutagenicities of five others, daunomycin hydrochloride, Adriamycin hydrochloride, mitomycin C, 6-mercaptopurine, and cyclophosphamide, were confirmed.

INTRODUCTION

Many chemotherapeutic agents are now widely used clinically for treatment of cancers and leukemias and sometimes for prophylaxis of recurrent cancer after surgical treatment.

However, a phenomenon known as Haddow's paradox (5) is that some cancer chemotherapeutic agents are carcinogenic to laboratory animals and also induce chromosomal aberrations in cultured mammalian cells. Thus, it is important to know whether chemotherapeutic agents are carcinogenic.

Recently, it has become widely accepted that many mutagens are carcinogens (15, 16, 24). Therefore, in this work we tested the mutagenic activities in *S. typhimurium* strains TA100, TA98, and TA92 of various kinds of chemotherapeutic agents now in use or being tested for use in treatment of human cancer. The compounds tested were antibiotics, antimetabolites, alkylating agents, alkaloids, an enzyme, and a hormone.

MATERIALS AND METHODS

Microbes. *S. typhimurium* strains TA100, TA98, and TA92 were kindly supplied by Dr. Bruce N. Ames, University of California, Berkeley, Calif. TA100 and TA98 are *uvrB* and *rfa* (deep rough) mutants (1). TA92 has the capacity for excision repair and an intact lipopolysaccharide barrier (LPS⁺) (1). TA100 and TA92 both contain the same base-pair change mutation, and TA98 contains a frameshift mutation at *his* locus (1). All these strains carry R-factor plasmid pKM101 (1).

Chemicals. Adriamycin hydrochloride, 5-fluorouracil, mitomycin C, and L-asparaginase were obtained from Kyowa Hakko Kogyo Co., Ltd., Tokyo, Japan. Cyclophosphamide and vinblastine sulfate were purchased from Shionogi & Co., Ltd., Osaka, Japan; Methotrexate and busulfan were from Takeda Chemical Industries, Ltd., Osaka, Japan; 1-β-D-arabinofuranosylcytosine and cyclocytidine were from Kohjin Co., Ltd., Tokyo, Japan. 6-Mercaptopurine and 8-azaguanine were from Sigma Chemical Co., St. Louis, Mo.; ACNU³ and carbazilquinone were from Sankyo Co., Ltd., Tokyo, Japan; and actinomycin D was from Nippon Merck-Banyu Co., Ltd., Tokyo, Japan. Bleomycin hydrochloride was from Nippon Kayaku Co., Ltd., Tokyo, Japan; daunomycin hydrochloride was from Meiji Seika Kaisha, Tokyo, Japan; and FT-207 was from Taiho Pharmaceutical Co., Ltd., Tokyo, Japan. Alkylcarbonyl derivatives of 5-fluorouracil were synthesized at Mitsui Pharmaceuticals, Inc., Tokyo, Japan. Vincristine sulfate was from Eli Lilly & Co., Indianapolis, Ind., and pipobroman was from Dainippon Pharmaceutical Co., Ltd., Osaka, Japan. Prednisolone was from Sanwa Research Institute, Tokyo, Japan. Glucose 6-phosphate, glucose-6-phosphate dehydrogenase (EC 1.1.1.49), and ATP were from Sigma; NADH and NADPH were from Oriental Yeast Co., Ltd., Tokyo, Japan; and dimethyl sulfoxide, spectrophotometric grade, was from Wako Pure Chemical, Inc., Osaka, Japan.

Eight compounds for drug use contained some vehicle, and the amounts of these compounds were expressed as μg of active principle. The names of these compounds and the amounts in 1 vial, 1 ampul, or 1 tablet were as follows: daunomycin·HCl (20 mg principle; actual weight, 120 mg/vial), Adriamycin·HCl (10 mg principle; actual weight, 50 mg/vial), bleomycin·HCl (15 mg potency; actual weight, 8 mg/ampul), actinomycin D (0.5 mg principle; actual weight, 20.5 mg/vial), mitomycin C (0.996 mg principle; actual weight, 1 mg/vial), cyclophosphamide (500 mg principle; actual weight, 725 mg/vial), L-asparaginase (10,000 IU; actual weight, 33 mg/vial), and prednisolone (5 mg principle; actual weight, 153 mg/tablet).

Mutation Test. The mutation test was carried out by our modification (20) of the method of Ames *et al.* (1). The test chemical in 0.1 ml dimethyl sulfoxide or H₂O was placed in a tube and mixed with 0.5 ml S-9 mix (150 μl of the S-9 fraction of rat liver pretreated with polychlorinated biphenyl, 2 μmol NADPH, 2 μmol NADH, 2.5 μmol glucose 6-phosphate, 0.25 unit glucose-6-phosphate dehydrogenase, 4 μmol MgCl₂, 16.5 μmol KCl, and 50 μmol sodium phosphate buffer, pH 7.4 and 0.1 ml of culture of the bacterial

¹ This work was supported in part by Grants-in-Aid for Cancer Research from the Ministry of Education, Science and Culture and the Ministry of Health and Welfare, Japan, and by the Princess Takamatsu Cancer Research Fund.

² To whom requests for reprints should be addressed.
Received April 14, 1977; accepted March 22, 1978.

³ The abbreviations used are: ACNU, 1-(4-amino-2-methylpyrimidine-5-yl)methyl-3-(2-chloroethyl)-3-nitrosourea hydrochloride; FT-207, 1-(2-tetrahydrofuryl)-5-fluorouracil.

tester strain (1 to 2×10^8 cells). Without metabolic activation $50 \mu\text{mol}$ sodium phosphate buffer, pH 7.4, in 0.5 ml were used instead of S-9 mix. The mixture was preincubated at 37° for 20 min and then mixed with 2 ml top agar (0.7% agar and 0.6% NaCl) at 45° and spread on a minimal-glucose agar plate containing $0.1 \mu\text{mol}$ each L-histidine and biotin. Plates were incubated at 37° for 2 days , and then his^+ revertant colonies were counted.

RESULTS AND DISCUSSION




The trivial names and structures of the 5 antibiotics, 19 antimetabolites, 5 alkylating agents, and other compounds tested are shown in Tables 1, 3, 5, and 7, respectively, with the results of mutagenicity tests on them and information on their carcinogenicities.

Antibiotics. Results on the mutagenicities of the antibiotics are listed in Table 2. The mutagenicities of daunomycin hydrochloride (16), Adriamycin hydrochloride (16), and mitomycin C (17), which had been reported previously, were confirmed. Daunomycin hydrochloride and Adriamycin hydrochloride intercalate DNA (33), resulting in inhibition of DNA-dependent DNA and RNA polymerases (32). These compounds were highly mutagenic to TA100 and TA98 without S-9 mix, the latter showing more mutagenicity in TA98 than in TA100. Addition of S-9 mix reduced the activity. Clear dose-response activities of these compounds were obtained, as shown in Table 2. These compounds have potent actions in producing malignant transformation

in vivo and *in vitro* and mutation in mammalian cells *in vitro* (14). Bleomycin hydrochloride was not mutagenic to TA100 or TA98, with or without S-9 mix, but it is known to cause DNA strand breakage (26). Actinomycin D, which binds preferentially to dG-dC pairs of DNA (13) and inhibits DNA-dependent RNA synthesis (12), was not mutagenic to TA100 or TA98 with or without S-9 mix. The *in vivo* carcinogenicity of actinomycin D in rats has been reported (30). Actinomycin D is also reported to have very weak mutagenic activity on mammalian cells cultured *in vitro* (14). Mitomycin C was not mutagenic to TA100 or TA98 but was weakly mutagenic to TA92, especially without S-9 mix, as shown in Table 2. Mitomycin C produces interstrand cross-linking of double-stranded DNA (27). In this case *uv* repair-deficient strain cells were killed and did not yield any mutants (19). TA92, which has *uv* repair capacity, gave more revertants than did the spontaneous revertants. Mitomycin C produces tumors *in vivo* (30). Bleomycin hydrochloride and actinomycin D were not mutagenic but were toxic to TA100 and TA98, as was mitomycin C. But these compounds were not mutagenic to TA92 (Table 2).

Antimetabolites. Results on antimetabolites are listed in Table 4. 5-Fluorouracil, an inhibitor of thymidylate synthetase, was not mutagenic to TA100 or TA98 with or without S-9 mix, as shown in Table 4. 5-Fluorouracil had a strong lethal effect on cells at a concentration of $5 \mu\text{g}/\text{plate}$ without S-9 mix. FT-207 also was not mutagenic to TA100 or TA98 with or without S-9 mix. These 2 compounds were also not mutagenic to TA92 with or without S-9 mix.

Table 1
Trivial names, structures, mutagenicities, and carcinogenicities of antibiotics

Trivial name	Structure	Mutagenicity	Carcinogenicity
Daunomycin · HCl	 <p>R: —CH₃, daunomycin —CH₂OH, Adriamycin</p>	+	+ (14)
Adriamycin · HCl		+	+ (14)
Bleomycin · HCl	Glycopeptide	—	
Actinomycin D		—	+ (30)
Mitomycin C ^a		+	+ (30)

^a Mitomycin C is mutagenic to *S. typhimurium* TA92.

Table 2
Results of mutation tests on antibiotics

Drug ^a	μg/plate	<i>his</i> ⁺ revertants/plate						<i>his</i> ⁺ revertants/ nmol ^b
		TA100		TA98		TA92		
		+ S-9 mix	- S-9 mix	+ S-9 mix	- S-9 mix	+ S-9 mix	- S-9 mix	
Daunomycin · HCl	0.083	144	286	24	340			2800 (8, -) ^c
	0.17	160	295	27	852			
	0.42	163	202	33	1555			
	0.83	147	144	62	1368			
Adriamycin · HCl	0.02	109	106	24	102			3130 (8, -)
	0.05	130	309	28	323			
	0.1	152	234	27	556			
	0.2	159	98	22	1034			
	0.5	270	72	31	776			
	1	303	36	55	260			
	2	170	20	106	64			
	5	185	67	423	45			
Bleomycin · HCl	10	168	0	443	0			0
	0.94	99	190	25	17	46	39	
	1.88	111	143	23	18	48	37	
	4.7	125	10	20	7	49	32	
	9.4	129	0	22	0	49	28	
Actinomycin D	18.8	106	0	12	0	45	9	0
	2.4	125	103	18	18	28	24	
	6.1	100	90	17	15	20	34	
	12.2	66	45	21	18	14	32	
Control ^d Mitomycin C	0.01	147	132	30	16			43 (2, -)
	1.0	110	0	21	0	48	90	
	2.5	70	0	21	0	40	120	
	5	75	0	22	0	80	137	
	10	103	0	17	0	61	72	
Control ^e		131	122	19	17	39	26	

^a Drug contained some vehicle, and the amount of the drug was expressed as μg of active principle.

^b The number of *his*⁺ revertants per nmol was calculated from the data obtained under the most active conditions in the linear dose-responsive range and a yield of 100 colonies more than the background yield.

^c Tester strain TA98; -, without S-9 mix; 2, TA92.

^d Average of values in 20 experiments. The numbers of revertants were: 117 to 183 of TA100 with S-9 mix, 89 to 169 of TA100 without S-9 mix, 19 to 38 of TA98 with S-9 mix, and 14 to 26 of TA98 without S-9 mix.

^e Average of values in 2 experiments.

Twelve alkylcarbamoyl derivatives of 5-fluorouracil (namely, 1-methyl, 1-ethyl, 1-propyl, 1-isopropyl, 1-butyl, 1-*tert*-butyl, 1-pentyl, 1-hexyl, 1-cyclohexyl, 1-phenyl, 1-heptyl, and 1-octyl-carbamoyl-5-fluorouracil) were all nonmutagenic to TA100 and TA98 with or without S-9 mix. All of these derivatives were cytotoxic at doses of 2.5 to 5 μg/plate. Therefore, these compounds were also tested in TA92 and were found not to be mutagenic, as shown in Table 4. These alkylcarbamoyl derivatives of 5-fluorouracil are not yet used clinically, although their antitumor activities against experimental tumors in mice were demonstrated by Hoshi et al. (9, 10).

6-Mercaptopurine, an inhibitor of purine synthesis, was weakly mutagenic to TA100 without S-9 mix, as shown in Table 4. This drug was reported to be mutagenic to *S. typhimurium* strains TA1535 and *his*G46 by the liquid method (8). The noncarcinogenicity of this compound was reported (25). Methotrexate, which inhibits the biosyntheses of purine and thymine from formate, gave negative results on TA100 and TA98 at doses of up to 312 μg/plate. It had a slight lethal effect on the cells. Methotrexate was also reported to be nonmutagenic to *S. typhimurium* TA1535, TA1536, TA1537, TA1538, and *his*G46 by the liquid method (8). A weak carcinogenicity of this compound on

experimental animals has been reported (22), but it has also been reported not to be carcinogenic (23) and to cause no chromosomal aberrations in mammalian culture cells (7). Thus, further studies are required to determine whether it is really carcinogenic.

8-Azaguanine, an inhibitor of guanylic acid synthesis, was not mutagenic to TA100, TA98, or TA92 with or without S-9 mix (Table 4). It had no effect on cell survival of TA100 or TA92 at doses of up to 50 μg/plate but killed TA98 at a dose of 25 μg/plate. This compound is noncarcinogenic (6).

1-β-D-Arabinofuranosylcytosine and cycloctidine were not mutagenic at doses of up to 30 and 25 mg/plate, respectively (Table 4). Arabinofuranosylcytosine is known to interfere with DNA polymerase (18) and to be incorporated into DNA (31), and it induced chromosomal aberrations both *in vivo* (2) and *in vitro* (4, 11).

Alkylating Agents. Five alkylating agents were all found to be mutagenic in this system, as shown in Table 6. Cyclophosphamide was mutagenic to TA100 with S-9 mix but was not mutagenic to TA98. This compound was reported to be mutagenic (8, 16) and carcinogenic (29). Busulfan was mutagenic to TA100 with or without S-9 mix but not to TA98. Busulfan was reported to be carcinogenic

Table 3
Trivial names, structures, mutagenicities, and carcinogenicities of antimetabolites

Trivial name	Structure	Mutagenicity	Carcinogenicity
5-Fluorouracil		-	
FT-207		-	
Carbamoyl derivatives of 5-fluorouracil			
R: Methyl		-	
Ethyl		-	
Propyl		-	
Isopropyl		-	
Butyl		-	
tert-Butyl		-	
Pentyl		-	
Hexyl		-	
Cyclohexyl		-	
Phenyl		-	
Heptyl		-	
Octyl		-	
6-Mercaptopurine		+	- (25)
Methotrexate		-	+ (22), - (23)
8-Azaguanine		-	- (6)
1-β-D-Arabinofuranosylcytosine		-	
Cyclocytidine		-	

Table 4
Results of mutation tests on antimetabolites

Antimetabolite	$\mu\text{g}/\text{plate}$	<i>his</i> ⁺ revertants/plate						<i>his</i> ⁺ revertants/nmol ^a
		TA100		TA98		TA92		
		+ S-9 mix	- S-9 mix	+ S-9 mix	- S-9 mix	+ S-9 mix	- S-9 mix	
5-FU ^b	1	183	82	30	13	33	19	0
	5	134	0	32	0	32	0	
	10	121	0	38	0	0	0	
	50	133	0	0	0	0	0	
FT-207	25	132	99	37	7	34	15	0
	50	131	101	14	1	21	13	
	250	87	0	0	0	22	9	
1-Methylcarbamoyl-5-FU	1	177	125	35	11	19	19	0
	5	153	0	49	0	21	0	
	10	173	0	27	0	17	0	
	50	0	0	0	0	6	0	
1-Ethylcarbamoyl-5-FU	2.5	145	125	27	14	26	16	0
	5	165	0	27	7	16	0	
	10	142	0	22	0	15	0	
	50	100	0	0	0	1	0	
1-Propylcarbamoyl-5-FU	2.5	158	133	28	10	20	12	0
	5	189	0	32	0	20	0	
	10	137	0	31	0	11	0	
	50	0	0	0	0	1	0	
1-Isopropylcarbamoyl-5-FU	0.5	129	103	14	9	26	18	0
	2.5	134	0	25	0	23	0	
	10	194	0	20	0	18	0	
	50	97	0	40	0	1	0	
1-Butylcarbamoyl-5-FU	0.5	162	154	24	21	26	17	0
	2.5	170	96	23	0	30	0	
	5	150	0	29	0	17	0	
	50	140	0	22	0	2	0	
1- <i>tert</i> -Butylcarbamoyl-5-FU	0.5	110	100	12	12	27	10	0
	2.5	126	0	21	0	28	0	
	10	142	0	24	0	30	0	
	50	160	0	0	0	3	0	
1-Pentylcarbamoyl-5-FU	0.5	170	118	36	21	21	13	0
	2.5	162	125	19	0	29	0	
	5	171	0	30	0	22	0	
	50	154	0	21	0	13	0	
1-Hexylcarbamoyl-5-FU	0.5	120	71	26	10	30	19	0
	2.5	107	0	29	0	22	0	
	5	114	0	16	0	27	0	
	10	148	0	24	0	3	0	
1-Cyclohexylcarbamoyl-5-FU	0.5	132	104	30	15	24	13	0
	1	149	115	37	11	25	0	
	5	146	0	35	0	23	0	
	10	138	0	31	0	2	0	
1-Phenylcarbamoyl-5-FU	0.5	121	110	27	9	25	13	0
	2.5	151	0	17	0	13	0	
	10	131	0	22	0	30	0	
	50	67	0	0	0	2	0	
1-Heptylcarbamoyl-5-FU	0.5	174	95	33	12	25	18	0
	2.5	146	0	19	0	17	0	
	10	151	0	29	0	17	0	
	50	0	0	0	0	12	0	

Table 4—Continued

Antimetabolite	$\mu\text{g}/\text{plate}$	<i>his</i> ⁺ revertants/plate						<i>his</i> ⁺ revertants/nmol ^a
		TA100		TA98		TA92		
		+ S-9 mix	- S-9 mix	+ S-9 mix	- S-9 mix	+ S-9 mix	- S-9 mix	
1-Octylcarbamoyl-5-FU	0.5	130	80	28	8	33	15	0
	2.5	135	0	22	0	31	0	
	10	130	0	31	0	13	0	
	50	0	0	0	0	11	0	
6-Mercaptopurine	250	114	276	22	20			0.06 ^c
	500	110	313	32	21			
	750	127	282	24	9			
	1,000	90	231	24	6			
Methotrexate ^d	104	142	65	36	33			0
	208	172	77	35	35			
	312	169	85	37	21			
8-Azaguanine	10	109	117	44	20	21	14	0
	25	126	139	37	0	36	19	
	50	123	127	36	0	33	18	
1- β -D-Arabinofuranosylcytosine	1,000	127	92	37	31			0
	5,000	145	134	30	30			
	10,000	158	125	28	22			
	25,000	135	132	34	19			
	30,000	131	141	28	20			
Cycloctidine	5,000	127	90	29	22			0
	10,000	91	77	19	25			
	20,000	132	105	31	37			
	25,000	150	135	38	22			
Control ^e		147	132	30	16	39	26	

^a The number of *his*⁺ revertants per nmol was calculated from the data obtained under the most active conditions in the linear dose-responsive range and a yield of 100 colonies more than the background yield.

^b 5-FU, 5-fluorouracil.

^c Tester strain TA100 without S-9 mix.

^d Drug contained some vehicle, and the amount of the drug was expressed as μg of active principle.

^e Average of values in 20 experiments. The numbers of revertants were: 117 to 183 of TA100 with S-9 mix, 89 to 169 of TA100 without S-9 mix, 19 to 38 of TA98 with S-9 mix, and 14 to 26 of TA98 without S-9 mix.

(28). ACNU, which is now being tested clinically, was mutagenic to TA100 with or without S-9 mix but was not mutagenic to TA98 at doses of up to 50 $\mu\text{g}/\text{plate}$. Carbazilquinone was mutagenic to TA100 with or without S-9 mix. Pipobroman was mutagenic to TA100 without S-9 mix but was not mutagenic to TA98.

Vinca Alkaloids. Results on *Vinca* alkaloids, including vinblastine sulfate and vincristine sulfate, are shown in Table 8. These compounds were not mutagenic to TA100 or TA98 with or without S-9 mix at doses of up to 500 $\mu\text{g}/\text{plate}$, as shown in Table 8. Moreover, a dose of 500 $\mu\text{g}/\text{plate}$ had no apparent lethal effect on the bacterial cells. These alkaloids were reported to inhibit the growth of cells in culture (21), but no data on their carcinogenicities are available.

Enzyme. L-Asparaginase was not mutagenic to either strain with or without S-9 mix (Table 8). Moreover, it had no lethal effect at doses of up to 3 mg/plate.

Adrenal Steroid Hormone. Prednisolone was not mutagenic to either strain with or without S-9 mix (Table 8), and it was not lethal at doses of up to 33 $\mu\text{g}/\text{plate}$. This compound was not tested at concentrations greater than 33 $\mu\text{g}/\text{plate}$.

In this work we tested the mutagenicities of various kinds

of antitumor agents. Some of them are now in clinical use, and some, such as various alkylcarbamoyl-5-fluorouracil derivatives, are now being tested in animals. Among 33 compounds tested 9 were found to be mutagenic; these were daunomycin hydrochloride, Adriamycin hydrochloride, mitomycin C, 6-mercaptopurine, cyclophosphamide, busulfan, ACNU, carbazilquinone, and pipobroman. The mutagenicities of daunomycin hydrochloride (3, 16), Adriamycin hydrochloride (3, 16), 6-mercaptopurine (3, 8), and cyclophosphamide (3, 8, 16) were reported, and the lack of mutagenicity found with methotrexate (3, 8) and actinomycin D (3) were confirmed in this study. Recently, Benedict *et al.* (3) reported that uracil mustard was mutagenic to TA1535 without S-9 mix. Uracil mustard has a bis(2-chloroethyl)amino group. This bis(2-chloroethyl)amino group may alkylate DNA and induce mutation. On the other hand, 5-fluorouracil and its alkylcarbamoyl derivatives seem to have no alkylating activity.

Certain substances used in cancer chemotherapy are endowed with mutagenic potential. There is a demonstrated correlation between mutagenicity in this bacterial test system and potential for causing cancer in laboratory animals. Therefore, the development of new chemotherapeutic agents that have no mutagenicity in such bacterial

Table 5
Trivial names, structures, mutagenicities, and carcinogenicities of alkylating agents

Trivial name	Structure	Mutagenicity	Carcinogenicity
Cyclophosphamide		+	+ (29)
Busulfan	$\text{CH}_3\text{-SO}_2\text{-O-(CH}_2\text{)}_4\text{-O-SO}_2\text{CH}_3$	+	+ (28)
ACNU		+	
Carbazilquinone		+	
Pipobroman		+	

Table 6
Results of mutation tests on alkylating agents

Agent	$\mu\text{g/plate}$	his^+ revertants/plate				his^+ revertants/ nmol^a
		TA100		TA98		
		+ S-9 mix	- S-9 mix	+ S-9 mix	- S-9 mix	
Cyclophosphamide ^b	70	158	90	16	13	0.26 (0, +) ^c
	175	309	110	27	5	
	525	437	106	22	2	
	700	401	133	25	0	
Busulfan	100	160	137	35	3	0.14 (0, +)
	500	390	267	24	4	
	750	578	578	35	7	
ACNU	1000	1132	1038	49	5	2.33 (0, -)
	5	140	182	30	21	
	10	178	218	28	23	
	25	234	262	23	23	
Carbazilquinone	50	500	366	38	30	7.48 (0, -)
	10	198	365	29	26	
	25	251	602	40	48	
Pipobroman	75	273	641	66	89	0.061 (0, -)
	100	317	420	29	85	
	250	127	176	35	31	
	500	147	218	35	29	
Control ^d	750	157	260	30	26	0.061 (0, -)
	1000	196	313	38	24	
		147	132	30	16	

^a The number of his^+ revertants per nmol was calculated from the data obtained under the most active conditions in the linear dose-responsible range and a yield of 100 colonies more than the background yield.

^b Drug contained some vehicle, and the amount of drug was expressed as μg of active principle.

^c 0, Tester strain TA100; +, with S-9 mix; -, without S-9 mix.

^d Average of values in 20 experiments. The numbers of revertants were: 117 to 183 of TA100 with S-9 mix, 89 to 169 of TA100 without S-9 mix, 19 to 38 of TA98 with S-9 mix, and 14 to 26 of TA98 without S-9 mix.

Table 7
Trivial names, structures, and mutagenicities of Vinca alkaloids, an enzyme, and an adrenal steroid hormone




Trivial name	Structure	Mutagenicity
Vinblastine sulfate Vincristine sulfate		- -
L-Asparaginase		-
Prednisolone		-

Table 8
Results of mutation tests on Vinca alkaloids, an enzyme, and an adrenal steroid hormone

	$\mu\text{g}/\text{plate}$	<i>his</i> ⁺ revertants/plate				<i>his</i> ⁺ revertants/nmol ^a
		TA100		TA98		
		+ S-9 mix	- S-9 mix	+ S-9 mix	- S-9 mix	
Vinblastine sulfate	100	124	102	25	9	0
	250	93	129	29	14	
	500	92	113	26	14	
Vincristine sulfate	100	113	121	32	18	0
	250	104	127	33	18	
	500	149	126	26	16	
L-Asparaginase ^b	500 (152 IU)	146	134	29	15	0
	1500 (455 IU)	148	156	23	26	
	3000 (909 IU)	165	135	25	23	
Prednisolone ^b	8.3	135	125	37	34	0
	17	145	145	39	33	
	33	123	144	35	23	
Control ^c		147	132	30	16	

^a The number of *his*⁺ revertants per nmol was calculated from the data obtained under the most active conditions in the linear dose-responsible range and a yield of 100 colonies more than the background yield.

^b Drug contained some vehicle, and the amount of the drug was expressed as μg of active principle.

^c Average of values in 20 experiments.

test systems would be most desirable because of the possibly decreased risk of latent drug-induced cancer in the patient undergoing chemotherapy.

REFERENCES

- Ames, B. N., McCann, J., and Yamasaki, E. Methods for Detecting Carcinogens and Mutagens with the *Salmonella*/mammalian-microsome Mutagenicity Test. *Mutation Res.*, 31: 347-364, 1975.
- Bell, W. R., Whang, J. J., Carbone, P. P., Breches, G., and Block, J. B. Cytogenetic and Morphologic Abnormalities in Human Bone Marrow Cells during Cytosine Arabinoside Therapy. *Blood*, 27: 771-781, 1966.
- Benedict, W. F., Baker, M. S., Haroun, L., Choi, E., and Ames, B. N. Mutagenicity of Cancer Chemotherapeutic Agents in the *Salmonella*/Microsome Test. *Cancer Res.*, 37: 2209-2213, 1977.
- Benedict, W. F., Harris, N., and Karon, M. Kinetics of 1- β -D-Arabinofuranosylcytosine-induced Chromosome Breaks. *Cancer Res.*, 30: 2477-2483, 1970.
- Haddow, A., Harris, R. J. C., Kon, G. A. R., and Roe, E. M. F. The Growth-inhibitory and Carcinogenic Properties of 4-Aminostilbene and Derivatives. *Phil. Trans. Roy. Soc. London Ser. A*. 241-247, 1948.
- Hadidian, Z., Fredrickson, T. N., Weisburger, E. K., Weisburger, J. H., Glass, R. M., and Mantel, N. Tests for Chemical Carcinogens. Report on the Activity of Derivatives of Aromatic Amines, Nitrosamines, Quinolines, Nitroalkanes, Amides Epoxides, Aziridines, and Purine Antimetabolites. *J. Natl. Cancer Inst.*, 41: 985-1036, 1968.
- Hampel, K. E., Kober, B., Rösch, D., Gerharts, H., and Meinig, K. H. The Action of Cytostatic Agents on the Chromosomes of Human Leukocytes *in Vitro*. *Blood*, 27: 816-823, 1966.
- Herbold, B., and Buselmaier, W. Induction of Point Mutations by Different Chemical Mechanisms in the Liver Microsomal Assay. *Mutation Res.*, 40: 73-84, 1976.
- Hoshi, A., Iigo, M., Yoshida, M., and Kuretani, K. Antitumor Activity of Carbamoyl Derivatives of 5-Fluorouracil by Oral Administration. *Gann*, 66: 673-674, 1975.
- Hoshi, A., Iigo, M., Yoshida, M., and Kuretani, K. Antitumor Activity of 1-Hexylcarbamoyl-5-fluorouracil in a Variety of Experimental Tumors. *Gann*, 67: 725-731, 1976.

11. Kihlman, B. A., Nichols, W. W., and Levan, A. The Effect of Deoxyadenosine and Cytosine Arabinoside on the Chromosomes of Human Leukocytes *in Vitro*. *Hereditas*, 50: 139-143, 1963.
12. Kirk, J. M. The Mode of Action of Actinomycin D. *Biochim. Biophys. Acta*, 42: 167-169, 1960.
13. Krugh, T. R. Association of Actinomycin D and Deoxyribonucleotides as a Model for Binding of the Drug to DNA. *Proc. Natl. Acad. Sci. U. S.*, 60: 1911-1914, 1972.
14. Marquardt, H., Philips, F. S., and Sternberg, S. S. Tumorigenicity *in Vivo* and Induction of Malignant Transformation and Mutagenesis in Cell Cultures by Adriamycin and Daunomycin. *Cancer Res.*, 36: 2065-2069, 1976.
15. McCann, J., and Ames, B. N. Detection of Carcinogens as Mutagens in the *Salmonella*/microsome Test: Assay of 300 Chemicals: Discussion. *Proc. Natl. Acad. Sci. U. S.*, 73: 950-954, 1976.
16. McCann, J., Choi, E., Yamasaki, E., and Ames, B. N. Detection of Carcinogens as Mutagens in the *Salmonella*/microsome Test: Assay of 300 Chemicals. *Proc. Natl. Acad. Sci. U. S.*, 72: 5135-5139, 1975.
17. McCann, J., Spingarn, N. E., Kobori, J., and Ames, B. N. Detection of Carcinogens as Mutagens: Bacterial Tester Strains with R Factor Plasmids. *Proc. Natl. Acad. Sci. U. S.*, 72: 979-983, 1975.
18. Müller, W. E. G., Yamazaki, Z., Sögtrop, H. H., and Zahn, R. K. Action of 1- β -D-Arabinofuranosylcytosine on Mammalian Tumor Cells-2. Inhibition of Mammalian and Oncogenic Viral Polymerases. *European J. Cancer*, 8: 421-428, 1972.
19. Murayama, I., and Otsuji, N. Mutation by Mitomycins in the Ultraviolet Light-sensitive Mutant of *Escherichia coli*. *Mutation Res.*, 18: 117-119, 1973.
20. Nagao, M., Yahagi, T., Seino, Y., Sugimura, T., and Ito, N. Mutagenicities of Quinoline and Its Derivatives. *Mutation Res.*, 42: 335-342, 1977.
21. Noble, R. L. Symposium on Vincalukoblastine (VLB). *Can. Cancer Conf.*, 4: 333-338, 1961.
22. Roshlaze, G., and Justus, J. Kanzerogene Wirkung von Methotrexat und Cyclophosphamid im Tierexperiment. *Deut. Gesundheitsw.*, 26: 219-222, 1971.
23. Rustia, M., and Shubik, P. Life-span Carcinogenicity Tests with 4-Amino- N^{10} -methylpteroylglutamic Acid (Methotrexate) in Swiss Mice and Syrian Golden Hamsters. *Toxicol. Appl. Pharmacol.*, 26: 329-338, 1973.
24. Sugimura, T., Sato, S., Nagao, M., Yahagi, T., Matsushima, T., Seino, Y., Takeuchi, M., and Kawachi, T. Overlapping of Carcinogens and Mutagens. *In*: P. N. Magee, S. Takayama, T. Sugimura, and T. Matsushima (eds.), *Fundamentals in Cancer Prevention*, pp. 191-215. Tokyo: University of Tokyo Press, 1976.
25. Sugiura, K., Teller, M. N., Parham, J. C., and Brown, G. B. A Comparison of the Oncogenicities of 3-Hydroxyxanthine, Guanine 3- N -Oxide, and Some Related Compounds. *Cancer Res.*, 30: 184-188, 1970.
26. Suzuki, H., Nagai, K., Yamaki, H., Tanaka, N., and Umezawa, H. On the Mechanism of Action of Bleomycin: Scission of DNA Strands *in Vitro* and *in Vivo*. *J. Antibiotics Tokyo Ser. A*, 22: 446-448, 1969.
27. Szybalski, W., and Iyer, V. N. The Mitomycins and Porfiriomycins. *In*: D. Gottlieb and P. D. Shaw (eds), *Antibiotics*, pp. 211-245. Berlin: Springer-Verlag, 1967.
28. Upton, A. C., Wolff, F. F., and Sniffen, E. P. Leukemogenic Effect of Myleran on the Mouse Thymus. *Proc. Soc. Exptl. Biol. Med.*, 108: 464-467, 1961.
29. Walker, S., and Bole, G. Augmented Incidence of Neoplasia in NZB/NZW Mice Treated with Long-term Cyclophosphamide. *J. Lab. Clin. Med.*, 82: 619-629, 1973.
30. Weisburger, J. H., Griswold, D. P., Prejean, J. D., Casey, A. E., Wood, H. B., and Weisburger, E. K. The Carcinogenic Properties of Some of the Principal Drugs Used in Clinical Cancer Chemotherapy. *Recent Results Cancer Res.*, 52: 1-17, 1975.
31. Zahn, R. K., Müller, W. E. G., Forster, W., Maidhoff, A., and Beyer, R. Action of 1- β -D-Arabinofuranosylcytosine on Mammalian Tumor Cells-1. Incorporation into DNA. *European J. Cancer*, 8: 391-396, 1972.
32. Zunino, F., Gambetta, R. A., and DiMarco, A. The Inhibition *in Vitro* of DNA Polymerase and RNA Polymerase by Daunomycin and Adriamycin. *Biochem. Pharmacol.*, 24: 309-311, 1975.
33. Zunino, F., Gambetta, R. A., and Zaccara, A. Interaction of Daunomycin with DNA. *Biochim. Biophys. Acta*, 277: 489-498, 1972.