

Control of Permeation of Bleomycin A₂ by Polyene Antibiotics in Cultured Chinese Hamster Cells¹

Shin-ichi Akiyama,² Katsuhiko Hidaka, Sohtarō Komiyama, and Michihiko Kuwano

Department of Biochemistry, Oita Medical School, Hazama-cho, Oita 879-56, Japan [S. A., K. H., M. K.], and Department of Otorhinolaryngology, Faculty of Medicine, Kyushu University, Fukuoka 812, Japan [S. K.]

ABSTRACT

Control of permeation of bleomycin A₂, a well-known antitumor antibiotic, in combination with various polyene macrolide antibiotics was analyzed in cultured Chinese hamster cells *in vitro*. Three polyene antibiotics, filipin, pentamycin, and pimaricin, were found to enhance the action of bleomycin A₂ remarkably, while amphotericin B or nystatin could not. Although DNA synthesis and colony-forming activity of polyene-sensitive Chinese hamster V79 cells were synergistically inhibited by the combination of filipin and bleomycin A₂, in a polyene-resistant subline (AMB^R-1) derived from V79, they were only slightly affected in the presence of both drugs. The cellular uptake of [¹⁴C]bleomycin A₂ by V79 was enhanced 2- to 4-fold in the presence of increasing doses of filipin or pentamycin, but not in the presence of amphotericin B. The treatment of V79 cells with filipin for 20 to 30 min was enough to block DNA synthesis almost completely when combined with 20 μg bleomycin A₂ per ml. The pretreatment of the hamster cells with 6 μg filipin per ml for 60 min continued to enhance the inhibitory action by bleomycin A₂ of DNA synthesis up to 5 hr after the removal of filipin from the cultured medium.

INTRODUCTION

Polyene macrolide antibiotics are known to enhance the permeation of combined second agents through interaction with sterols of cell membranes (1, 4, 9, 17), and the synergistic combination with polyenes is found to be a useful therapeutic method not only against fungi-infectious diseases (10) but also against tumors (15, 18). A polyene antibiotic, pentamycin, could potentiate bleomycin A₂, a potent antitumor agent (8, 19, 22), but not fusidic acid, an inhibitor of protein synthesis (7). In contrast, another polyene, amphotericin B, could potentiate fusidic acid (8, 16). In this report, we examine whether the cytotoxic action of bleomycin A₂ *in vitro* is enhanced by other polyene antibiotics besides pentamycin, and we discuss diversity among polyene-induced membranous alterations.

MATERIALS AND METHODS

Cell Culture and Cell Lines. Chinese hamster cell line V79 and its amphotericin B-resistant subline (AMB^R-1) were obtained and maintained as described previously (7). The cells were routinely grown in monolayers in glass Petri dishes in

minimal essential medium (Nissui Seiyaku Co., Tokyo, Japan) supplemented with 1 mg bacto-peptone per ml, 0.292 mg L-glutamine per ml, 10% fetal bovine serum (Microbiological Associates, Bethesda, Md.), and penicillin (100 units/ml). These cells could grow with a doubling time of 9 to 10 hr.

Chemicals. [³H]Thymidine (20 Ci/mmol) (New England Nuclear, Boston, Mass.) was used. Radioactive [¹⁴C]bleomycin A₂ (10 μCi/400 μg; Nippon Kayaku Co., Tokyo, Japan) was prepared and highly purified when bleomycin A₂ was methylated with ¹⁴C-containing methyl iodide at 42° in methanol, as described previously (5). Amphotericin B and nystatin were given to us by Sankyo Co., Tokyo, Japan, and pentamycin was given by Nikken Chemical Co., Tokyo, Japan. Filipin was kindly given by The Upjohn Co., Kalamazoo, Mich. Pimaricin was given by Torii Pharmaceutical Co., Tokyo, Japan. Nonradioactive bleomycin A₂ which does not contain other bleomycin molecules was kindly donated by Nippon Kayaku Co., and bleomycin A₂ was used as bleomycin throughout this study. Aquasol was purchased from New England Nuclear.

Synergistic Study *in Vitro* by [³H]Thymidine Incorporation. V79 and AMB^R-1 (1 to 2 × 10⁵ cells/ml) cultured in 3 ml medium in Petri dishes were exposed to various doses of bleomycin with or without each of the polyene antibiotics for 10 or 18 hr and were then exposed to [³H]thymidine (1 μCi/ml) for another 4 hr in the presence of drugs. After exposure to [³H]thymidine, DNA synthesis was measured by counting the radioactivity of 10% trichloroacetic acid-insoluble fraction retained on glass filter paper as described previously (13). Polyene antibiotics used in this study, including amphotericin B, pentamycin, pimaricin, filipin, and nystatin, were prepared by dissolving in dimethyl sulfoxide before each experiment, and all control experiments were done by adding the same amount of dimethyl sulfoxide alone.

Synergistic Study by Colony Formation. Cell survival of V79 or AMB^R-1 cells under synergistic conditions was also measured by colony formation. Cells (250 each) were plated in duplicate 60-mm dishes in the absence of any drug and incubated for 18 hr. Then the cells were exposed to bleomycin A₂ with or without filipin and incubated at 37° for 8 days. Colony number was counted when stained with Giemsa as described previously (7, 8).

Cellular Uptake of Radioactive Bleomycin. Cells were treated with each polyene antibiotic for 2 hr and then exposed to [¹⁴C]bleomycin A₂ (30,000 cpm/dish) for another 4 hr in the presence of the polyene. The cells in each dish were harvested and washed 3 times with phosphate-buffered saline containing 137 mM NaCl, 2.68 mM KCl, 8.1 mM Na₂HPO₄, 1.47 mM KH₂PO₄, and 1 mM bleomycin A₂ by repeated centrifugation. Finally, the cellular pellets were suspended in 1 ml of H₂O and mixed thoroughly with 10 ml of Aquasol. Then radioactivities were counted. Protein concentrations were determined by the

¹ This work was supported by a Grant-in-Aid from the Ministry of Education, Science and Culture, Japan.

² To whom requests for reprints should be addressed, at Department of Biochemistry, Oita Medical School, Hazama-cho, Oita 879-56, Japan.

Received February 15, 1979; accepted September 10, 1979.

method of Lowry *et al.* (16), with bovine serum albumin as a standard.

RESULTS

Enhancement of the Action of Bleomycin by Filipin, Pentamycin, and Pimaricin, but not by Amphotericin B and Nystatin. The Chinese hamster V79 cell is not highly sensitive to bleomycin (8), which is possibly due to difficulty in membrane permeability by bleomycin. The cytotoxic activity of bleomycin has been shown to be enhanced remarkably when combined with pentamycin in Chinese hamster V79 cells (8). To test if any other polyene antibiotics can potentiate bleomycin, we examined the effect of bleomycin in combination with 5 polyene antibiotics such as amphotericin B, filipin, nystatin, pentamycin, and pimaricin. We used 2 different concentrations of each polyene antibiotic with respect to inhibitory extent of DNA synthesis, low concentration of polyene (0 to 20% inhibition of DNA synthesis), and high concentration of polyene (40 to 60% inhibition). Cellular DNA synthesis was synergistically inhibited when 10 μg bleomycin A_2 per ml was combined with polyene antibiotics such as pentamycin, filipin, and pimaricin, but not when combined with amphotericin B or nystatin (Table 1). Bleomycin A_2 could not be potentiated even when combined with higher doses of amphotericin B or nystatin.

Synergistic Effect of Bleomycin and Filipin upon DNA Synthesis, as well as Colony Formation of Amphotericin B-sensitive and -resistant Cell Lines. Three polyene antibiotics, filipin, pentamycin, and pimaricin, effectively potentiate bleomycin in Chinese hamster cells, and the effect of pentamycin is in good agreement with our previous results (8, 19). To study an effective combination of bleomycin and filipin in more detail, we compared the synergistic effect between V79 and an amphotericin B-resistant (AMB^R-1) cell line derived from V79 (7). Dose response of V79 and AMB^R-1 cells to bleomycin was compared when combined with or without 3 or 6 μg filipin per ml (Chart 1). In the absence of filipin, both V79 and AMB^R-1 show almost identical dose-response curves to bleomycin, while addition of filipin was found to efficiently enhance the inhibition of DNA synthesis of V79, but not of AMB^R-1. DNA synthesis of V79 which is amphotericin B sensitive, was blocked by 70 to 75% when 5 to 10 μg bleomycin A_2 per ml was combined with 6 μg filipin per ml (Chart 1a). Almost the




Chart 1. Dose response to bleomycin in the absence (O) or presence (Δ, 3 $\mu\text{g}/\text{ml}$; ▲, 6 $\mu\text{g}/\text{ml}$) of filipin of V79 (a) and AMB^R-1 (b). Both V79 and AMB^R-1 cells were exposed to various doses of bleomycin- A_2 with or without filipin for 18 hr. Cellular DNA synthesis was determined as described in "Materials and Methods." Normalized activity of DNA synthesis from triplicate experiments is shown when 100% corresponds to 6741 cpm (V79) and 5454 cpm (AMB^R-1) in the absence of filipin, 6717 cpm (V79) and 5489 cpm (AMB^R-1) in the presence of filipin (3 $\mu\text{g}/\text{ml}$), and 5900 cpm (V79) and 4067 cpm (AMB^R-1) in the presence of filipin (6 $\mu\text{g}/\text{ml}$). Bars, S.E.

same synergistic effect of bleomycin and filipin as in DNA synthesis was observed when assayed by colony formation of V79 and AMB^R. As shown in Chart 2, bleomycin A_2 alone, below the concentration of 0.8 $\mu\text{g}/\text{ml}$, inhibited colony formation of both V79 and AMB^R cells by about 40 to 50% of the control activity. Colony-forming activity of V79, however, was decreased to below 10^{-2} of the control value by 0.3 μg bleomycin A_2 per ml, in combination with 3 μg filipin per ml (Chart 2a). In contrast, colony formation of AMB^R cells was blocked by 75% only when 0.8 μg bleomycin A_2 per ml was combined with 3 μg filipin per ml (Chart 2b). In addition, filipin (4 $\mu\text{g}/\text{ml}$) decreased the survival fraction of V79 to 7×10^{-3} of the control value when combined with 0.1 μg bleomycin A_2 per ml, while the same dose of the polyene antibiotic decreased that of AMB^R to 10^{-2} when combined with 0.8 μg bleomycin A_2 per ml. Thus, 2 independent assays (DNA synthesis and colony formation) reinforced an argument that filipin potentiated bleomycin A_2 and also that, to obtain the same degree of synergistic effect, AMB^R cells required about an 8-fold higher dose of bleomycin A_2 than did the V79 cells.

The differential effect of filipin and bleomycin A_2 upon V79 and AMB^R-1 cells was also confirmed by examining the dose response to filipin with or without bleomycin A_2 . More than 70% of DNA synthesis activity in V79 cells was inhibited by 10 μg bleomycin A_2 per ml in combination with 5 μg filipin per ml (Chart 3a), whereas DNA synthesis of AMB^R cells was inhibited only by 25% of the control when bleomycin A_2 (10 $\mu\text{g}/\text{ml}$) was combined with 20 μg filipin per ml (Chart 3b). The finding that AMB^R-1 cells are apparently resistant to filipin is consistent with our previous report (8). Since resistant phenotype of AMB^R to polyene antibiotics is due to the altered cholesterol content (7),³ it is suggested that the extent of synergistic effect of bleomycin and filipin is mediated through the level of cholesterol.

³ K. Hidaka, S. Akiyama, and M. Kuwano, submitted for publication.

Table 1

Synergistic effect of bleomycin and various polyene antibiotics upon DNA synthesis of V79 cells

Polyene antibiotics	Concentration (μg/ml)	cpm at bleomycin concentration of	
		0 μg/ml	10 μg/ml
None	0	2489 ± 120 ^a (100) ^b	2385 ± 122 (100)
Amphotericin B	20	2344 ± 140 (94)	2367 ± 153 (99)
	60	1115 ± 58 (45)	1125 ± 46 (47)
Nystatin	20	1921 ± 61 (77)	2125 ± 86 (89)
	60	1083 ± 59 (44)	982 ± 68 (41)
Filipin	6	2215 ± 160 (89)	546 ± 90 (23)
	15	1492 ± 54 (60)	28 ± 10 (1)
Pentamycin	1	2248 ± 293 (90)	897 ± 88 (38)
	2	1555 ± 64 (64)	29 ± 7 (1)
Pimaricin	50	2403 ± 196 (97)	120 ± 32 (5)
	75	1218 ± 45 (49)	119 ± 36 (5)

^a Mean ± SE of triplicate experiments.

^b Numbers in parentheses, DNA synthesis activity of polyenes alone, or of each combination of polyene and 10 μg bleomycin A_2 per ml when 100% corresponds to the activity in the absence (2489 cpm) and the presence (2385 cpm) of 10 μg bleomycin A_2 per ml, respectively.




Chart 2. Dose response of the parental cell V79 (a) and AMB^R-1 (b) to bleomycin A₂ in the absence or presence of filipin, assayed by colony-forming ability. Cells (250 each) of V79 and AMB^R-1 were plated and exposed to various concentrations of bleomycin A₂ alone (●) or in combination with filipin [1 (Δ), 2 (▲), 3 (○), or 4 (○) μg/ml] for 8 days. The curves drawn for each combination show the survival values obtained from triplicate experiments. Bars, S.E. Number of colonies in the absence of bleomycin A₂ with or without 1 to 4 μg filipin per ml was 150 to 216 in V79 (a) and 139 to 193 in AMB^R-1 (b).




Chart 3. Dose response of V79 (a) and AMB^R-1 (b) to filipin in the absence or presence of bleomycin A₂. After cells were exposed to various concentrations of filipin alone (○) or in combination with bleomycin A₂ (10 μg/ml) (●) for 18 hr, [³H]thymidine incorporation into acid-insoluble fraction was measured. The normalized activity of DNA synthesis is presented from triplicate experiments when 100% corresponds to 3825 cpm (V79) and 4253 cpm (AMB^R-1) in the absence of bleomycin A₂ and 3901 (V79) and 3785 cpm (AMB^R-1) in the presence of bleomycin A₂. Bars, S.E.

Duration of the Effect of Filipin to Enhance the Action of Bleomycin-A₂ in Chinese Hamster Cells. To examine when the synergistic effect of bleomycin A₂ and filipin appears after treatment of cells with polyene antibiotics, V79 cells were

exposed to 6 μg filipin per ml for various times, and then the culture medium was changed to remove filipin. At indicated times after the treatment, Bleomycin A₂ (20 μg/ml) was added immediately and was followed by incubation for another 10 hr. As shown in Chart 4, short time (20 to 30 min) exposure of V79 cells to filipin was enough to inhibit more than 80% of DNA synthesis when combined with bleomycin A₂. There appeared to be no inhibition of DNA synthesis in the absence of filipin.

We then examined how long the effect of filipin to potentiate the antitumor antibiotic continued after the removal of polyene antibiotics. The cells which were pretreated with filipin for 60 min were incubated in fresh medium and then 20 μg bleomycin A₂ per ml were added immediately (at Time 0) or at 1, 3, 5, 7, and 24 hr. It was found that the DNA synthesis of V79 cells was markedly inhibited by bleomycin A₂ even at 5 hr after the removal of filipin (Chart 5). The inhibition of DNA synthesis, however, could not be observed when V79 cells were cultured in the absence of the polyene for 7 hr or longer. It is concluded that the effect of filipin to enhance the action of bleomycin A₂ appears when V79 cells are exposed to the polyene briefly for 20 to 30 min and also that the cells treated with filipin could be susceptible to bleomycin A₂ during 5 hr after the removal of filipin.

Cellular Uptake of [¹⁴C]Bleomycin A₂ in the Presence of Polyene Antibiotics. Our previous study showed that a synergistic effect of bleomycin A₂ and polymyxin B, which is not a polyene antibiotic, could be partly due to enhancement of permeation of bleomycin A₂ by polymyxin B in cultured rat cells (14). In this work, we found that pentamycin, pimaricin, and filipin could potentiate bleomycin (Table 1). To test whether or not the synergistic effect is due to altered permeation, we assayed cellular uptake of bleomycin in the absence or presence of polyene antibiotics. Autoradiographic study with radioactive bleomycin *in vitro* showed that the radioactivity was observed near the nuclear membrane after exposure of the cultured cell for 4 hr (6). We therefore assayed the cellular




Chart 4. Exposure time to filipin and synergistic effect. V79 cells were exposed to filipin (6 μg/ml) for various times, and filipin was removed by changing medium at indicated time (●). Immediately after the removal of filipin, 20 μg bleomycin A₂ per ml were added, followed by incubation for 10 hr. Cellular DNA synthesis was determined from triplicate samples, as described in "Materials and Methods." Bars, S.E. As control (○), cells were cultured in the absence of filipin, but bleomycin A₂ alone was added at indicated times. Normalized activity of DNA synthesis is shown when 100% corresponds to 2063 cpm at time 0.




Chart 5. Duration of the effect of filipin to potentiate bleomycin A₂. V79 cells were exposed to 6 μg filipin per ml for 60 min (○), followed by incubation in fresh medium without filipin, and then 20 μg bleomycin A₂ per ml were added at indicated times. Cellular DNA synthesis was assayed by labeling with [³H]thymidine for 4 hr after the cells were incubated with bleomycin A₂ for 10 hr. As control (●), V79 cells were not treated with filipin, but 20 μg bleomycin A₂ per ml were added. Values are the mean of 3 samples. Bars, S.E.




Chart 6. Incorporation of [¹⁴C]bleomycin A₂ into V79 cells in the absence or presence of polyenes. Cells were treated with various doses of pentamycin (○), filipin (●), or amphotericin B (Δ). Incorporation of [¹⁴C]bleomycin A₂ was determined from triplicate samples, as described in "Materials and Methods." Bars, S.E.

uptake of bleomycin A₂ when V79 cells were incubated with [¹⁴C]bleomycin A₂ for 4 hr. As shown in Chart 6, the cellular uptake of [¹⁴C]bleomycin A₂ was stimulated 2- to 4-fold reproducibly in the presence of pentamycin or filipin. The percentage of uptake of bleomycin A₂ ranged from 0.42 to 0.59% of input count in the presence of 4 μg pentaene polyenes per ml, while the uptake was 0.17% in the absence of the polyene.

DISCUSSION

Among 5 polyene antibiotics (filipin, amphotericin B, nystatin, pentamycin, and pimaricin) examined, filipin, pentamycin, and pimaricin could remarkably potentiate an anticancer agent, bleomycin A₂ (23). In accordance with synergistic effects,

cellular uptake of radioactive bleomycin A₂ was enhanced by filipin or pentamycin, but not by amphotericin B. Since bleomycin A₂ was shown to induce breakdown of DNA chains *in vitro* (3, 20), enhanced permeation of the antitumor agent into the cells might cause reduction in cell survival. Bleomycin A₂, the structure of which was recently determined (21), may not be able to permeate the pore being formed by amphotericin B or nystatin (1, 4). One may assume that the membranous alteration induced by pentamycin or pimaricin is akin to that indicated by filipin, because these polyenes, filipin, pentamycin, and pimaricin, enhance the action of bleomycin A₂ remarkably. Recently, polyene antibiotics were classified into 2 groups: Group 1, antibiotics that caused K⁺ leakage and cell death (or hemolysis) at the same doses of added polyenes; and Group 2, antibiotics that caused K⁺ leakage at low doses and cell death (or hemolysis) at high doses (11). They found that Group 1 drugs included pimaricin (tetraene) or filipin (pentaene), while Group 2 included amphotericin B (heptaene) or nystatin (degenerated heptaene). Our work thus far appears to indicate that bleomycin A₂ is potentiated by Group 1 polyenes, but not by Group 2 antibiotics. In addition, amphotericin B or nystatin could make macrophages tumoricidal, but filipin and pimaricin, which are not ionophoretic, could not stimulate the tumoricidal activity of macrophages (2). It is interesting to note that the immunological study by Chapman and Hibbs (2) and our synergistic study thus far are consistent with the classification of polyene antibiotics, according to chemical structure and biological effects proposed by Kotler-Brajtbrug *et al.* (11).

Continuous incubation of animal cells with amphotericin B is necessary to enhance the cellular uptake of gigantic *Escherichia coli* DNA (12). We also found that the cells preexposed to amphotericin B lost their ability to potentiate the combined agents immediately after washing the cells once to remove the polyene.⁴ By contrast, the cells treated briefly with filipin could continue to enhance the effect of bleomycin A₂ during 5 to 6 hr, even in the absence of the polyene antibiotic (Chart 5). This result may be due to higher affinity of filipin than of amphotericin B to membranous sterols. Alternatively, some membranous damage caused by the pentaene antibiotic may not be repaired rapidly, presumably until new synthesis of membrane structure resumes.

On the other hand, bleomycin A₂ was well known to show differential response to various tumors in humans, and it was suggested that cellular level of inactivation enzyme of bleomycin A₂ was correlated with the sensitivity of tumors to the anticancer agent (23). However, differences in permeation of the antibiotic may be involved in the different responses of tumors to bleomycin A₂, as discussed previously in the bacterial system (24). Further study will be necessary to verify the latter possibility with mammalian cells or tumors.

ACKNOWLEDGMENTS

We thank K. Matsui for her technical assistance. We are deeply indebted to Dr. Y. Ikehara for the critical reading of this manuscript.

REFERENCES

1. Andreoli, T. E. The structure and function of amphotericin B cholesterol pores in lipid bilayer membranes. *Ann. N. Y. Acad. Sci.*, 235: 466-468, 1974.
2. Chapman, H. A., and Hibbs, J. B. Modulation of macrophage tumoricidal

⁴ S. Akiyama and M. Kuwano, unpublished data.

- capability by polyene antibiotics: support for membrane lipid as a regulatory determinant of macrophage function. *Proc. Natl. Acad. Sci. U. S. A.*, **75**: 4349-4353, 1978.
3. D'Andrea, A. D., and Haseltine, W. A. Sequence specific cleavage of DNA by the antitumor antibiotics neocarzinostatin and bleomycin. *Proc. Natl. Acad. Sci. U. S. A.*, **75**: 3608-3612, 1978.
 4. Dekruiff, B., and Demel, R. A. Polyene antibiotic sterol interactions in membranes of *Acholeplasma laidlawii* cells and lecithin liposomes. *Biochim. Biophys. Acta*, **339**: 57-70, 1974.
 5. Fujii, A., Takita, T., Maeda, K., and Umezawa, H. Chemistry of bleomycin. XI. The structures of the terminal amines. *J. Antibiot. (Tokyo)*, **26**: 398-399, 1973.
 6. Fujimoto, J. Radioautographic studies on the intracellular distribution of [¹⁴C]bleomycin in mouse tumor cells. *Cancer Res.*, **34**: 2969-2974, 1974.
 7. Hidaka, K., Endo, H., Akiyama, S., and Kuwano, M. Isolation and characterization of amphotericin B-resistant cell lines in Chinese hamster cells. *Cell*, **14**: 415-421, 1978.
 8. Hidaka, K., Matsui, K., Endo, H., Akiyama, S., and Kuwano, M. Differential control of synergistic effect with polyene macrolide antibiotics upon Chinese hamster cells *in vitro*. *Cancer Res.*, **38**: 4650-4653, 1978.
 9. Kinsky, S. C. Antibiotic interaction with model membranes. *Annu. Rev. Pharmacol.*, **10**: 119-142, 1970.
 10. Kobayashi, G. S., and Medoff, G. Antifungal agents: recent developments. *Ann. Rev. Microbiol.* **31**: 291-308, 1977.
 11. Kotler-Brajtburg, J., Medoff, G., Kobayashi, G. S., Boggs, S., Schlessinger, D., Pandey, R. C., and Rinehalt, K. L., Jr. Classification of polyene antibiotics according to chemical structure and biological effects. *Antimicrob. Agents Chemother.*, **15**: 716-722, 1979.
 12. Kumar, B. V., Medoff, G., Kobayashi, G., and Schlessinger, D. Uptake of *Escherichia coli* DNA in HeLa cells enhanced by amphotericin B. *Nature (Lond.)*, **250**: 323-325, 1974.
 13. Kuwano, M., Akiyama, S., Endo, H., and Kohga, M. Potentiation of fusidic acid and lentinan effects upon normal and transformed fibroblastic cells by amphotericin B. *Biochem. Biophys. Res. Commun.*, **49**: 1241-1248, 1972.
 14. Kuwano, M., Kamiya, T., Endo, H., and Komiyama, S. Potentiation of 5-fluorouracil, chromomycin A3 and bleomycin by amphotericin B or polymyxin B in transformed fibroblastic cells. *Antimicrob. Agents Chemother.*, **3**: 580-584, 1973.
 15. Laurent, G., Atassi, G., and Hildebrand, J. Potentiation of 1-(2-chloroethyl)-3-cyclohexyl-1-nitrosourea by amphotericin B in murine ependymoblastoma. *Cancer Res.*, **36**: 4069-4073, 1976.
 16. Lowry, O. H., Rosebrough, N. J., Farr, A. L., and Randall, R. J. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.*, **193**: 265-275, 1951.
 17. Medoff, G., Schlessinger, D., and Kobayashi, G. S. Sensitivity of transformed and nontransformed cells to amphotericin B and several rifamycin derivatives. *Cancer Res.*, **34**: 823-826, 1974.
 18. Medoff, G., Valenriote, F., Lynch, R., Schlessinger, D., and Kobayashi, G. S. Synergistic effect of amphotericin B and 1,3-bis(2-chloroethyl)-1-nitrosourea against a transplantable AKR leukemia. *Cancer Res.*, **34**: 974-978, 1974.
 19. Nakashima, T., Kuwano, M., Matsui, K., Komiyama, S., Hiroto, I., and Endo, H. Potentiation of bleomycin by antifungal polyene, pentamycin, in transformed animal cells. *Cancer Res.*, **34**: 3258-3261, 1974.
 20. Suzuki, H., Nagai, K., Yamaki, H., Tanaka, N., and Umezawa, H. On the mechanism of action of bleomycin: scission of DNA strands *in vitro* and *in vivo*. *J. Antibiot. (Tokyo) Ser. A*, **11**: 26-29, 1958.
 21. Takita, T., Muraoka, Y., Nakatani, T., Fujii, A., Umezawa, Y., Naganawa, H., and Umezawa, H. Chemistry of bleomycin. XIX. Revised structures of bleomycin and phleomycin. *J. Antibiot. (Tokyo)*, **31**: 801-804, 1978.
 22. Twentyman, P. R. The effect of bleomycin and pentamycin in combination on the survival of EMTB mouse tumor cells *in vivo*. *Br. J. Cancer*, **33**: 459-464, 1976.
 23. Umezawa, H. Bleomycin. *In*: J. W. Corcoran and F. E. Hahn (eds.), *Antibiotics*, Vol. 3, pp. 21-33. Berlin: Springer Verlag, 1975.
 24. Yamagami, H., Ishizawa, M., and Endo, H. Phenotypic and genetic characteristics of bleomycin sensitive mutants of *Escherichia coli*. *Gann*, **65**: 61-67, 1974.