

A Case-Cohort Study of Diet and Stomach Cancer¹

Po-Huang Chyou,² Abraham M. Y. Nomura, Jean H. Hankin, and Grant N. Stemmermann

Japan-Hawaii Cancer Study, Kuakini Medical Center, Honolulu, Hawaii 96817 [P. H. C., A. M. Y. N., G. N. S.], and Epidemiology Program, Cancer Research Center of Hawaii, Honolulu, Hawaii 96813 [J. H. H.]

ABSTRACT

In this case-cohort study, from 1965 to 1968, 8006 Hawaiian men of Japanese ancestry were interviewed with a 24-h dietary recall questionnaire. After a follow-up period of 18 years, 111 stomach cancer incident cases were identified. Dietary data from these patients and from 361 cancer-free men were analyzed for intake of selected foods, food groups, and nutrients. We found that the consumption of all types of vegetables was protective against stomach cancer. Specifically, subjects in the highest group of vegetable consumption (≥ 80 g/day) had a relative risk of 0.6 (95% confidence interval, 0.3-0.9) in comparison with nonconsumers. This statistically significant inverse trend persisted after adjustment for age at examination and cigarette-smoking status. Similar but weaker protective effects from consumption of green and cruciferous vegetables were also observed. In addition, an inverse association between stomach cancer risk and intake of fruits was noted ($P = 0.05$), but this inverse trend was weakened after the effect of cigarette smoking was taken into account. There were no other dietary factors significantly associated with the risk of gastric cancer.

INTRODUCTION

Researchers suspect that diet affects the risk of stomach cancer. Past dietary studies strongly suggest that a high intake of various types of vegetables is protective against stomach cancer (1-9). Epidemiological studies in different countries have also found that frequent ingestion of fruits (2, 6, 8, 10-12) and vitamin C (1, 2, 6, 11) is inversely associated with gastric cancer risk. On the other hand, the results from several other reports have not supported these findings (13-15).

Observational studies have also shown that increased risk of gastric cancer is associated with the high intakes of dried/salted fish (3, 13), pickled vegetables (3, 10, 13), cured meats (2, 11), and carbohydrate (7, 11, 16, 17). However, other investigators have not been able to substantiate these observations (5, 8, 9, 14, 17).

Due to the discrepant findings in past dietary investigations, we undertook the present study. Previously, we reported the results of a prospective study of stomach cancer and its relation to diet in our study population of Japanese men (15). At that time, a brief food frequency questionnaire obtained at time of examination was analyzed. We found that cases had more frequent intakes of pickles and cured meats but less frequent intakes of fruits and fried vegetables than did noncases. None of these differences was statistically significant. The computerized 24-h diet recall questionnaire data were limited to the intakes of major nutrients, so we were unable to report findings concerning the consumption of other specific foods or food groups recorded in the questionnaire. Subsequently, we recoded in detail the 24-h intake data from this population to assess further the relation of diet to gastric cancer risk by means of this case-cohort study.

Received 6/11/90; accepted 8/30/90.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ Supported by Grant R01 CA 33644 from the National Cancer Institute, NIH, Bethesda, MD.

² To whom requests for reprints should be addressed, at Japan-Hawaii Cancer Study, Kuakini Medical Center, 347 N. Kuakini Street, Honolulu, HI 96817.

MATERIALS AND METHODS

The population base for this study consisted of 8006 American men of Japanese ancestry who were born from 1900 to 1919. The median year of birth was 1913. They were interviewed and examined from 1965 to 1968 on the Hawaiian island of Oahu. Since then, newly diagnosed cases of stomach cancer have been identified through continuous surveillance of the major Oahu hospitals. To reduce the possibility of missing incident cases, computer linkage was periodically done with data files from the Hawaii Tumor Registry, which is a population-based cancer registry and a member of the Surveillance, Epidemiology, and End Results Program of the National Cancer Institute. From 1966 to July 1983, 115 incident stomach cancer cases [International Classification of Diseases, Rev. 8, code 151] with histological tissue confirmation of their diagnoses were identified. Based on a separate 19-year follow-up survey of the study subjects since their examination, we found that only 1.3% of the men could not be located on Oahu. Consequently, the surveillance for incident cases of stomach cancer should be nearly complete.

To identify a suitable control group for this investigation, we chose a random sample of 391 cancer-free men to represent the large cohort, using the case-cohort study design approach (18).

Four cases and 30 noncases were excluded because their previous 24-h dietary intake was self-reported as atypical, leaving 111 stomach cancer cases and 361 noncases for further data analysis.

Based on the Lauren histopathological classification (19), there were 83 cases with the intestinal-mixed-other type, 23 with the diffuse type, and 5 with unknown type. Because of their small number, the diffuse cases were not analyzed separately.

A 24-h dietary recall questionnaire was administered at the time of examination of the cohort subjects. The interviewing and coding methods have been described in detail elsewhere (20). Briefly, to prepare the recall data for analysis at the time of examination, a set of precoded forms was used in transcribing the 24-h dietary information. The nutritionists tallied the consumption frequencies of predefined units of 54 common food items grouped on the form on the basis of similarity of macronutrients. For example, vegetables were grouped as starchy, cooked, or raw and coded according to number of half-cup portions, while beef items were grouped and coded as number of ounces eaten. For specific foods other than the 54 items that were consumed over the past 24 h, the nutritionists recorded the data and calculated by hand the nutrients consumed. The original nutrient data bank included only calories, carbohydrate, fat, protein and their components, cholesterol, and sodium.

For the present study, we recoded the 24-h recalls, recording actual items consumed such as tomatoes, carrots, oranges, etc. This was done so that specific foods and food groups, as well as the intakes of vitamins and minerals, could be analyzed, using our updated comprehensive food consumption data base. In this data base, information for more than 1000 items, 50 nutrients, and other dietary components is available in 100-g units. Sources include United States Department of Agriculture Handbook 8 (21), United States Department of Agriculture Handbook supplements (22), food composition tables from Japan (23, 24), and other publications.

The age-adjusted mean intakes of selected nutrients and various generic food groups were calculated for cases and noncases. A test based on one-way, unbalanced analysis of covariance methods was used to determine whether there was a statistically significant difference ($P \leq 0.05$) between cases and noncases in mean intakes of individual nutrients and food groups (25).

For the food groups which showed statistically significant or border-

Table 1 Age-adjusted mean daily intake of selected nutrients for stomach cancer cases and noncases

Nutrient (units/day)	Cases (n = 111)		Noncases (n = 361)		P value ^a
	Mean	SE	Mean	SE	
Calories	2126.4	60.1	2082.2	33.4	0.52
Carbohydrate (g)	248.4	7.8	251.5	4.3	0.73
Fat (g)	76.1	3.2	72.9	1.8	0.37
Protein (g)	89.1	3.1	88.7	1.7	0.92
Sodium (g)	3.0	0.2	3.1	0.1	0.66
Vitamin A (IU)	6155.2	1469.6	7794.9	814.9	0.33
Retinol (μg)	672.0	156.3	477.8	86.7	0.28
β-Carotene (μg)	1944.4	695.4	3086.8	385.6	0.15
Other carotene (μg)	802.7	249.8	1261.0	138.5	0.11
Vitamin C (mg)	101.3	9.3	114.5	5.2	0.21
Vitamin D (IU)	27.6	6.1	36.5	3.4	0.20
Vitamin E (IU)	6.2	0.4	6.3	0.2	0.76
Calcium (mg)	607.0	33.3	576.5	18.4	0.42

^a To compare the two mean values, adjustment for age at the time of examination was by analysis of covariance.

line ($P = 0.07$) mean differences, we separately estimated the RR³ and CI values of stomach cancer for consumers compared with nonconsumers by the proportional hazards regression approach (26). All RR values were adjusted for age at examination. We also evaluated the RR and CI of stomach cancer with simultaneous adjustments for age and current smoking status (yes/no), since current smoking was a significant risk factor for stomach cancer in our cohort (15). However, there was no dose-response effect of cigarette smoking based on number of cigarettes per day or years of smoking among current smokers. In addition, likelihood ratio tests were used to evaluate the linear trend associations between continuous dietary intake variables and stomach cancer risk. Finally, parameters for the proportional hazards regression were estimated by a method of maximum likelihood (27).

RESULTS

The mean age at time of examination was 57.7 years for the 111 gastric cancer cases and 57.6 years for the 361 noncases. Seventy-eight % of the cases and 81 % of the noncases were born in Hawaii or elsewhere in the United States. Of the 111 cancer cases, 33 (29.7%) were diagnosed from 1966 to 1973, another 37 (33.3%) were identified from 1974 to 1978, and the remaining 41 (37%) were diagnosed during the last 5 years of follow-up from 1979 to 1983.

The age-adjusted mean intakes of selected nutrients for cases and noncases are listed in Table 1. The cases consumed smaller amounts of β- and other carotenes, vitamin C, and vitamin D than did noncases, but none of the differences were statistically significant at $P \leq 0.05$. The intakes of vitamin A and retinol did not differ markedly between the two groups of subjects. They were also similar in their consumption of carbohydrate, fat, protein, and sodium.

Table 2 summarizes the comparison between cases and noncases in their mean intakes of selected food groups, as well as their intakes of nitrate, nitrite, and dimethylnitrosamine. The cases consumed fewer vegetables and fruits than the noncases. Statistically significant differences were observed in the intakes of all vegetables ($P < 0.01$) and green vegetables ($P = 0.02$). There were also borderline differences in the consumption of cruciferous vegetables and all fruits. The cases and noncases did not differ greatly in their consumption of dried fish, processed meats, pickles, grains, and nitrate-related compounds.

The relative risks by increasing amounts of consumption of different vegetables are shown in Table 3. There was a statistically significant inverse trend in the age-adjusted relative risk

Table 2 Age-adjusted mean daily dietary intake of selected food groups and nitrate-related compounds for stomach cancer cases and noncases

Food group (units/day)	Cases (n = 111)		Noncases (n = 361)		P value ^a
	Mean	SE	Mean	SE	
All vegetables (g)	45.3	8.2	76.8	4.5	<0.01
Green vegetables (g)	33.0	5.8	48.4	3.2	0.02
Cruciferous vegetables (g)	10.4	4.3	19.9	2.4	0.06
Yellow vegetables (g)	4.2	5.7	15.2	3.2	0.09
All fruits (g)	174.2	21.2	218.4	11.7	0.07
Yellow fruits (g)	86.7	12.6	101.1	7.0	0.32
Citrus fruits (g)	52.9	12.0	58.1	6.6	0.70
Dried fish (g)	0.4	1.0	1.3	0.5	0.44
Processed meats (g)	32.7	4.0	26.5	2.2	0.18
Pickles (g)	15.7	2.4	15.6	1.3	0.97
Breakfast cereals (g)	9.8	4.5	11.3	2.5	0.77
Grains: breads, rice, pasta, crackers (g)	472.1	24.2	468.0	13.4	0.88
Legumes, seeds, nuts (g)	22.4	6.4	28.5	3.5	0.40
Nitrate (mg)	78.2	8.1	86.7	4.5	0.36
Nitrite (mg)	2.1	0.1	2.1	0.1	0.94
Dimethylnitrosamine (μg)	0.8	0.1	0.6	0.1	0.10

^a To compare the two mean values, adjustment for age at the time of examination was by analysis of covariance.

Table 3 Adjusted relative risk of stomach cancer by level of intake of all vegetables, green vegetables, and cruciferous vegetables

Cutoff points for daily intakes were as follows. All vegetables: 0, none; <40, low; <80, medium; 80+, high. Green vegetables: 0, none; <30, low; <60, medium; 60+, high. Cruciferous vegetables: 0, none; >0, consumer.

Intake of vegetables (g/day)	No. of cases	No. of noncases	Age-adjusted		Age- and smoking-adjusted ^a	
			RR	CI	RR	CI
All vegetables						
None ^b	38	107	1.0		1.0	
Low	27	61	1.2	0.7-1.9	1.2	0.8-2.0
Medium	20	61	0.9	0.5-1.6	0.9	0.5-1.5
High	26	132	0.6	0.3-0.9	0.7	0.4-1.1
P value for linear trend			<0.001		0.001	
Green vegetables						
None ^b	43	137	1.0		1.0	
Low	22	54	1.2	0.7-2.0	1.3	0.8-2.2
Medium	25	58	1.3	0.8-2.1	1.4	0.9-2.3
High	21	112	0.6	0.4-1.0	0.7	0.4-1.2
P value for linear trend			0.01		0.06	
Cruciferous vegetables						
None ^b	95	282	1.0		1.0	
Consumer	16	79	0.6	0.4-1.0	0.7	0.4-1.2
P value for linear trend			0.03		0.07	

^a Current smoking status: yes/no.

^b Referent group.

of stomach cancer with the intake of green, cruciferous, and all vegetables. However, after adjustment for current smoking status, only the association with all vegetables remained significant ($P = 0.001$).

The relative risk of gastric cancer by increasing intake of all fruits is presented in Table 4. There was an inverse linear trend in risk that was statistically significant ($P = 0.05$), but it did not persist after adjustment for cigarette smoking.

The analyses in Tables 3 and 4 were repeated for the 83 intestinal-mixed-other histopathological cases of stomach cancer. The results were similar. The relative risks were 0.8 for the high consumers of all vegetables (for linear trend, $P = 0.01$) and 0.6 for the high consumers of all fruits (for linear trend, $P = 0.06$) with adjustment for age at the time of examination and current smoking status.

To determine whether the high consumption of both vegetables and fruits had an additive protective effect, the subjects were classified into four categories in which those with less than the median intake of both foods were chosen as the baseline

³ The abbreviations used are: RR, relative risk; CI, 95% confidence interval.

Table 4 Adjusted relative risk of stomach cancer by level of intake of all fruits
Cutoff points for daily intake of all fruits were as follows: 0, none; ≤ 150 , low; ≤ 300 , medium; 301+, high.

Intake of all fruits (g/day)	No. of Cases	No. of Noncases	Age-adjusted		Age- and Smoking- ^a adjusted	
			RR	CI	RR	CI
None ^b	37	96	1.0		1.0	
Low	29	84	0.9	0.6–1.5	1.0	0.6–1.6
Medium	23	80	0.8	0.5–1.3	0.9	0.5–1.4
High	22	101	0.6	0.4–1.0	0.8	0.4–1.3
P value for linear trend			0.05		0.20	

^a Current smoking status: yes/no.

^b Referent group.

group. The three relative risks of the subjects with a high intake of either vegetables or fruits or both were identical at 0.7 with adjustment for age at the time of examination and current smoking status.

DISCUSSION

The consumption of all types of vegetables was found to be protective against stomach cancer in this investigation. The highest consumers (≥ 80 g/day) of vegetables had an age-adjusted relative risk of 0.6 (CI 0.3–0.9) compared with nonconsumers. This statistically significant inverse association persisted after the effect of current smoking was taken into account. Weaker, but similar, associations were also observed for the intakes of cruciferous vegetables and green vegetables.

Cabbage (five types) constituted 77% of the total amount of cruciferous vegetables consumed by study subjects. In addition, the combination of cabbage and lettuce accounted for 67% of the green vegetables consumed, whereas cabbage, lettuce, and tomatoes accounted for 60% of all vegetables eaten. The subjects consumed 43 different raw or cooked vegetables. The predominance of cabbage and lettuce in the diet helps to explain the similar results for the vegetable categories.

Evidence of an inverse association between stomach cancer and the intake of vegetables has been supported by a number of earlier studies. You *et al.* (6) in China recently reported that the risk of gastric cancer was reduced in frequent consumers of vegetables. Similar results were observed in Greece (9), Poland (8), and the United States (2, 4). La Vecchia *et al.* (11) in Italy found an inverse association of gastric cancer with green vegetable consumption, which was supported by a prospective study of green-yellow vegetable intake in Japan (28). Risch *et al.* (1) also reported a negative association with both the intakes of green and cruciferous vegetables in Canada. However, other investigators could not confirm the inverse association of gastric cancer with green-yellow vegetable intake in Japan (10, 13) or with cruciferous vegetable consumption in Italy (11).

Some of the discrepancies in the findings from separate studies could be due to intercountry differences in the varieties of vegetables consumed, the methods of consumption (raw or cooked), the number of specific vegetable items included in the individual studies, or the validity of the dietary methods. If vegetables are protective against stomach cancer, then it would be helpful to identify the specific ones that have the greatest effect. Dark green vegetables contain predominantly lutein and β -carotene, while tomatoes are rich in lycopene (29). β -Carotene is the major carotenoid in carrots and other deep yellow-orange vegetables (29). Carrots were not consumed to a great extent

among our subjects, and their low intake may help to explain the lack of a significant association between β -carotene and stomach cancer in this study (Table 1). Cruciferous vegetables contain certain indoles and isothiocyanates which have been found to inhibit carcinogenesis by polycyclic aromatic hydrocarbons in the forestomach of experimental animals (30). Other investigators have found that the mutagen-depressing effect of vegetables was markedly weakened by cooking (31), so the form in which vegetables are consumed could also be very important. Thus, any number of components found in vegetables could have a protective effect against gastric cancer.

An inverse association was also noted between stomach cancer risk and intake of all types of fruit. Subjects whose daily fruit intake exceeded 300 g had an age-adjusted relative risk of 0.6 (CI 0.4–1.0) compared with nonconsumers, but the trend was only of borderline statistical significance. The association diminished too with adjustment for current smoking status. The correlation (Spearman's) coefficient between fruit intake and pack-years of cigarette smoking was -0.17 , indicating fruit intake and cigarette smoking are weakly related. Earlier studies done in five countries all showed a protective effect of fruit intake against stomach cancer (2, 6, 8, 10, 11). Moreover, two of these studies (8, 11) found a clear dose-response relationship. Yet, other investigators (3, 9, 13) did not support this finding.

The subjects in our study consumed 54 different fruits. Papayas, oranges, apples, guavas, and mangos accounted for 66% of the total amount of fruits consumed. These fruits (except apples) are rich in vitamin A or vitamin C. Vitamin A has been found to be protective against tumor promotion in laboratory animals (32), and vitamin C inhibits the formation of *N*-nitroso compounds (33), which have been suspected to play a role in gastric carcinogenesis (34). Noncases consumed more vitamin A, β - and other carotenes, and vitamin C than cases in our study. However, none of these nutrients was found to be significantly protective against stomach cancer ($P > 0.10$).

Although this study has the advantage of obtaining the dietary data prospectively before the cases of stomach cancer were identified, one limitation is that the dietary intake was based on only a 24-h diet recall. A 7-day diet record completed by 329 men in the cohort was done to assess the within-person variability of the 24-h data. Similar levels of nutrient intakes were recorded by both methods (35). Thirty-four subjects, whose dietary intake on the previous day was atypical, were excluded from the study to improve the representativeness of the 24-h dietary recall. Nonetheless, the possibility exists that intraindividual variation in daily diet could decrease the statistical power to detect specific differences of the nutrient intakes between the two groups of subjects. Consequently, associations found in this study may actually be stronger than reported.

The consumption of dried fish, processed meats, pickles, and foods rich in nitrate-related compounds was relatively low in these subjects. This low consumption in persons, aged 45–68 years at time of interview, may have partially contributed to the lack of any association of these foods with stomach cancer risk. Some past studies have found that persons who consume foods high in precursors of *N*-nitroso compounds are at increased risk for gastric cancer (1, 3, 34, 36–38), although other investigations have not supported this association (5, 8–10, 14, 17).

In conclusion, the results of our investigation support the potential importance of vegetable and probably fruit intake in reducing the risk of stomach cancer. Further research is required to identify the specific vegetables and their components which may be protective agents. Furthermore, we would rec-

ommend that future epidemiological studies in populations at high risk of developing gastric cancer be conducted. Quantitative diet histories covering a longer time should be included in the study design to estimate the usual intakes of foods associated with gastric cancer risk.

REFERENCES

- Risch, H. A., Jain, M., Choi, N. W., Fodor, J. G., Pfeiffer, C. J., Howe, G. R., Harrison, L. W., Craib, K. J. P., and Miller, A. B. Dietary factors and the incidence of cancer of the stomach. *Am. J. Epidemiol.*, *122*: 947-959, 1985.
- Correa, P., Fontham, E., Pickle, L. W., Chen, V., Lin, Y., and Haenszel, W. Dietary determinants of gastric cancer in south Louisiana inhabitants. *J. Natl. Cancer Inst.*, *75*: 645-654, 1985.
- Haenszel, W., Kurihara, M., Segi, M., and Lee, R. K. C. Stomach cancer among Japanese in Hawaii. *J. Natl. Cancer Inst.*, *49*: 969-988, 1972.
- Graham, S., Schotz, W., and Martino, P. Alimentary factors in the epidemiology of gastric cancer. *Cancer (Phila.)*, *30*: 927-938, 1972.
- Haenszel, W., Kurihara, M., Lock, F. B., Shimizu, H., and Segi, M. Stomach cancer in Japan. *J. Natl. Cancer Inst.*, *56*: 265-274, 1976.
- You, W., Blot, W. J., Chang, Y., Ershow, A. G., Yang, Z., An, Q., Henderson, B., Ku, G., Fraumeni J. F., and Wang, T. Diet and high risk of stomach cancer in Shandong, China. *Cancer Res.*, *48*: 3518-3523, 1988.
- Hu, J., Zhang, S., Jia, E., Wang, Q., Liu, S., Liu, Y., Wu, Y., and Cheng, Y. Diet and cancer of the stomach: a case-control study in China. *Int. J. Cancer*, *41*: 331-335, 1988.
- Jedrychowski, W., Wahrendorf, W., Popiela, T., and Rachtan, J. A case-control study of dietary factors and stomach cancer risk in Poland. *Int. J. Cancer*, *37*: 837-842, 1986.
- Trichopoulos, D., Ouranos, G., Day, N. E., Tzonou, A., Manousos, O., Papadimitriou, C., and Trichopoulos, A. Diet and cancer of the stomach: a case-control study in Greece. *Int. J. Cancer*, *36*: 291-297, 1985.
- Kono, S., Ikeda, M., Tokudome, S., and Kuratsune, M. A case-control study of gastric cancer and diet in northern Kyushu, Japan. *Jpn. J. Cancer Res. (Gann)*, *79*: 1067-1074, 1988.
- La Vecchia, C., Negri, E., Decarli, A., Avanzo, B. D., and Franceschi, S. A case-control study of diet and gastric cancer in northern Italy. *Int. J. Cancer*, *40*: 484-489, 1987.
- Correa, P., Cuello, C., Fajardo, L. F., Haenszel, W., Bolanos, O., and Ramirez, B. Diet and gastric cancer: nutrition survey in a high risk area. *J. Natl. Cancer Inst.*, *70*: 673-678, 1983.
- Tajima, K., and Tominaga, S. Dietary habits and gastrointestinal cancers: a comparative case-control study of stomach and large intestinal cancers in Nagoya, Japan. *Jpn. J. Cancer Res. (Gann)*, *76*: 705-716, 1985.
- Acheson, E. D., and Doll, R. Dietary factors in carcinoma of the stomach: a study of 100 cases and 200 controls. *Gut*, *5*: 126-131, 1964.
- Nomura, A., Grove, J., Stemmermann, G. N., and Severson, R. K. A prospective study of stomach cancer and its relation to diet, cigarettes, and alcohol consumption. *Cancer Res.*, *50*: 627-631, 1990.
- Graham, S., Haughey, B., Marshall, J., Brasure, J., Zielezny, M., Freudenheim, J., West, D., Nolan, J., and Wilkinson, G. Diet in the epidemiology of gastric cancer. *Nutr. Cancer*, *13*: 19-34, 1990.
- Modan, B., Lubin, F., Barell, V., Greenberg, R. A., Modan, M., and Graham, S. The role of starches in the etiology of gastric cancer. *Cancer (Phila.)*, *34*: 2087-2092, 1974.
- Prentice, R. L. A case-cohort design for epidemiologic cohort studies and disease prevention trials. *Biometrika*, *73*: 1-12, 1986.
- Lauren, P. The two histological main types of gastric carcinoma. Diffuse and so-called intestinal-type. *Acta Pathol. Microbiol. Scand.*, *64*: 31-49, 1965.
- Tillotson, J. L., Kato, H., Nichaman, M. Z., Miller, D. C., Gay, M. L., Johnson, K. G., and Rhoads, G. G. Epidemiology of coronary heart disease and stroke in Japanese men living in Japan, Hawaii, and California: methodology for comparison of diet. *Am. J. Clin. Nutr.*, *26*: 177-184, 1973.
- Composition of Foods. . . Raw, Processed, Prepared. Data set 8-1-1. Hyattsville, MD: U. S. Department of Agriculture, 1972.
- US Department of Agriculture. Composition of Foods. . Raw, Processed, Prepared. Agriculture Handbook Suppl. 8-1: Dairy and Egg Products (1976); 8-4: Fats and Oils (1979); 8-5: Poultry Products (1979); 8-6: Soups, Sauces, and Gravies (1980); 8-7: Sausages and Luncheon Meats (1980); 8-8: Breakfast Cereals (1982); 8-9: Fruits and Fruit Juices (1982); 8-10: Pork Products (1983). Washington, DC: Government Printing Office, 1976-1983.
- Science and Technology Agency, Resources Survey Group. Table on Components of Japanese Foods, Ed. 3. Tokyo: Ishiyaku, 1980.
- Japan Dietetic Association. Standard Tables of Food Composition, Rev. 3. Tokyo: Daiichi Shuppan, 1964.
- Freund, R. J., and Littell, R. C. Statistical Analysis System (SAS) for Linear Models. Cary, NC: SAS Institute Inc., 1981.
- Cox, D. R. Regression models and life tables (with discussion). *J. R. Stat. Soc. B*, *34*: 187-220, 1972.
- Harrell, F. The PHGLM procedure. *In: SUGI Supplemental Library User's Guide*, pp. 267-294. Cary, NC: SAS Institute Inc., 1983.
- Hirayama, T. Epidemiology of stomach cancer. *Jpn. J. Cancer Res. (Gann)*, *11*: 3-19, 1971.
- Micozzi, M. S., Beecher, G. R., Taylor, P. R., and Khachik, F. Carotenoid analyses of selected raw and cooked foods associated with a lower risk for cancer. *J. Natl. Cancer Inst.*, *82*: 282-285, 1990.
- Wattenberg, L. W. Inhibitors of chemical carcinogenesis. *Adv. Cancer Res.*, *26*: 197-225, 1977.
- Kamiyama, S., and Michioka, O. Mutagenic components of diets in high- and low-risk areas for stomach cancer. *In: H. F. Stich (ed.), Carcinogens and Mutagens in the Environment*, pp. 29-42. Cleveland, OH: CRC Press, 1983.
- Bertram, J. S., Kolonel, L. N., and Meyskens, F. L. Rationale and strategies for chemoprevention of cancer in humans. *Cancer Res.*, *47*: 3012-3031, 1987.
- Mirvish, S. S., Wallcave, L., Eagen, M., and Shubik, P. Ascorbate-nitrite reaction: possible means of blocking the formation of carcinogenic *N*-nitroso compounds. *Science (Washington DC)*, *177*: 65-67, 1972.
- Mirvish, S. S. The etiology of gastric cancer: intragastric nitrosamide formation and other theories. *J. Natl. Cancer Inst.*, *71*: 629-647, 1983.
- McGee, D., Rhoads, G., Hankin, J., Yano, K., and Tillotson, J. Within-person variability of nutrient intake in a group of Hawaiian men of Japanese ancestry. *Am. J. Clin. Nutr.*, *36*: 657-663, 1982.
- Howson, C. P., Hiyama, T., and Wynder, E. L. The decline in gastric cancer: epidemiology of an unplanned triumph. *Epidemiol. Rev.*, *8*: 1-27, 1986.
- Nomura, A. Stomach. *In: D. Schottenfeld and J. Fraumeni, Jr. (eds.), Cancer Epidemiology and Prevention*, pp. 624-637. Philadelphia: W. B. Saunders, 1982.
- Forman, D. Dietary exposure to *N*-nitroso compounds and the risk of human Cancer. *Cancer Surv.*, *6*: 719-738, 1987.