



Inequity in uptake of hospital-based childbirth care in rural Tanzania: analysis of the 2015–16 Tanzania Demographic and Health Survey

Manuela Straneo¹, Lenka Benova^{2,3}, Claudia Hanson^{4,5,*}, Piera Fogliati⁶, Andrea B Pembe⁷, Tom Smekens⁸ and Thomas van den Akker^{1,9}

¹Athena Institute, VU Amsterdam, De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands

²Sexual and Reproductive Health Group, Department of Public Health, Institute of Tropical Medicine, Nationalestraat 155, 2000 Antwerp, Belgium

³Faculty of Epidemiology and Population Health, LSHTM, Keppel Street, London WC1E 7HT, UK

⁴Karolinska Institutet, 171 77 Stockholm, Sweden

⁵Faculty of Infectious and Tropical Diseases, LSHTM, Keppel Street, London WC1E 7HT, UK

⁶Doctors with Africa-CUAMM, Av. Mártires da Machava n.º 859 R/C, Cidade de Maputo, Moçambique

⁷Department of Obstetrics and Gynecology, Muhimbili University of Health and Allied Sciences, PO Box 65001, Dar es Salaam, United Republic of Tanzania

⁸Department of Public Health, Institute of Tropical Medicine, Nationalestraat 155, 2000 Antwerp, Belgium

⁹Department of obstetrics and Gynecology, Leiden University Medical Center, Rapenburg 70, 2311 EZ Leiden, The Netherlands

*Corresponding author. Health Systems and Policy, Global Public Health, Karolinska Institutet, K9 Global folkhälsa, K9 GH Stålsby Lundborg Hanson, Stockholm 171 77, Sweden. E-mail: claudia.hanson@ki.se

Accepted on 22 June 2021

Abstract

Proportions of facility births are increasing throughout sub-Saharan Africa, but obstetric services vary within the health system. In Tanzania, advanced management of childbirth complications (comprehensive emergency obstetric care) is offered in hospitals, while in frontline, primary health care (PHC) facilities (health centres and dispensaries) mostly only routine childbirth care is available. With over half (54%) of rural births in facilities, we hypothesized the presence of socio-economic inequity in hospital-based childbirth uptake in rural Tanzania and explored whether this relationship was modified by parity. This inequity may compound the burden of greater mortality among the poorest women and their babies. Records for 4456 rural women from the 2015–16 Tanzania Demographic and Health Survey with a live birth in the preceding 5 years were examined. Proportions of births at each location (home/PHC/hospital) were calculated by demographic and obstetric characteristics. Multinomial logistic regression was used to obtain crude and adjusted odds ratios of home/PHC and hospital/PHC births based on household wealth, including interaction between wealth and parity. Post-estimation margins analysis was applied to estimate childbirth location by wealth and parity. Hospital-based childbirth uptake was inequitable. The gap between poorest and richest was less pronounced at first birth. Hospital-based care utilization was lowest (around 10%) among the poorest multiparous women, with no increase at high parity (≥ 5) despite higher risk. PHC-based childbirth care was used by a consistent proportion of women after the first birth (range 30–51%). The poorest women utilized it at intermediate parity, but at parity ≥ 5 mostly gave birth at home. In an effort to provide effective childbirth care to all women, context-specific strategies are required to improve hospital-based care use, and poor, rural, high parity women are a particularly vulnerable group that requires specific attention. Improving childbirth care in PHC and strengthening referral linkages would benefit a considerable proportion of women.

Keywords: Obstetrics, maternal and child health, maternal services, equity, primary health care, health inequalities, poverty, rural, hospital, health facilities, health care utilization

Introduction

Mortality around the time of childbirth is essentially a disease of poverty. An inverse relationship between poverty and maternal health has been known for over a century. Wide inequities in maternal and perinatal mortality exist between nations, with low- and middle-income countries being the most affected (Graham *et al.*, 2016). Sub-Saharan Africa (SSA), with only 14% of global population, accounted for 66% (201 000) of maternal deaths, 40% (1 027 000) of newborn deaths and 31% (1 060 000) of stillbirths

in 2015 (Alkema *et al.*, 2016; Blencowe *et al.*, 2016; World Bank, 2020; WHO, 2015a). Wide gradients also exist within countries, with the poorest disproportionately affected (Ronsmans and Graham, 2006; Houweling *et al.*, 2007; Filippi *et al.*, 2016). Such inequities are often masked by national averages (Kinney *et al.*, 2010).

Providing effective childbirth care is challenging where resources are limited, and rural SSA is a particularly arduous setting (Campbell *et al.*, 2016). In SSA countries, primary health care (PHC) has been a central strategy to ensure access

Key messages

- The proportion of facility births is increasing in rural Tanzania, but obstetric care at different levels of the health system varies. We hypothesized a differential use of hospitals for childbirth and assessed interaction between wealth and parity. Inequity in hospital-based childbirth care use may contribute to the burden of greater mortality among the poorest women and their babies.
- Hospital-based childbirth care use was inequitable among women in rural Tanzania. The gap between the poorest and richest use was less pronounced among women at first parity. Uptake of hospital-based care was lowest (around 10%) among poorest multiparous women, remaining low at high parity (≥ 5) despite higher risk of complications and death.
- Rural women's use of PHC childbirth care after the first birth was noteworthy, ranging between 30 and 51% depending on wealth and parity.
- As part of efforts to reach all women with appropriate, timely care, strategies are required to improve uptake of hospital-based care particularly among poor, rural, high parity ones. Improving quality of childbirth care in PHC and referral linkages would benefit a substantial proportion of women.

to services, including intrapartum care, for rural populations. Tanzania, with a population of 59 million (2020) (World Bank, 2020), has been at the forefront of PHC development after independence with its founding principles set out in the Arusha Declaration in 1967 (Bustreo *et al.*, 2019; Dominicus and Akamatsu, 1989). A vision of high-quality PHC for all is expressed in current policy (Vision 2025) (Tanzania URO, 2000). PHC facilities are dispensaries at village level and health centres at ward level (Tanzania MoHSW, 2015). Women can give birth at all levels of the health system, and childbirth in PHC facilities is encouraged for women with no known risk factors at onset of labour (Hanson *et al.*, 2015). In spite of a capillary PHC network, with 85% of the population living within 5 km from a facility, Tanzania's maternal mortality ratio in 2017 remained high at 524 per 100 000 live births (a reduction of 39% from 2000) (WHO, 2015b; 2019) and was among 10 countries worldwide with the highest absolute numbers of maternal, newborn deaths and stillbirths (Lawn *et al.*, 2016).

Obstetric care at different levels of the health system varies markedly in SSA (Campbell *et al.*, 2016). There is growing evidence that outcomes for mothers and their babies improve when women give birth in units offering high-quality care, not just in any facility (Gabrysch *et al.*, 2019; Hanson *et al.*, 2015; Lohela *et al.*, 2019). In Tanzania, advanced management of childbirth complications (including surgery and blood transfusions, equivalent to comprehensive emergency obstetric care [EmOC]) is available in hospitals, while lower-level, PHC facilities are generally able to provide routine childbirth care only (Campbell *et al.*, 2016; Kruk *et al.*, 2016). Although dispensaries and health centres differ in size, number of beds and staffing levels, both types of facilities have similar obstetric capacity and often do not reach a description of basic EmOC facility (Campbell *et al.*, 2016; Hanson *et al.*, 2013). Challenges to the provision of high-quality obstetric care in

lower-level facilities in Tanzania have been amply described and include insufficient staffing, poor infrastructure and low birth volumes (Hanson *et al.*, 2013; Benova *et al.*, 2014; Kruk *et al.*, 2014; Hanson *et al.*, 2015; Straneo *et al.*, 2014; Baker *et al.*, 2015). Although national efforts are underway to increase obstetric care in health centres up to comprehensive EmOC, at the time of analysis very few had been upgraded. Higher mortality among poorer women and their babies may be compounded by their reduced hospital-based childbirth care uptake, where higher-quality obstetric care is more commonly found.

Over the past decade, a shift from home to facility births has been described in SSA, with increasing proportions of women giving birth in facilities in rural and urban contexts and across wealth groups, and Tanzania is no exception (Montagu *et al.*, 2017; Doctor *et al.*, 2019). However, strong socio-demographic differentials continue to be reported (Kyei-Nimakoh *et al.*, 2017; Moyer and Mustafa, 2013; Gabrysch and Campbell, 2009; Campbell *et al.*, 2016; Virgo *et al.*, 2017; Dunlop *et al.*, 2018). There is limited information on how socio-economic groups uptake obstetric care at different levels of the health system. Within the background of a renewed discussion of the most efficient configuration of childbirth care from an equity, quality and cost-efficiency perspective (Kruk *et al.*, 2016; Hanson and Schellenberg, 2019; Kruk *et al.*, 2018; Roder-DeWan, 2020), we aimed to estimate the levels of use of hospital-based childbirth care in rural Tanzania and its association with women's socio-economic status. Given the association of poverty and high parity, and the latter's implications for obstetric care, we investigated whether the association between wealth and hospital-based childbirth depended on parity.

Methods**Study setting**

In Tanzania, the most recent (2015–16) Demographic and Health Survey (DHS) estimated that 63% of births in the 5-year period preceding the survey took place in health facilities (54% in rural and 86% in urban areas) (Tanzania MoHCDGEC, 2016). In the same period, there were 6790 facilities (including public, faith-based, parastatal and private), from which 257 (3.8%) were hospitals (Tanzania MoHSW, MoHMZ, National Bureau of Statistics [NBS], Office of the Chief Government Statistician [OCGS], and ICF, 2015). There were an estimated 12 million women of reproductive age and approximately 1.9 million births in 2015.

Data and population

Data from the 2015–16 Tanzania DHS were used. DHS are cross-sectional, nationally representative surveys of households, with women of reproductive age (15–49 years) self-reporting on the use of reproductive and maternal healthcare. Approximately 12 500 households were visited, and 13 000 women interviewed. Records of women living in rural areas in mainland Tanzania were used in this analysis, if they reported a live birth in the 5 years preceding the survey. Classification as 'rural' in DHS was based on census enumeration units (Tanzania NBS and Zanzibar OCGS, 2013).

Definitions

The outcome variable was the location of the most recent live birth, in three categories: home (respondent's home, other home and en route to provider), PHC facility (dispensary, health centre, maternity home and 'other facility') and hospital (district, regional, referral or tertiary/university). All public and private (non-profit/profit) PHC facilities and hospitals were included.

Socio-economic status (SES): In DHS, SES is based on availability of durable household assets (Vyas and Kumaranayake, 2006). A wealth score is generated for each sampled household using principal component analysis and the households are then subdivided into equal-size wealth quintiles. Distribution of wealth is uneven across different contexts, with the highest (wealthiest) SES quintile households under-represented in rural contexts. In rural Tanzania (DHS 2015–16), there were only 14.6% and 2.5% women in Quintile 4 (richer) and Quintile 5 (richest), respectively. Thus, for the purpose of this analysis, the two highest wealth quintiles were merged, resulting in the creation of four wealth groups (poorest, poorer, middle and wealthiest). The terms richest/wealthiest refer to relative wealth in a poor, rural context, thus indicate women from the least poor households. To analyse the interaction of SES and parity, a binary SES variable was created, by generating two equal groups based on wealth scores (wealth score < median recoded as poorer, wealth score \geq median coded as richer 50%).

Parity group refers to a woman's parity at index pregnancy (0, 1–2, 3–4 and \geq 5). Grand multiparity was defined as parity \geq 5 (Mgaya *et al.*, 2013).

Maternal age at index birth was coded in 5-year age groups, grouping categories at the extremes of age because they had fewer than 100 observations (\leq 19, 20–24, 25–29, 30–34, 35–39 and \geq 40 years). The 20–24 years' group was used as reference.

Maternal education was recoded into three categories: no education, completed primary and completed secondary or higher.

Marital status at survey was recoded into currently married/cohabiting and not currently married/cohabiting.

Zone of residence: Tanzania is divided into 21 administrative regions, grouped into eight zones (Tanzania MoHCDGEC, MoH [Zanzibar], and ICF, 2016). We used the eight zones to account for sub-national variation in outcomes and service availability (Armstrong *et al.*, 2016). All eight zones include rural areas; the Eastern zone includes the Dar es Salaam urban conglomerate.

Antenatal care (ANC) for the index pregnancy was categorized into no visits, 1–3 visits and \geq 4 visits.

Other obstetric characteristics studied were the following: multiple index birth, a previous birth in the recall period by Caesarean section (CS), death of a previous child (born in the recall period) aged 1–12 months, death of a previous child (born in the recall period) aged <1 month, a short preceding birth interval (\leq 12 months).

Statistical analysis

Analysis was performed using STATA IC 15 software. Complex survey design and non-response (stratification, clustering and survey weights) were accounted for using svyset commands. Characteristics of the sample were analysed with proportions and 95% confidence intervals (CIs) of outcome and

exposure variables. There were no missing data for the variables examined. Proportions of subgroups of women at each level of outcome (hospital/PHC/home birth) for each exposure variable were examined using bivariate analysis. As the interaction between SES and parity was of interest, the proportion of women giving birth at each location by combinations of parity levels and a binary SES variable (poorer/richer) were determined. Associations between the outcome variable and dependent variables (demographic, geographical characteristics, SES, ANC care received and available obstetric factors) were assessed in bivariate analysis. Variables which were significant at $P < 0.05$ level in bivariate analysis were included in the final multivariable model. Multinomial logistic regression was used as the outcome variable had three categories, thus allowing to include all births in one model. The baseline outcome was birth in PHC, thus the model produced odds ratios (ORs) of home vs PHC and hospital vs PHC birth. In the final multivariable model, we tested for an interaction between SES and parity group. We calculated the margins to obtain predicted percentages of women giving birth at each of the three locations, depending on their SES and parity group combination. Results were used to calculate the difference or gap in hospital or PHC uptake for birth between the wealthiest and poorest women.

Ethical approval

The DHS receive government permission, use informed consent and assure respondents of confidentiality. Permission to use the dataset for the purpose of this analysis was obtained from the DHS programme.

Results

Population characteristics

Observations of 4456 women living in rural mainland Tanzania and the circumstances of their most recent live birth in the 5 years preceding the survey were included in the analysis. Home birth was reported by 41% of women and a slight majority reported a facility birth (59%): 35% in PHCs and 24% in hospitals. Women from the wealthiest households were under-represented, with only 17% in the highest group compared with 28% in the lowest. Approximately one in five women was nulliparous at index birth (22%), while 25% had parity five or higher. Background characteristics are summarized in Supplementary Table S1.

Results of bivariate analysis are reported in Table 1. The percentage of rural women using hospitals for childbirth increased with higher SES, from 16% in the poorest group to 45% in the wealthiest. PHC births also rose with increasing wealth, although less steeply than hospital births. As parity increased, hospital births reduced sharply, while PHC births had a less clear trend across SES and parity and varied only marginally at around one-third of births. Hospital births increased with higher maternal age, maternal education and number of ANC visits. There was a wide variation in hospital births across the eight zones, ranging from 16% in the Lake Zone to 39% in the Southern Highlands. PHC facilities provided a substantial proportion of childbirth care in rural areas of all zones (range 23–48%, median 38%).

Examining SES and parity together, hospital births were more frequent in women from wealthier households in all parity groups, with percentages reducing as parity increased

Table 1. Percentages of home, primary care facility or hospital birth by subgroups among rural women with a recent live birth (TDHS 2015–16) *n* = 4456

Variable	Women in category (%)	Home births		PHC facility births		Hospital births	
		Number	% (95% CI)	Number	% (95% CI)	Number	% (95% CI)
SES							
Poorest	1351 (29.3)	742	55.5 (50.5–60.3)	413	29.1 (25.4–33.0)	196	15.5 (12.3–19.1)
Poorer	1239 (28.4)	559	45.7 (41.5–50.0)	467	36.5 (32.7–40.5)	213	17.8 (14.9–21.1)
Medium	1104 (25.2)	383	35.5 (31.1–40.1)	448	39.7 (35.5–44.0)	273	24.9 (21.5–28.6)
Wealthiest	762 (17.1)	131	18.1 (14.1–23.0)	288	36.9 (31.9–42.2)	343	45.0 (39.4–50.7)
Parity							
0	963 (22.2)	245	25.6 (22.2–29.4)	333	32.9 (29.2–36.8)	385	41.5 (37.1–46.1)
1–2	1342 (30.6)	537	40.6 (36.4–45.1)	514	37.3 (33.5–41.3)	291	22.1 (19.0–25.5)
3–4	1057 (23.6)	475	45.4 (41.1–50.0)	410	38.9 (35.0–42.9)	172	15.7 (12.9–18.9)
≥5	1094 (23.6)	558	52.7 (48.2–57.2)	359	30.9 (27.5–34.6)	177	16.4 (13.6–19.6)
Maternal age at index birth (years)							
≤19	743 (17.6)	241	32.0 (27.7–36.6)	280	36.2 (32.0–40.7)	222	31.8 (27.2–36.8)
20–24	1124 (25.0)	459	41.6 (37.7–45.7)	412	35.7 (32.1–39.4)	253	22.7 (19.6–26.1)
25–29	946 (21.2)	400	42.9 (38.2–47.8)	343	35.7 (31.6–40.0)	203	21.4 (18.0–25.2)
30–34	744 (16.3)	321	43.9 (38.7–49.3)	255	33.4 (28.9–38.1)	168	22.7 (18.9–27.1)
35–39	594 (13.1)	249	43.7 (38.2–49.4)	228	35.7 (30.8–41.0)	117	20.6 (16.4–25.4)
40–49	305 (6.8)	145	47.9 (41.4–54.5)	98	32.3 (26.8–38.4)	62	19.7 (15.2–25.3)
Maternal education at survey							
No education	1054 (23.8)	577	54.7 (49.8–59.5)	343	32.5 (28.7–36.7)	134	12.8 (10.2–15.9)
Completed primary	3028 (67.5)	1191	40.2 (37.0–43.5)	1120	35.6 (32.9–38.5)	717	24.2 (21.6–27.0)
Secondary and above	374 (8.7)	47	13.1 (9.6–17.7)	153	38.9 (33.1–45.0)	174	48.0 (41.7–54.4)
Marital status at survey							
Currently married or cohabiting	3695 (82.9)	1548	43.6 (39.1–46.1)	1334	35.0 (32.4–37.8)	813	22.4 (20.0–25.1)
Not currently married or cohabiting	761 (17.1)	267	35.1 (30.5–40.0)	282	35.9 (31.6–40.5)	212	29.0 (24.8–33.6)
Zone of residence							
Western	515 (14.0)	255	50.1 (39.4–60.9)	170	31.8 (24.3–40.4)	90	18.1 (11.8–26.6)
Northern	420 (10.1)	164	40.7 (28.7–54.0)	95	22.6 (16.8–29.8)	161	36.6 (26.1–48.6)
Central	569 (14.0)	246	41.6 (32.6–51.2)	165	31.1 (24.4–38.8)	158	27.3 (20.9–34.8)
Southern Highlands	394 (6.0)	48	14.3 (7.2–26.6)	189	46.3 (34.9–58.0)	157	39.4 (30.1–49.5)
Southern	273 (5.5)	53	19.5 (12.5–29.0)	128	47.6 (39.5–55.8)	92	33.0 (25.3–41.7)
Southern West Highlands	588 (11.0)	256	37.5 (27.9–48.1)	255	43.5 (34.1–53.3)	77	19.1 (12.9–27.3)
Lake	1440 (32.3)	732	51.2 (46.4–55.9)	487	32.6 (28.8–36.5)	221	16.3 (13.0–20.3)
Eastern	257 (7.1)	61	24.6 (15.7–36.5)	127	47.8 (35.9–60.0)	69	27.6 (19.1–38.2)

(continued)

Table 1. (Continued)

Variable	Women in category (%)		Home births		PHC facility births		Hospital births	
	Number	% (95% CI)	Number	% (95% CI)	Number	% (95% CI)	Number	% (95% CI)
Antenatal visits								
None	78	77.7 (68.3–84.9)	14	13.9 (7.8–23.5)	7	8.5 (3.9–17.5)		
1–3	2344 (52.4)	46.9 (42.9–51.0)	783	32.4 (29.2–35.8)	473	20.7 (18.1–23.5)		
≥4	2013 (45.4)	33.0 (29.7–36.4)	819	39.4 (36.5–42.4)	545	27.6 (24.7–30.8)		
Index pregnancy was multiple								
No	4371 (98.0)	41.3 (38.1–44.6)	1591	35.4 (32.8–38.0)	997	23.4 (20.9–26.0)		
Yes	85 (2.0)	40.6 (29.4–53.0)	25	26.4 (17.6–37.5)	28	33.0 (22.8–45.1)		
Previous birth was by CS								
No or no previous birth	4428 (99.1)	41.5 (38.2–44.8)	1612	35.3 (32.7–37.9)	1005	23.3 (20.9–25.9)		
Yes	28 (0.6)	13.9 (5.1–32.9)	4	17.0 (6.1–39.5)	20	69.1 (47.9–84.4)		
Short previous birth interval (≤12 months)								
No or no previous birth	4410 (99.1)	41.2 (38.0–44.5)	1601	35.2 (32.6–37.8)	1018	23.6 (21.2–26.2)		
Yes	46 (0.9)	51.6 (36.0–66.9)	15	33.1 (19.5–50.3)	7	15.3 (7.2–29.7)		
Death of newborn preceding index birth								
No or no previous birth	4377 (98.0)	41.3 (38.1–44.6)	1586	35.1 (32.6–37.8)	1008	23.6 (21.2–26.2)		
Yes	79 (2.0)	38.5 (27.4–51.0)	30	38.1 (26.7–51.0)	17	23.4 (14.2–36.0)		
Death of child born before index birth aged >1 month and <12 months								
No	4372 (98.0)	41.0 (37.8–44.4)	1589	35.3 (32.7–37.9)	1012	23.7 (21.3–26.3)		
Yes	84 (2.0)	53.3 (41.1–65.1)	27	31.1 (21.2–43.1)	13	15.6 (8.6–26.7)		
Parity by SES (<i>n</i> = 4456)								
Parity 0 (<i>n</i> = 963)								
Poorer 50%	412 (43.5)	34.2 (28.4–40.6)	142	32.6 (27.3–38.4)	125	33.2 (26.5–40.7)		
Richer 50%	551 (56.5)	19.0 (15.1–23.6)	191	33.1 (28.2–38.4)	260	47.9 (42.5–53.4)		
Parity 1–2 (<i>n</i> = 1342)								
Poorer 50%	647 (46.5)	53.7 (48.4–59.0)	238	35.5 (30.8–40.6)	72	10.7 (8.1–14.1)		
Richer 50%	695 (53.5)	29.2 (24.3–34.7)	276	38.8 (33.7–44.2)	219	32.0 (27.5–36.8)		
Parity 3–4 (<i>n</i> = 1057)								
Poorer 50%	533 (49.5)	58.7 (53.2–64.0)	170	31.4 (27.0–36.4)	60	10.0 (7.4–13.3)		
Richer 50%	524 (50.5)	32.5 (27.3–38.1)	240	46.3 (40.7–52.0)	112	21.3 (17.1–26.1)		
Parity ≥5 (<i>n</i> = 1094)								
Poorer 50%	636 (57.8)	59.1 (53.6–64.4)	187	27.4 (23.1–32.0)	83	13.6 (10.5–17.4)		
Richer 50%	458 (42.2)	44.0 (38.4–49.9)	172	35.8 (30.9–41.0)	94	20.2 (16.3–24.8)		

across all wealth groups. The drop of hospital use for childbirth was seen among poorer women already at Parity 1–2, while among wealthier women, this reduction was seen at Parity 3–4. Despite the decrease, the percentage of births in hospitals remained higher among wealthier than poorer women in all parity groups. The gap between the poorest and wealthiest women in hospital births was greatest at Parity 1–2 (Supplementary Graph 1).

Logistic regression

Results of bivariate and multivariate logistic regression are reported in Table 2.

In adjusted analysis, compared with women from the poorest households' group, all wealthier women were less likely to have given birth at home vs in PHC. The wealthiest were 66% less likely to do so. High-parity women (≥ 5) had higher odds of home birth (OR 1.54, 95% CI 1.05–2.25) compared with the reference group Parity 1–2, while odds in other parity groups were not significantly different from baseline. Higher odds of a home birth were seen in women with no ANC or 1–3 ANC visits compared with women with ≥ 4 visits. Compared with women with primary education, those with no education had higher odds of a home birth, while those with secondary or higher education had reduced odds. Women residing in four zones (Southern Highlands; Southern and Southern West Highlands; and Eastern) had reduced odds compared with those residing in the reference Lake Zone.

In adjusted analysis, the wealthiest rural women had higher odds (OR 1.78, 95% CI 1.26–2.50) of a hospital vs a PHC birth compared with the poorest, while other wealth groups were not significantly different from the poorest. Higher odds of hospital vs PHC birth were found in Parity 0 women compared with baseline Parity 1–2 (OR 3.22, 95% CI, 2.34–4.43), while the odds were reduced in higher-parity groups. The effect of maternal age was confounded in crude analysis; in adjusted analysis, the odds of hospital vs PHC birth increased with age. Women with a previous birth by CS had higher odds of a hospital birth compared with those with no previous CS, while women with no education, compared with those with primary education, had reduced odds of hospital vs PHC birth. Higher odds were observed in women residing in two zones (Northern and Central) compared with those residing in the reference Lake Zone.

Interaction between SES and parity

To assess the joint effects of parity and SES, the final adjusted multinomial logistic regression model was run with an interaction term between the two variables. A likelihood ratio test comparing the model with and without interaction indicated better fit of the model with interaction ($P = 0.006$). The reference group included the poorest women at parity ≥ 5 , as this group had the lowest use of hospital-based childbirth care and was the most numerous wealth/parity subgroup ($n = 406$). Results are shown in Supplementary Tables S2a and S2b.

All combinations of SES and parity had lower odds than the baseline category of a home vs a PHC birth, although not all reached statistical significance at $P < 0.05$. The richest women at high parity (≥ 5) had the lowest adjusted OR (0.29, 95% CI 0.15–0.57) compared with the reference group. The poorest women at Parity 0 had an OR of 8.03 (95% CI, 4.45–14.46) compared with the baseline of a hospital vs PHC birth, while

at other parity levels the ORs were not significantly different. Women at Parity 1–2 from poorer, medium and richest groups had higher odds of hospital vs PHC childbirth compared with the baseline group; the OR was non-significant in the poorest group. In other groups, ORs were not significantly different from the baseline.

We predicted the percentages of childbirth for each combination of SES and parity in each location using margins analysis; results are reported in Table 3 and displayed in Graph 1(A–C). Across all SES groups, hospital-based childbirth (Graph 1A) was highest at first birth, at $>40\%$. Use of hospitals reduced in all SES groups with increasing parity, but the shape of this decline varied. Among the wealthiest women, hospital use decreased gradually, reaching its lowest (around 25%) at parity 3–4. Among the poorest, the decline was abrupt after parity 0, levelling out at 12% at parity 1–2. The effect of wealth on PHC births was more complex (Graph 1B). The predicted percentages at this level were lowest among nulliparous women in all wealth groups. Among the wealthiest, the percentage rose with parity, reaching its maximum (51%) at parity 3–4. Among the poorest women, the predicted utilization reached the highest level at parity 1–2 (39%) and then levelled off at around 30%. In all wealth groups, after parity 0, $\geq 30\%$ of women were predicted to give birth in a PHC. Median utilization of PHC facilities in parous women was 35% (range 30–51%), while in women at first parity it was 27% (range 20–31%). The percentage of births at home (Graph 1C) increased as parity rose in all wealth groups and was lowest among the wealthiest women in all parity groups.

The profiles of birth location among the two extremes of wealth (poorest and richest women) are compared in Graph 2. Among the richest women, there was a shift in the location of births from mainly hospital (at parity 0) to mainly PHC facilities (at parity ≥ 5). In the poorest women's group, between parity groups 0 and ≥ 5 , decline in hospital births was accompanied by a sharp rise in home births, with a small increase in PHC births.

Discussion

This study explored rural women's differential use of childbirth care in Tanzania. We report three key findings. First, there was a socio-economic inequity in rural women's use of hospital-based childbirth, which additionally varied with parity. Second, the poorest multiparous women had the lowest use of hospital-based care for childbirth (around 10%), with no increase in uptake at grand multiparity despite increased risk. This group also had lower uptake of PHC care. Third, PHC facilities provided care to a sizeable proportion of women after the first birth, with a median uptake of 35% (range 30–51%) by women after the first birth (compared with a median of 27% by women at first birth).

The poorest women in rural Tanzania were less likely than those from the wealthiest households to give birth in hospitals, where advanced management of childbirth complications was available. The study adds to existing evidence that wealth is not just a determinant for facility birth, but also for uptake of hospital-based childbirth care within the health system. It expands findings of a previous sub-national study (Straneo *et al.*, 2014) and earlier studies (Benova *et al.*, 2014). We found that the gap between the poorest and wealthiest women in use of hospitals was less pronounced among nulliparous

Table 2. Crude and adjusted ORs by multinomial logistic regression of home vs PHC birth (left) and hospital vs primary care births (right) in rural women, with a live birth in the last 5 years (Tanzania, DHS 2015–16)

Variable	Home birth vs PHC birth			Hospital birth vs PHC birth		
	Crude OR	P-value	Adjusted OR ^a	Crude OR	P-value	Adjusted OR ^a
SES						
Poorest	ref		ref	ref		ref
Poorer	0.66 (0.51–0.84)	1	0.75 (0.58–0.96)	0.92 (0.67–1.26)	0.6	0.90 (0.66–1.23)
Medium	0.47 (0.36–0.61)	<0.001	0.56 (0.43–0.74)	1.18 (0.86–1.62)	0.3	1.05 (0.78–1.41)
Wealthiest	0.26 (0.18–0.37)	<0.001	0.34 (0.23–0.50)	2.30 (1.61–3.27)	<0.001	1.78 (1.26–2.50)
Parity						
0	0.71 (0.56–0.91)	0.006	0.84 (0.63–1.12)	2.13 (1.67–2.73)	<0.001	3.22 (2.34–4.43)
1–2	ref		ref	ref		ref
3–4	1.07 (0.87–1.31)	0.5	1.07 (0.82–1.39)	0.68 (0.51–0.90)	0.007	0.46 (0.32–0.65)
≥5	1.57 (1.26–1.95)	<0.001	1.54 (1.05–2.25)	0.89 (0.67–1.19)	0.4	0.59 (0.39–0.89)
Maternal age at birth						
≤19	0.76 (0.60–0.96)	0.02	0.80 (0.59–1.08)	1.38 (1.05–1.80)	0.019	0.86 (0.62–1.19)
20–24	ref		ref	ref		ref
25–29	1.03 (0.82–1.29)	0.796	0.86 (0.66–1.13)	0.94 (0.72–1.24)	0.664	1.89 (1.37–2.60)
30–34	1.13 (0.89–1.43)	0.322	0.80 (0.56–1.13)	1.07 (0.81–1.42)	0.636	2.69 (1.78–4.08)
35–39	1.05 (0.81–1.35)	0.724	0.61 (0.40–0.93)	0.90 (0.64–1.27)	0.551	2.49 (1.51–4.13)
40–49	1.27 (0.91–1.76)	0.156	0.69 (0.44–1.08)	0.96 (0.65–1.41)	0.829	2.64 (1.58–4.42)
Maternal education at survey						
No education	1.49 (1.23–1.80)	<0.001	1.19 (0.97–1.45)	0.58 (0.44–0.75)	<0.001	0.70 (0.53–0.93)
Primary	ref		ref	ref		ref
Secondary and above	0.30 (0.21–0.43)	<0.001	0.38 (0.25–0.59)	1.82 (1.39–2.39)	<0.001	1.03 (0.76–1.41)
Marital status at survey						
Currently married or cohabiting	ref		ref	ref		ref
Not currently married or cohabiting	0.80 (0.64–1.01)	0.059	0.80 (0.64–1.01)	1.26 (1.00–1.59)	0.049	1.03 (0.76–1.41)

(continued)


Table 2. (Continued)

Variable	Home birth vs PHC birth			Hospital birth vs PHC birth		
	Crude OR	P-value	Adjusted OR ^a	Crude OR	P-value	Adjusted OR ^a
Zone of residence						
Western	1.00 (0.63–1.60)	0.989	0.82 (0.51–1.31)	1.13 (0.66–1.93)	0.645	1.28 (0.73–2.23)
Northern	1.15 (0.67–1.96)	0.621	1.40 (0.86–2.28)	3.23 (1.95–5.34)	<0.001	2.54 (1.56–4.13)
Central	0.85 (0.54–1.34)	0.483	0.89 (0.57–1.39)	1.75 (1.12–2.74)	0.014	1.83 (1.16–2.88)
Southern Highlands	0.20 (0.09–0.46)	<0.001	0.24 (0.11–0.57)	1.70 (1.01–2.86)	0.045	1.52 (0.90–2.29)
Southern	0.26 (0.15–0.45)	<0.001	0.27 (0.16–0.47)	1.39 (0.89–2.16)	0.151	1.44 (0.90–2.29)
Southern West Highlands	0.55 (0.34–0.90)	0.016	0.55 (0.34–0.91)	0.88 (0.51–1.52)	0.637	0.85 (0.49–1.49)
Lake	ref		ref	ref		ref
Eastern	0.33 (0.18–0.61)	0.001	0.43 (0.22–0.83)	1.15 (0.65–2.06)	0.63	1.17 (0.67–2.07)
ANC (visits)						
None	6.71 (3.53–12.74)	<0.001	5.75 (3.23–10.23)	0.87 (0.29–2.58)	0.801	0.87 (0.25–3.07)
1–3	1.73 (1.45–2.07)	<0.001	1.53 (1.28–1.83)	0.91 (0.76–1.08)	0.276	1.10 (0.92–1.32)
≥4	ref		ref	ref		ref
Multiple live birth at index pregnancy						
No	ref		ref	ref		ref
Yes	1.32 (0.75–2.31)	0.332	1.17 (0.68–2.00)	1.89 (1.02–3.50)	0.042	2.24 (1.18–4.26)
Previous birth was by CS						
No	ref		ref	ref		ref
Yes	0.69 (0.16–3.03)	0.628	0.71 (0.18–2.84)	6.15 (1.89–20.0)	0.003	6.96 (2.15–22.57)
Short preceding birth interval (≤12 months)						
No	ref		ref	ref		ref
Yes	1.33 (0.63–2.79)	0.45		0.69 (0.26–1.84)	0.457	
Previous neonatal death						
No	ref		ref	ref		ref
Yes	0.86 (0.49–1.51)	0.596		0.92 (0.46–1.83)	0.803	
Previous baby died						
No	ref		ref	ref		ref
Yes	1.47 (0.85–2.56)	0.169		0.75 (0.35–1.58)	0.441	

^aAdjusted for wealth, parity, maternal age at index birth, maternal education, marital status, ANC visits and multiple index pregnancy.

Table 3. Predicted margins (%) for each outcome (home/PHC/hospital birth) in rural women, Tanzania 2015–16 DHS, by SES and parity, adjusted for wealth, parity, maternal age at index birth, maternal education, marital status, ANC visits and multiple index pregnancy


Predicted margins (%) of rural women by outcome, SES group and parity					
Outcome	SES group	Parity 0	Parity 1–2	Parity 3–4	Parity ≥ 5
Home birth	Poorest	28 (21–35)	51 (44–58)	59 (52–65)	58 (51–65)
	Poorer	28 (21–35)	42 (35–49)	47 (40–54)	56 (48–63)
	Medium	21 (14–28)	33 (27–39)	39 (33–46)	53 (45–61)
	Wealthiest	16 (10–22)	21 (11–30)	25 (17–33)	30 (19–41)
PHC birth	Poorest	20 (14–25)	39 (33–46)	31 (25–37)	30 (24–36)
	Poorer	31 (22–39)	34 (28–40)	44 (38–51)	32 (26–39)
	Medium	28 (22–35)	42 (35–49)	46 (39–52)	35 (27–42)
	Wealthiest	25 (18–32)	35 (26–43)	51 (41–61)	47 (35–59)
Hospital birth	Poorest	52 (43–61)	10 (6–14)	10 (6–14)	12 (8–16)
	Poorer	42 (32–51)	23 (17–30)	8 (6–11)	12 (8–16)
	Medium	51 (43–59)	25 (19–30)	15 (10–20)	12 (8–16)
	Wealthiest	59 (51–66)	44 (37–52)	25 (17–32)	23 (15–31)

**Graph 1.** Predicted margins of birth at each location, rural Tanzania 2015-16, 95% CI

women. Health policy in Tanzania (Tanzania MoHSW, 2008) recommends that women's first births should take place in hospitals. In spite of the existing recommendations, uptake by rural women at first birth was not universal, as just over half used hospital-based care. We found that hospital-based care use among nulliparous women was very similar across wealth groups. Factors other than wealth are likely to limit hospital use at first birth; amongst these distance to hospital from a woman's residence, which could not be accounted for, stands out, and interaction between distance and wealth has been described (Bai *et al.*, 2002; Hanson *et al.*, 2015; Wong *et al.*, 2020). Our finding is in line with that of other researchers indicating that women and their families recognize the first birth as a higher-risk one (Jahn *et al.*, 1998; Dunlop *et al.*, 2018). Utilization of hospitals decreased at different rates in SES groups with increasing parity and the gap between the poorest and wealthiest was widest at Parity Level 1–2 but persisted across all higher-parity groups. A switch in birth location away from facilities between first- and

second-order births was found to be less likely in wealthier households across low- and middle-income countries (Benova *et al.*, 2017). What this study adds is that among the poorest women between first-order and successive births (Parity 1–2), there was a switch within the health system, from hospital-based care to PHC-based care, and to home-based care.

Utilization of hospital-based childbirth is lowest among the poorest, multiparous women. Use, as estimated by margins analysis, dropped to around 10% at all levels of parity after the first-order birth. Despite greater risk in women with ≥ 5 previous births of adverse pregnancy outcomes, including haemorrhagic complications (Bai *et al.*, 2002; Mgaya *et al.*, 2013; Filippi *et al.*, 2016), there was no increased hospital care uptake among the poorest women. Factors contributing to this may be inadequate counselling during ANC on hospital-based childbirth resulting in low perceived risk (Pembe *et al.*, 2008), childcare duties at home, and greater economic constraints due to larger families. Reducing facility



Graph 2. Predicted margins (percentages) of Hospital/PHC/Home births of poorest and wealthiest rural women by parity, Tanzania 2015-16

use with increasing parity is well documented (Kyei-Nimakoh *et al.*, 2017; Moyer and Mustafa, 2013; Gabrysch and Campbell, 2009; Berhan and Berhan, 2014). Results of our study add that, in rural Tanzania, multiparous women will opt for a home birth when economic means are limited but will uptake PHC-based childbirth when resources are available. Poor multiparous women constitute a disadvantaged group, least served by hospitals or indeed by any facility. Qualitative studies in southern Tanzania (Kowalewski *et al.*, 2000) indicated that in the community, these women were perceived as vulnerable due 'to fatigue and (being) overburdened with household duties', and precisely these factors prevented them from accessing health services.

Although our analysis focused on hospital births, rural women's uptake of childbirth care in lower-level facilities is noteworthy. PHC units (health centres and dispensaries) have a critical role in childbirth care in rural Tanzania, as 61% of all facility births took place here, and utilization ranged from 30 to 51% (median 35%) across all SES groups after Parity 0. The greatest use is among the wealthiest, at intermediate and high parity. This has relevance in the current debate on reorganization of maternity care in low-income countries (Campbell *et al.*, 2016; Kruk *et al.*, 2018; Hanson and Schellenberg, 2019). From these data, childbirth care in PHC facilities is used by all wealth groups: among the wealthiest, uptake increases with parity, while among the poorest, utilization is mostly at intermediate levels of parity. Comparing the shift in births between the two extremes of parity in the poorest and richest women suggests that, in this context, a reduction in the availability of facilities providing childbirth care without other measures may result in an increase in the already-high level of home births among the poorest women.

Policy recommendations arising from this study include three main points. Firstly, aiming attention on the poorest women allows identification of health system adjustments to mitigate the effects of poverty on childbirth-related deaths (WHO, 2008). A subsidized voucher scheme has been applied in Kenya (Dennis *et al.*, 2019; Wong *et al.*, 2020). Maternity waiting homes may contribute to facilitating access to

hospitals (Virgo *et al.*, 2017), and there is evidence that they are utilized preferentially by poorer women (Fogliati *et al.*, 2017). Secondly, high-parity women's low use of hospital-based childbirth care, particularly among the poorest, requires urgent action. All women should receive appropriate, timely care. National policy should focus attention on grand-multiparous women as a particularly higher-risk group. Guidelines should be in place to prepare these women for hospital-based births. They may include adapted birth preparedness plans and emergency transport during labour to improve geographic accessibility. Thirdly, the current debate on centralization of childbirth care must take into account the sizeable proportions of women using PHC facilities for obstetric care. Care at childbirth is part of essential care, as defined in the Alma Ata declaration of PHC (Beard and Redmond, 1979), which also includes the 'scientifically sound' concept. Effective coverage is increasingly advocated, in place of contacts with care (Campbell *et al.*, 2016; Marsh *et al.*, 2020). In this context, to achieve effective coverage for the large proportion of women who uptake PHC-based care, quality-adjusted coverage (Marsh *et al.*, 2020) must be available at the base of the health system pyramid (Hanson and Schellenberg, 2019; Straneo *et al.*, 2014; Fogliati *et al.*, 2015). Comprehensive EmOC in strategically identified rural health centres is one possible solution (Nyamtema *et al.*, 2016); a locally adapted and community-participated reduction of birth sites may be necessary to balance quality and coverage of care (Fogliati *et al.*, 2015). From a policy perspective, the position of childbirth care in PHC should be reappraised.

Limitations

This analysis is the first, to the best of our knowledge, to examine the use of different levels of the health system for childbirth among rural women in Tanzania and analysed the interaction between wealth and parity. It is based on nationally representative data, from a country that has consistently supported the development of a PHC network (Tanzania MoHSW, 2007; Tanzania MoHSW, MoHMZ, NBS, OCGS,

and ICF, 2015), and thus is a model for countries developing rural obstetric care. The Tanzania DHS is unique in allowing identification of facility type (hospital, health centre or dispensary) in both the public and the private sectors, thus providing a more detailed picture of where women report giving birth (Tanzania MoHCDGEC, 2016). The DHS data set was complete, with very little non-response and missing data. Multinomial logistic regression allowed us to include all three locations in one model and thus study factors significant in use of hospital vs PHC facilities, and home vs PHC. Some caution should be applied when interpreting the findings, in terms of the cross-sectional nature of the DHS data and the possible response bias. Since in Swahili all facilities may be referred to as 'hospitali', lower-level public health facilities may be misreported as district hospitals. The DHS interviewers are instructed to circle a type of facility, if known, and if not, to write down the name of the facility, which is later coded as a specific type of facility by the field supervisor. This non-differential misclassification of facilities may bias results and may have led to weaker associations. This DHS collects limited information on obstetric risk factors (Virgo *et al.*, 2017); thus, use of facilities due to risk factors identified during ANC (such as hypertension or maternal infections) cannot be fully captured. Even the risk factors for which some information is available may not fully reflect women's knowledge prior to index birth; e.g., limitations in DHS questionnaire identification of twin pregnancies have been described recently (Hanson *et al.*, 2019). This is unlikely to modify findings, as in a study on four East African countries no association between obstetric risk and birth location was found (Virgo *et al.*, 2017), while wealth and education were strong determinants. Information on referral was lacking; thus, hospital births may include women who had been referred during labour from a PHC unit. In previous studies we found that intra-partum referral rates were very low (Straneo *et al.*, 2016; Fogliati *et al.*, 2017); thus, this too is unlikely to change the findings. Distance travelled to facilities could not be taken into account. Recent studies indicate that travel time is an important factor for hospital births in Tanzania (Wong *et al.*, 2020), and current distribution leaves the rural poor underserved (Wong *et al.*, 2019). Additionally, the study broadly categorized facilities by level (PHC and hospitals) but could not account for variation of quality of care within levels, such as more limited quality at hospital level or more advanced care in health centres (Nyamtema *et al.*, 2016; Tanzania MoHSW, MoHMZ, NBS, OCGS, and ICF, 2015).

In conclusion, the study found that in rural Tanzania the use of hospital-based childbirth was not equitable. Inequity varied with parity level: at first birth, uptake varied only minimally with wealth, while in successive births it was strongly dependent on SES and parity. Uptake was the lowest amongst the poorest, multiparous women, with no increase in uptake at grand multiparity (≥ 5), in spite of increased risk. PHC-based childbirth accounted for a median of 35% of births after the first in this setting.

To leave no one behind in attaining Sustainable Development Goal 3 on maternal and preventable newborn mortality (Boerma *et al.*, 2018), it is necessary to identify who is underserved at childbirth and make adjustments to improve the use of high-quality care, bearing in mind that from a human rights' perspective health care should contribute to equity

(Tudor Hart, 1971). Strategies are needed to improve uptake of hospital-based care among the poorest, rural women, particularly at high parity. A reassessment of the whole district health system, which may involve re-evaluation of childbirth care in PHC and strengthening referral linkages, would benefit a substantial proportion of women.

Supplementary data

Supplementary data are available at *Health Policy and Planning* online.

Data availability statement

The dataset is available to the public at www.measuredhs.com.

Acknowledgements

The authors wish to acknowledge ICF (Rockville, Maryland USA) for providing access to the Tanzanian DHS database.

Conflict of interest statement The authors declare that they have no conflict of interest.

References

- Alkema L, Chou D, Hogan D *et al.* and UN Maternal Mortality Estimation Inter-Agency Group. 2016. Global, regional, and national levels and trends in maternal mortality between 1990 and 2015, with scenario-based projections to 2030: a systematic analysis by the UN Maternal Mortality Estimation Inter-Agency Group. *Lancet* 387: 462–74.
- Armstrong CE, Martínez-Álvarez M, Singh NS *et al.* 2016. Subnational variation for care at birth in Tanzania: is this explained by place, people, money or drugs? *BMC Public Health* 16: 795.
- Bai J, Wong FW, Bauman A, Mohsin M. 2002. Parity and pregnancy outcomes. *American Journal of Obstetrics and Gynecology* 186: 274–8.
- Baker U, Peterson S, Marchant T *et al.* 2015. Identifying implementation bottlenecks for maternal and newborn health interventions in rural districts of the United Republic of Tanzania. *Bulletin of the World Health Organization* 93: 380–9.
- Beard TC, Redmond S. 1979. Declaration of Alma-Ata. *The Lancet* 313: 217–8.
- Benova L, Cumming O, Gordon BA, Magoma M, Campbell OMR. 2014. Where there is no toilet: water and sanitation environments of domestic and facility births in Tanzania. *PLoS One* 9: 1–10.
- Benova L, Macleod D, Radovich E, Lynch CA. 2017. Should I stay or should I go?: consistency and switching of delivery locations among new mothers in 39 Sub-Saharan African and South/Southeast Asian countries. *Health Policy and Planning* 32: 1294–308.
- Berhan Y, Berhan A. 2014. A meta-analysis of socio-demographic factors predicting birth in health facility. *Ethiopian Journal of Health Sciences* 24: 81–92.
- Blencowe H, Cousens S, Jassir FB *et al.* 2016. National, regional, and worldwide estimates of stillbirth rates in 2015, with trends from 2000: a systematic analysis. *The Lancet Global Health* 4: e98–108.
- Boerma T, Requejo J, Victora CG *et al.* 2018. Countdown to 2030: tracking progress towards universal coverage for reproductive, maternal, newborn, and child health. *The Lancet* 391: 1538–48.
- Bustreo F, Mshinda H, Hinton R, Hausmann-Muela S, Tanner M. 2019. Commentary: primary health care in Tanzania - leading the way through innovation. *EClinicalMedicine* 13: 12–3.

- Campbell OMR, Calvert C, Testa A *et al.* 2016. The scale, scope, coverage, and capability of childbirth care. *The Lancet* 388: 2193–2208.
- Dennis ML, Benova L, Abuya T *et al.* 2019. Initiation and continuity of maternal healthcare: examining the role of vouchers and user-fee removal on maternal health service use in Kenya. *Health Policy and Planning* 34: 120–31.
- Doctor HV, Radovich E, Benova L. 2019. Time trends in facility-based and private-sector childbirth care: analysis of demographic and health surveys from 25 sub-Saharan African countries from 2000 to 2016. *Journal of Global Health* 9: 020406.
- Dominicus DA, Akamatsu T. 1989. Health policy and implementations in Tanzania. *The Keio Journal of Medicine* 38: 192–200.
- Dunlop CL, Benova L, Campbell O. 2018. Effect of maternal age on facility-based delivery: analysis of first-order births in 34 countries of sub-Saharan Africa using demographic and health survey data. *BMJ Open* 8: e02023.
- Filippi V, Chou D, Ronsmans C, Graham W, Say L. 2016. Levels and causes of maternal mortality and morbidity. In: Black RE, Laxminarayan R, Temmerman M, Walker N (eds). *Reproductive, Maternal, Newborn, and Child Health: Disease Control Priorities*. Vol. 2, 3rd edn. Washington, DC: The International Bank for Reconstruction and Development/The World Bank, 51–70.
- Fogliati P, Straneo M, Brogi C *et al.* 2015. How can childbirth care for the rural poor be improved? A contribution from spatial modelling in rural Tanzania. *PLoS One* 10: e0139460.
- Fogliati P, Straneo M, Mangi S *et al.* 2017. A new use for an old tool: maternity waiting homes to improve equity in rural childbirth care. Results from a cross-sectional hospital and community survey in Tanzania. *Health Policy and Planning* 10: 1354–60.
- Gabrysch S, Campbell OMR. 2009. Still too far to walk: literature review of the determinants of delivery service use. *BMC Pregnancy and Childbirth* 9: 34.
- Gabrysch S, Nesbitt RC, Schoeps A *et al.* 2019. Does facility birth reduce maternal and perinatal mortality in Brong Ahafo, Ghana? A secondary analysis using data on 119 244 pregnancies from two cluster-randomised controlled trials. *Lancet Global Health* 7: e1074–87.
- Graham W, Wood S, Byass P *et al.* 2016. Diversity and divergence: the dynamic burden of poor maternal health. *Lancet* 388: 2164–75.
- Hanson C, Cox J, Mbaruku G *et al.* 2015. Maternal mortality and distance to facility-based obstetric care in rural southern Tanzania: a secondary analysis of cross-sectional census data in 226 000 households. *The Lancet Global Health* 3: e387–95.
- Hanson C, Munjanja S, Binagwaho A *et al.* 2019. National policies and care provision in pregnancy and childbirth for twins in Eastern and Southern Africa: a mixed-methods multi-country study. *PLoS Medicine* 16: e1002749.
- Hanson C, Ronsmans C, Penfold S *et al.* 2013. Health system support for childbirth care in Southern Tanzania: results from a health facility census. *BMC Research Notes* 6: 435.
- Hanson C, Schellenberg J. 2019. Redesigning maternal health services: is centralisation the answer in low-resource settings? *BMJ Global Health* 4: e001488.
- Houweling TAJ, Ronsmans C, Campbell OMR, Kunst AE. 2007. Huge poor-rich inequalities in maternity care: an international comparative study of maternity and child care in developing countries. *Bulletin of the World Health Organization* 85: 745–54.
- Jahn A, Kowalewski M, Kimatta SS. 1998. Obstetric care in southern Tanzania: does it reach those in need? *Tropical Medicine and International Health* 3: 926–32.
- Kinney MV, Kerber KJ, Black RE *et al.* On behalf of the Science in Action: Saving the lives of Africa's mothers, newborns, and children working group. 2010. Sub-Saharan Africa's mothers, newborns, and children: where and why do they die? *PLoS Medicine* 7: e1000294.
- Kowalewski M, Jahn A, Kimatta SS. 2000. Why do at-risk mothers fail to reach referral level? Barriers beyond distance and cost. *African Journal of Reproductive Health* 4: 100–9.
- Kruk ME, Gage AD, Arsenault C *et al.* 2018. High-quality health systems in the Sustainable Development Goals era: time for a revolution. *The Lancet Global Health* 6: e1196–252.
- Kruk ME, Hermosilla S, Larson E, Mbaruku GM. 2014. Bypassing primary care clinics for childbirth: a cross-sectional study in the Pwani region, United Republic of Tanzania. *Bulletin of the World Health Organization* 92: 246–53.
- Kruk ME, Leslie HH, Verguet S *et al.* 2016. Quality of basic maternal care functions in health facilities of five African countries: an analysis of national health system surveys. *The Lancet Global Health* 4: 1–11.
- Kyei-Nimakoh M, Carolan-Olah M, McCann TV. 2017. Access barriers to obstetric care at health facilities in sub-Saharan Africa—a systematic review. *Systematic Reviews* 6: 110.
- Lawn JE, Blencowe H, Waiswa P *et al.* 2016. Stillbirths: rates, risk factors, and acceleration towards 2030. *The Lancet* 387: 1–17.
- Lohela TJ, Nesbitt RC, Pekkanen J, Gabrysch S. 2019. Comparing socioeconomic inequalities between early neonatal mortality and facility delivery: cross-sectional data from 72 low- and middle-income countries. *Scientific Reports* 9: 9786.
- Marsh AD, Muzigaba M, Diaz T *et al.* 2020. Effective coverage measurement in maternal, newborn, child, and adolescent health and nutrition: progress, future prospects, and implications for quality health systems. *Lancet Global Health* 8: e730–6.
- Mgaya AH, Massawe SN, Kidanto HL, Mgaya HN. 2013. Grand multiparity: is it still a risk in pregnancy? *BMC Pregnancy and Childbirth* 13: 241.
- Montagu D, Sudhinaraset M, Diamond-Smith N *et al.* 2017. Where women go to deliver: understanding the changing landscape of childbirth in Africa and Asia. *Health Policy and Planning* 122: 1045–9.
- Moyer CA, Mustafa A. 2013. Drivers and deterrents of facility delivery in sub-Saharan Africa: a systematic review. *Reproductive Health* 10: 40.
- Nyamtema AS, Mwakatundu N, Dominico S *et al.* 2016. Enhancing maternal and perinatal health in under-served remote areas in Sub-Saharan Africa: a Tanzanian model. *PLoS One* 11: e0151419.
- Pembe AB, Urassa DP, Darj E, Carlsted A, Olsson P. 2008. Qualitative study on maternal referrals in rural Tanzania: decision making and acceptance of referral advice. *African Journal of Reproductive Health* 12: 120–31.
- Roder-Dewan S, Nimako K, Twum-Danso NAY *et al.* 2020. Health system redesign for maternal and newborn survival: rethinking care models to close the global equity gap. *BMJ Global Health* 5: e002539.
- Ronsmans C, Graham WJ. 2006. Maternal mortality: who, when, where, and why. *Lancet* 368: 1189–200.
- Straneo M, Fogliati P, Azzimonti G, Mangi S, Kisika F. 2014. Where do the rural poor deliver when high coverage of health facility delivery is achieved? Findings from a community and hospital survey in Tanzania. *PLoS One* 9: e113995.
- Straneo M, Fogliati P, Pellis I *et al.* 2016. On the way to universal coverage of maternal services in Iringa rural district in Tanzania. Who is yet to be reached? *African Health Sciences* 16: 420–8.
- Tanzania MoHCDGEC, MoHMZ, National Bureau of Statistics (NBS), Office of the Chief Government Statistician (OCGS), and ICF. 2016. *Tanzania Demographic and Health Survey and Malaria Indicator Survey (TDHS-MIS) 2015–16*. Dar es Salaam, and Rockville, MD: MoHCDGEC, MoH, NBS, OCGS, and ICF.
- Tanzania MoHSW. 2008. Reproductive and Child Health Section. In: *The National Road Map Strategic Plan to Accelerate Reduction of Maternal, Newborn and Child Deaths in Tanzania, 2008–15*. Dar es Salaam: MoHSW.
- Tanzania MoHSW, MoHMZ, National Bureau of Statistics (NBS), Office of the Chief Government Statistician (OCGS), and ICF. 2015.

- Tanzania Service Provision Assessment Survey (TSPA) 2014-15. Dar es Salaam: NBS, MOF, OCGS, Ministry of State, President's Office.
- Tanzania NBS and Zanzibar OGCS. 2013. *Census 2012: population distribution by age and sex*.
- Tanzania MoHSW. 2007. *Primary Health Services Development Programme*, pp. 2007-17.
- Tanzania URO. 2000. *Vision_2025_en.pdf*. Dar es Salaam Tanzania. www.tanzania.go.tz, accessed 14 January 2020.
- Tudor Hart J. 1971. The inverse care law. *The Lancet* 297: 405-12.
- UNDP. 2020. <https://population.un.org/wpp/DataQuery/>, accessed 22 July 2020.
- Virgo S, Gon G, Cavallaro FL, Graham W, Woodd S. 2017. Who delivers where? The effect of obstetric risk on facility delivery in East Africa. *Tropical Medicine & International Health* 22: 1081-98.
- Vyas S, Kumaranayake L. 2006. Constructing socio-economic status indices: how to use principal components analysis. *Health Policy and Planning* 21: 459-68.
- WHO. 2008. *Closing the Gap in a Generation. Health Equity through Action on the Social Determinants of Health. Final Report of the Commission on the Social Determinants of Health*. Geneva: World Health Organization.
- WHO. 2015a. *State of Inequality. Reproductive, Maternal, Newborn and Child Health*.
- WHO. 2015b. *Trends in Maternal Mortality: 1990 to 2015: Estimates by WHO, UNICEF, UNFPA, World Bank Group and the United Nations Population Division*. www.who.int, accessed 16 February 2021.
- WHO. 2019. *Trends in Maternal Mortality 2000 to 2017: Estimates by WHO, UNICEF, UNFPA, World Bank Group and the United Nations Population Division*. Geneva: World Health Organization.
- Wong KLM, Brady OJ, Campbell OMR, Banke-Thomas A, Benova L. 2020. Too poor or too far? Partitioning the variability of hospital-based childbirth by poverty and travel time in Kenya, Malawi, Nigeria and Tanzania. *International Journal for Equity in Health* 19: 15.
- Wong KL, Brady OJ, Campbell OMR *et al.* 2019. Current realities versus theoretical optima: quantifying efficiency and sociospatial equity of travel time to hospitals in low-income and middle-income countries. *BMJ Global Health* 4: e001552.
- World Bank. 2020. *World Bank Open Data*. <https://data.worldbank.org>, accessed 10 April 2020.