

## Section 4: Oscillations in $\beta$ -Cells

# Control Mechanisms of the Oscillations of Insulin Secretion In Vitro and In Vivo

Patrick Gilon, Magalie A. Ravier, Jean-Christophe Jonas, and Jean-Claude Henquin

The mechanisms driving the pulsatility of insulin secretion in vivo and in vitro are still unclear. Because glucose metabolism and changes in cytosolic free  $\text{Ca}^{2+}$  ( $[\text{Ca}^{2+}]_c$ ) in  $\beta$ -cells play a key role in the control of insulin secretion, and because oscillations of these two factors have been observed in single isolated islets and  $\beta$ -cells, pulsatile insulin secretion could theoretically result from  $[\text{Ca}^{2+}]_c$  or metabolism oscillations. We could not detect metabolic oscillations independent from  $[\text{Ca}^{2+}]_c$  changes in  $\beta$ -cells, and imposed metabolic oscillations were poorly effective in inducing oscillations of secretion when  $[\text{Ca}^{2+}]_c$  was kept stable, which suggests that metabolic oscillations are not the direct regulator of the oscillations of secretion. By contrast, tight temporal and quantitative correlations between the changes in  $[\text{Ca}^{2+}]_c$  and insulin release strongly suggest that  $[\text{Ca}^{2+}]_c$  oscillations are the direct drivers of insulin secretion oscillations. Metabolism may play a dual role, inducing  $[\text{Ca}^{2+}]_c$  oscillations (via changes in ATP-sensitive  $\text{K}^+$  channel activity and membrane potential) and amplifying the secretory response by increasing the efficiency of  $\text{Ca}^{2+}$  on exocytosis. The mechanisms underlying the oscillations of insulin secretion by the isolated pancreas and those observed in vivo remain elusive. It is not known how the functioning of distinct islets is synchronized, and the possible role of intrapancreatic ganglia in this synchronization requires confirmation. That pulsatile insulin secretion is beneficial in vivo, by preventing insulin resistance, is suggested by the greater hypoglycemic effect of exogenous insulin when it is infused in a pulsatile rather than continuous manner. The observation that type 2 diabetic patients have impaired pulsatile insulin secretion has prompted the suggestion that such dysregulation contributes to the disease and justifies the efforts toward understanding of the mechanism underlying the pulsatility of insulin secretion both in vitro and in vivo. *Diabetes* 51 (Suppl. 1):S144–S151, 2002

Oscillations occurring at highly variable rhythms characterize many biological events such as contraction, neurotransmission, and secretion. At the cellular level, they may involve oscillations of signals as diverse as membrane potential,  $\text{Ca}^{2+}$  concentration, metabolism and activity of protein kinases, protein phosphatases, or pumps. Insulin secretion is also a pulsatile phenomenon. The mechanisms underlying this pulsatility have been widely studied but are still a matter of debate. They will be briefly reviewed in this article, which complements recent articles on the same subject (1,2) but presents different interpretations.

**In vivo oscillations of insulin secretion.** The plasma insulin concentration oscillates even during postabsorptive periods (3–6). Two major frequencies characterize these oscillations. Ultradian oscillations (period of  $\sim 120$  min) probably result from a feedback loop between glucose production and insulin secretion and will not be further discussed here (7,8). There also exist faster oscillations that were initially reported to have a period of  $\sim 15$  min in peripheral blood (3–5,9), and turned out later to have a period between 5 and 10 min in portal blood (10,11) or in peripheral blood using deconvolution algorithms (6,12). The finding that the C-peptide concentration oscillates nearly in phase with insulin indicates that cyclic secretion rather than cyclic clearance produces the oscillations of insulinemia (13–15). Pulsatile insulin secretion has been reported to account for  $\sim 70\%$  of total insulin secretion in normal human subjects (6,10), and the major effect of glucose is to increase the amplitude of the oscillations (11,12,16,17).

**In vitro oscillations of insulin secretion.** The first convincing evidence for pulsatile insulin secretion in vitro was obtained with the isolated dog pancreas perfused with a constant concentration of glucose (18,19). The oscillations had a period of 5–10 min. Similar findings were made with the isolated pancreases of other species including monkeys (20) and humans (21). Oscillations of insulin secretion from groups of 12–200 islets have also been reported; their period was 5–10 min (22–24) or 15–30 min (25–28). Monitoring insulin secretion from single islets requires a sensitive radioimmunoassay (29) or enzyme-linked immunosorbent assay (30), or uses an indirect approach, based on the amperometric detection of 5-hydroxytryptamine (5-HT) that has accumulated in insulin granules during preincubation of the islet with the amine

From the Unité d'Endocrinologie et Métabolisme, University of Louvain Faculty of Medicine, Brussels, Belgium.

Address correspondence and reprint requests to gilon@endo.ucl.ac.be.

Accepted for publication 13 June 2001.

5-HT, 5-hydroxytryptamine;  $[\text{Ca}^{2+}]_c$ , cytosolic free  $\text{Ca}^{2+}$  concentration;  $\text{IP}_3$ , inositol 1,4,5-trisphosphate;  $\text{K}_{\text{ATP}}$  channel, ATP-sensitive  $\text{K}^+$  channel.

The symposium and the publication of this article were made possible by an unrestricted educational grant from Servier, Paris.




FIG. 1. Various patterns of  $[Ca^{2+}]_c$  oscillations induced by 10 or 15 mmol/l glucose (G10 or G15) in single  $\beta$ -cells (A, B), clusters of 5–10 islet cells (C, D), and single islets (E–G) from normal mice. Slow oscillations were present in the three types of preparations. Mixed oscillations were rare in single cells, more common in clusters of islet cells, and frequent in islets. Regular and rapid  $[Ca^{2+}]_c$  oscillations were seen only in islets.

(31). Depending on the sampling frequency, variable periods were reported for pulsatile insulin secretion from the islet. Thus, oscillations with a period of 2–5 min were detected with a sampling interval of 10–20 s (29,30,32,33). These oscillations could be resolved into faster oscillations (period of 12–30 s) when the sampling interval was decreased to 3 s (32,34) or by using the amperometric real-time measurement of 5-HT release (35). Glucose increased the amplitude of the oscillations of the perfused pancreas (18) and of single islets (30,33,36) or groups of islets (24). At the single-cell level, several methods can monitor exocytosis of insulin granules with an excellent temporal resolution. They include the amperometric detection of 5-HT (37,38) or insulin itself (39),  $\beta$ -cell capacitance recording (38), and fluorimetric measurements of either quinacrine accumulated in secretory granules (40) or  $Zn^{2+}$  coreleased with insulin (41). Except in one study (37), no spontaneous oscillations of secretion were observed during glucose stimulation of single  $\beta$ -cells, probably because these techniques only detect large changes, cannot be used for long recordings, or put major constraints on the experimental protocols.

**Mechanisms of control of insulin secretion.** Glucose stimulates insulin secretion by activating two pathways that require metabolism of the sugar by  $\beta$ -cells (42). The triggering pathway, also referred to as the  $K_{ATP}$  channel-

dependent pathway, depends on a rise in the ATP/ADP ratio that closes ATP-sensitive  $K^+$  channels ( $K_{ATP}$  channels) in the plasma membrane. The resulting decrease in  $K^+$  permeability leads to membrane depolarization, opening of voltage-dependent  $Ca^{2+}$  channels,  $Ca^{2+}$  influx, and eventual rise of the cytosolic  $Ca^{2+}$  concentration ( $[Ca^{2+}]_c$ ) that triggers exocytosis. The amplifying pathway, also referred to as the  $K_{ATP}$  channel-independent pathway, depends on an already elevated  $[Ca^{2+}]_c$  but does not involve a further change in  $[Ca^{2+}]_c$ . It increases the efficiency of  $Ca^{2+}$  on exocytosis. The pulsatility of insulin secretion might result from oscillations in either of these transduction pathways. Because metabolism and  $[Ca^{2+}]_c$  play key roles in the control of insulin secretion and have been reported to oscillate, many efforts have been spent to investigate which of these two mechanisms is the primum movens of pulsatile insulin secretion.

**$[Ca^{2+}]_c$  oscillations in  $\beta$ -cells.** Isolated pancreatic islets have widely been used to study the effects of glucose on  $\beta$ -cell  $[Ca^{2+}]_c$ . Three types of  $[Ca^{2+}]_c$  oscillations can be observed during stimulation by the sugar (Fig. 1E–G): regular and rapid (frequency of 2–3/min), regular and slow (frequency of  $\sim 0.2$ /min), or mixed, characterized by rapid oscillations superimposed on slow ones (35,43,44). Oscillations of  $[Ca^{2+}]_c$  are also detected in islets in vivo (45). Image analysis has demonstrated that the changes in


[Ca<sup>2+</sup>]<sub>c</sub> are synchronized between all β-cells within an islet (46,47). [Ca<sup>2+</sup>]<sub>c</sub> also oscillates in glucose-stimulated single β-cells (2,48), but these oscillations are much less regular than those in islets and often occur at a low frequency only (Fig. 1A) (49). Mixed oscillations also occur in isolated cells (49) (Fig. 1B), suggesting that this peculiar pattern does not necessarily result from the sum of different signals produced in distinct β-cell populations of the islet (50). [Ca<sup>2+</sup>]<sub>c</sub> oscillations are more regular in clusters of islet cells than in single β-cells, and this regularity (not the frequency) increases with the number of coupled cells (Fig. 1C and D) (49).

The essential role of Ca<sup>2+</sup> influx in the generation of [Ca<sup>2+</sup>]<sub>c</sub> oscillations by glucose, in either whole islets or single β-cells, is demonstrated by their abrogation upon omission of extracellular Ca<sup>2+</sup> (2,47) or blockade of voltage-dependent Ca<sup>2+</sup> channels (51). [Ca<sup>2+</sup>]<sub>c</sub> oscillations are linked to oscillations of the membrane potential in β-cells (46,47), and it is assumed that mixed [Ca<sup>2+</sup>]<sub>c</sub> oscillations result from an irregular (so-called "periodic") electrical activity (35,52,53). Synchronization of the β-cell electrical activity (54) by gap junctions is likely to underlie the synchronization of [Ca<sup>2+</sup>]<sub>c</sub> oscillations between β-cells within the islet (47,49,55). Synchronization by extracellular signals has been proposed (56,57), but our experiments do not support this hypothesis (58).

The endoplasmic reticulum also contributes to glucose-induced [Ca<sup>2+</sup>]<sub>c</sub> oscillations. We have recently shown that it dampens the amplitude of the oscillations by rapidly taking up Ca<sup>2+</sup> during the upstroke phase and releasing Ca<sup>2+</sup> during the descending phase (59). Our model contrasts with a more widely received opinion that the endoplasmic reticulum amplifies glucose-induced [Ca<sup>2+</sup>]<sub>c</sub> oscillations by releasing Ca<sup>2+</sup> during the periods of depolarization in β-cells (50,60–63). Such a release could be triggered by depolarization (depolarization-induced Ca<sup>2+</sup> release, mediated by type 1 ryanodine receptors), by inositol 1,4,5-trisphosphate (IP<sub>3</sub>-induced Ca<sup>2+</sup> release), or by Ca<sup>2+</sup> itself (Ca<sup>2+</sup>-induced Ca<sup>2+</sup> release, mediated by type 2 or type 3 ryanodine receptors or IP<sub>3</sub> receptors). Because amplification of the Ca<sup>2+</sup> signal through some of these mechanisms usually requires the presence of caffeine or cAMP-producing agents (50,62,63), it is unclear whether it is operative during stimulation by glucose alone.


#### Mechanisms of the oscillations of insulin secretion by the islet

**The Ca<sup>2+</sup> hypothesis.** Because a rise in β-cell [Ca<sup>2+</sup>]<sub>c</sub> is required for glucose to stimulate insulin secretion (64), and because glucose-induced [Ca<sup>2+</sup>]<sub>c</sub> oscillations occur synchronously within all β-cells of an islet, it has been proposed that oscillations of insulin secretion are driven by [Ca<sup>2+</sup>]<sub>c</sub> oscillations. This proposal was directly supported by simultaneous measurements of insulin secretion and [Ca<sup>2+</sup>]<sub>c</sub> in the same islet. They showed that each [Ca<sup>2+</sup>]<sub>c</sub> oscillation induced by glucose is accompanied by a synchronous oscillation of insulin secretion even when the frequency of the oscillations is modified (Fig. 2) (29,31,35,36,65). A similar temporal correlation was observed at the single-cell level (37). It also holds when the mitochondrial substrate, α-ketoglutarate, or the hypoglycemic sulfonylurea tolbutamide are used as stimuli (29,66).



**FIG. 2.** Effect of glucose on the oscillations of [Ca<sup>2+</sup>]<sub>c</sub> and insulin secretion measured simultaneously in single mouse islets. **A:** During perfusion with 15 mmol/l glucose and 10 mmol/l Ca<sup>2+</sup> (G15–Ca10), the islet displayed regular oscillations of [Ca<sup>2+</sup>]<sub>c</sub>. Frequent sampling (every 10 s) of the effluent for insulin assay made it possible to follow the associated oscillations of insulin secretion. **B:** During perfusion with 12 mmol/l glucose and 2.5 mmol/l Ca<sup>2+</sup> (G12–Ca2.5), the islet displayed mixed oscillations of [Ca<sup>2+</sup>]<sub>c</sub>. The slower sampling rate (every 30 s) only permitted detection of the slow insulin secretion oscillations. **C:** Increasing the glucose (G) concentration of the medium from 15 to 30 mmol/l did not affect the peak of [Ca<sup>2+</sup>]<sub>c</sub> oscillations but increased that of insulin secretion oscillations, indicating that glucose increases the efficiency of [Ca<sup>2+</sup>]<sub>c</sub> on secretion. The Ca<sup>2+</sup> concentration of the perfusion medium was 10 mmol/l throughout. Insulin was measured in fractions of effluent collected every 10 s (adapted from Gilon and Henquin [36]).

When [Ca<sup>2+</sup>]<sub>c</sub> was forced to oscillate by imposed, repetitive depolarizations of β-cells with 30 mmol/l K<sup>+</sup> (in the presence of diazoxide and 15 mmol/l glucose), insulin secretion also oscillated. In contrast, when [Ca<sup>2+</sup>]<sub>c</sub> was stably elevated by 30 mmol/l K<sup>+</sup>, no oscillations of insulin secretion were detected (Fig. 3) (67). Although these data suggest that Ca<sup>2+</sup> is the oscillophore and the moment-to-moment driver of insulin secretion, the hypothesis was challenged by reports of dissociations between [Ca<sup>2+</sup>]<sub>c</sub>



**FIG. 3.** Insulin secretion does not oscillate at high and stable  $[Ca^{2+}]_c$ . Single mouse islets were perfused with a medium containing 15 mmol/l glucose (G) and 250  $\mu$ mol/l diazoxide (Dz) throughout.  $[Ca^{2+}]_c$  was varied by changing intermittently the  $K^+$  concentration of the medium between 4.8 and 30 mmol/l as indicated on the top of each panel.  $[Ca^{2+}]_c$  and insulin secretion were measured simultaneously. Rapid (every 12 s, *A*) or slow (every 30 s, *B*) sampling of the effluent for insulin assay did not reveal oscillations of insulin secretion at high and sustained  $[Ca^{2+}]_c$ . (Adapted from Jonas et al. [67].)


changes and insulin secretion. In *ob/ob* mouse islets, insulin secretion was found to oscillate at basal and stable  $[Ca^{2+}]_c$  when glucose was omitted from the medium, in the absence or presence of clonidine or diazoxide (68). Oscillations of insulin secretion were also observed during sustained and stable elevation of  $[Ca^{2+}]_c$  with high tolbutamide,  $K^+$ , or glucose concentrations in *ob/ob* and rat islets (24,33,69). Because many of the experiments reporting dissociations between  $[Ca^{2+}]_c$  and insulin secretion were performed with islets from *ob/ob* mice, we recently investigated the possibility that the islets from hypoleptinemic, hyperglycemic, and hyperinsulinemic *ob/ob* mice differ from those of normal mice (70). We indeed found that *ob/ob* mouse islets are more prone than islets from young normal mice to display fluctuations of insulin secretion at stably elevated  $[Ca^{2+}]_c$ . However, even in these islets, the fraction of total insulin secretion that occurred in a pulsatile manner did not exceed 10–13%, which is far less than the proportion of pulsatile secretion when  $[Ca^{2+}]_c$  oscillates in  $\beta$ -cells (70).

**The metabolic hypothesis.** Oscillations of different met-

abolic variables, such as  $O_2$  consumption, ATP/ADP ratio, NAD(P)H, dihydroxyacetone-phosphate, glucose 6-phosphate, and lactate release, have been observed in intact or permeabilized normal or clonal insulin-secreting cells (22,27,71–77). These oscillations have frequencies compatible with those of insulin secretion. The identification of these metabolic oscillations and the observation of pulsatile insulin secretion independent from  $[Ca^{2+}]_c$  changes (24,33,68,69) prompted the suggestion of metabolism-driven oscillations of insulin secretion (1,78).

Oscillations of a metabolic signal in  $\beta$ -cells may result from intrinsic properties of various pathways (for example glycolysis, see [78]) or be secondary to oscillations of  $[Ca^{2+}]_c$  (79–81). In the first case only can the metabolic oscillations be regarded as the true regulators of the oscillations of secretion. This metabolic regulation can be achieved in two ways. Through an action on  $K_{ATP}$  channels, intrinsic metabolic oscillations may induce oscillations of the membrane potential and  $Ca^{2+}$  influx, but  $[Ca^{2+}]_c$  oscillations remain the ultimate effector of the insulin pulsatility. Alternatively, intrinsic metabolic oscillations may modulate the action of  $Ca^{2+}$  on insulin secretion through the amplifying pathway. The proof of this second possibility requires the demonstration that oscillations of a metabolic signal do occur and drive large oscillations of insulin secretion in the absence of  $[Ca^{2+}]_c$  changes. However, only few experiments performed with intact cells support the existence of metabolic oscillations at stable  $[Ca^{2+}]_c$ . Thus, oscillations of the  $K_{ATP}$  channel activity were observed in intact single  $\beta$ -cells perfused with 3 mmol/l glucose, when  $[Ca^{2+}]_c$  was presumably low and stable. Oscillations of  $O_2$  consumption nearly in phase with oscillations of insulin secretion were detected in the presence of 3 mmol/l glucose or of a combination of 11 mmol/l glucose and a high concentration of tolbutamide (1 mmol/l) (77). Strangely, increasing the glucose concentration in the medium did not change the amplitude of the oscillations of  $O_2$  consumption but markedly increased that of the oscillations of insulin secretion. In single clonal  $\beta$ -cells (HIT), oscillations of  $O_2$  consumption were detected in the presence, not in the absence, of glucose. They had a period of  $\sim 3$  min, similar to that of insulin secretion oscillations, and persisted at the same frequency in a  $Ca^{2+}$ -free medium (76). It is important to keep in mind that clonal cells might behave differently from normal  $\beta$ -cells, because their glycolysis is preferentially anaerobic whereas glycolysis in normal  $\beta$ -cells is preferentially aerobic (82). Different results were obtained by simultaneous measurements of  $[Ca^{2+}]_c$  and metabolic parameters in single intact mouse islets. Oscillations of glucose and  $O_2$  consumption were detected only in the presence of stimulating concentrations of glucose (74). These oscillations were synchronous to those of  $[Ca^{2+}]_c$  and abolished by omission of  $Ca^{2+}$  or damped by addition of nifedipine (74,75). They thus seem to correspond to  $Ca^{2+}$ -induced metabolic oscillations.

Other observations also cast doubts on the existence of metabolic oscillations independent from  $[Ca^{2+}]_c$  changes. We did not detect oscillations of the  $K_{ATP}$  channel activity in single metabolically intact  $\beta$ -cells under conditions of low and stable  $[Ca^{2+}]_c$  (83). No oscillations of ATP concentration were detected in the mitochondrial and cytosol-



**FIG. 4.** Imposed metabolic oscillations are much less effective than imposed [Ca<sup>2+</sup>]<sub>c</sub> oscillations on insulin secretion. **A:** Insulin secretion was measured (every 30 s) simultaneously either with NAD(P)H or [Ca<sup>2+</sup>]<sub>c</sub> in single islets. The perfusion medium contained 30 mmol/l K<sup>+</sup> and 250 μmol/l diazoxide (Dz) throughout. The glucose (G) concentration was changed between 10 and 20 mmol/l as indicated on the top of the panel. Values are means ± SE for eight experiments of insulin secretion, of which four were combined with [Ca<sup>2+</sup>]<sub>c</sub> and four with NAD(P)H measurements. **B:** NAD(P)H, [Ca<sup>2+</sup>]<sub>c</sub>, and insulin secretion were measured separately. The islets were perfused with 10 mmol/l glucose (G) and 250 μmol/l diazoxide (Dz) throughout. The K<sup>+</sup> concentration was changed between 4.8 and 30 mmol/l as indicated on the top of the panel. All values in **A** and **B** are expressed as a percentage of the values measured within each experiment during the last minute preceding the first application of 20 mmol/l glucose (**A**) or the first depolarization with 30 mmol/l K<sup>+</sup> (**B**). [Ca<sup>2+</sup>]<sub>c</sub> corresponding to 100% were 222 ± 11 and 135 ± 33 nmol/l in **A** and **B**, respectively. Insulin secretion corresponding to 100% were 257 ± 32 and 12 ± 2 pg/min/islet in **A** and **B**, respectively. (Adapted from Ravier et al. [124].)

lic compartments of glucose-stimulated rat islets transfected with adenoviruses and expressing targeted luciferase (84). With the exception of one group that reported oscillations of NAD(P)H fluorescence in single islet cells (40), most authors do not detect NAD(P)H oscillations in islets that display [Ca<sup>2+</sup>]<sub>c</sub> oscillations under the same conditions (47,85).

If Ca<sup>2+</sup>-independent metabolic oscillations really exist in β-cells and play a role in the control of insulin secretion, models other than glycolytic oscillations will be required to explain the observation that the mitochondrial substrate α-ketoglutarate can also induce oscillations of insulin secretion (66).

**[Ca<sup>2+</sup>]<sub>c</sub> is the main driver of the oscillations of insulin secretion.** To establish whether metabolic or [Ca<sup>2+</sup>]<sub>c</sub> oscillations are the main driver of the oscillations of insulin secretion, each parameter was forced to oscillate while the other was stabilized as much as possible. In the presence of diazoxide, to prevent the effect of glucose on the membrane potential, alternating between 10 and 20 mmol/l glucose in the continuous presence of 30 mmol/l K<sup>+</sup> was without effect on [Ca<sup>2+</sup>]<sub>c</sub> but induced oscillations of metabolism [NAD(P)H fluorescence]. This resulted in small oscillations of insulin secretion (Fig. 4A). Conversely, when the islets were alternatively perfused with 4.8 and 30 mmol/l K<sup>+</sup> in the continuous presence of 10 mmol/l glucose and diazoxide, large [Ca<sup>2+</sup>]<sub>c</sub> oscillations occurred, accompanied by only minor oscillations of NAD(P)H fluorescence. However, the resulting oscillations

of insulin secretion had an amplitude ~100 times larger than that of the oscillations of secretion induced by imposed metabolic oscillations (Fig. 4B). In a model of permeabilized cells, oscillations of the Ca<sup>2+</sup> concentration in the medium (hence within the cells) triggered oscillations of insulin secretion in the presence of a stable concentration of ATP and without possible glucose metabolism (86). Altogether, these data demonstrate that metabolic oscillations are poor effectors compared with [Ca<sup>2+</sup>]<sub>c</sub> oscillations and strongly suggest that [Ca<sup>2+</sup>]<sub>c</sub> oscillations are the direct drivers of the oscillations of insulin secretion at the islet level.

**Mechanisms of the oscillations of insulin secretion in vivo.** In vivo, insulinemia, glucagonemia, and glycemia oscillate at the same frequency. These oscillations are unaffected by agents modifying the function of the central nervous system (87). Initially, plasma insulin was reported to cycle nearly in and glucagon nearly out of phase with glucose (3,9,88,89), suggesting that pulses of insulin/glucagon secretion might induce pulses of glucose production by the liver; the resulting pulses of glycemia might in turn induce pulses of hormone release by the pancreas. Although very attractive, this hypothesis was challenged by subsequent studies that failed to find a consistent relationship between the periodic fluctuations in insulin, glucagon, and glucose (4,13,90). Moreover, suppression of glucose oscillations by a glucose clamp did not prevent the oscillations of insulinemia (91), and pulses of glucose or insulin failed to reset the cycles (17).

The observation that insulin and glucagon are secreted in a pulsatile manner by the perfused pancreas (18–21) has led to the proposal that a pacemaker system is present in the gland itself. An intrapancreatic network was suggested to pace and synchronize insulin secretion between all islets within the pancreas (92). Several observations are consistent with this proposal. Intrapaneatic neurons of the cat pancreas showed oscillations of their electrical activity with a period similar to that of the oscillations of insulin release (93). Pulsatile insulin secretion from the perfused dog (94) and rat (2) pancreas was altered by blockade of neurotransmission with tetrodotoxin or inhibition of postsynaptic nicotinic receptors (92). However, further identification of the nature of the neurotransmitter controlling the pulsatility of insulin secretion proved impossible because pharmacological blockade of endorphin, muscarinic, and α- or β-adrenergic receptors was without effect on the frequency of the oscillations in vivo (17,95) and in vitro (92). Several studies have reported oscillations of insulinemia after intrasplenic (96) or intrahepatic grafts (97,98) of isolated islets, or transplantation of a denervated pancreas (99,100). But the opinion that pulsatility of insulin secretion only reappears upon reinnervation of the grafted islets (98) is not shared by most groups (96, 97,99,100). The role of intrapancreatic ganglia in synchronizing the signals between islets is also challenged by the observation of pulses of insulin secretion by pieces of pancreas containing several islets and electrically silent ganglia (101). Despite all these studies, it remains unclear whether intrapancreatic ganglia are really the pacemaker of the pulsatility of insulin secretion in vivo. The alternative proposal, that the in vivo pulsatility of secretion reflects an intrinsic property of each islet, also has weaknesses. Thus, the electrical activity that controls [Ca<sup>2+</sup>]<sub>c</sub>

has been reported not to be synchronized between islets in situ (102). Whether insulin secretion was pulsatile under these conditions is, however, unknown.

**Functional advantages of  $[Ca^{2+}]_c$  and insulin secretion oscillations.** What is the reason for the oscillatory behavior of  $\beta$ -cells?  $[Ca^{2+}]_c$  oscillations might be energetically less costly than a sustained  $[Ca^{2+}]_c$  elevation and, by limiting the time during which  $[Ca^{2+}]_c$  is elevated, might decrease the risk of  $Ca^{2+}$ -cytotoxicity (103). This has never been documented for  $\beta$ -cells.  $[Ca^{2+}]_c$  oscillations might be important for the activation of intracellular signals that control functions other than insulin secretion, such as gene expression (104). Changes in the frequency or duration of  $[Ca^{2+}]_c$  oscillations might control insulin secretion more accurately than changes in the amplitude of a sustained  $[Ca^{2+}]_c$  rise, and avoid inactivation or downregulation of transduction or effector pathways of stimulus-secretion coupling. However, our experiments have shown that, regardless of the oscillatory or steady pattern of the  $[Ca^{2+}]_c$  elevation, insulin secretion at constant glucose is determined by the product mean  $[Ca^{2+}]_c \times \text{time}$  (67,105). No functional advantages of  $[Ca^{2+}]_c$  oscillations have yet been identified in  $\beta$ -cells, but it should be emphasized that the long-term impact of a loss of oscillations has not been evaluated.

Oscillations of insulinemia might help in preventing insulin receptor downregulation (106) and development of insulin resistance (107). Indeed, insulin infused in a pulsatile fashion had a greater hypoglycemic effect than continuously infused insulin (108–111), being more potent in reducing hepatic glucose production and stimulating glucose utilization (112–115). This superior efficacy of pulsatile insulin is not unanimously accepted (116–118), however, perhaps because its disclosure critically depends on various in vivo factors, such as glucagonemia or insulinemia or the frequency of the pulsatility (5,109,119).

Some studies (16,17,120), but not all (12,121,122), have reported disorganized oscillations of insulinemia in patients with type 2 diabetes and their near relatives with mild glucose intolerance. It is unknown, however, whether these alterations are secondary to the prolonged hyperglycemia. Insulin resistance of obese patients also decreases the regularity of pulsatile insulin secretion (123). Because oscillations of insulinemia favor optimal glucose homeostasis, one may speculate that a disturbance of plasma insulin oscillations in type 2 diabetes contributes to the disease.

#### ACKNOWLEDGMENTS

This work was supported by grant 3.4552.98 from the Fonds de la Recherche Scientifique Médicale (Brussels), grant ARC 00/05–260 from the General Direction of Scientific Research of the French Community of Belgium, and by the Interuniversity Poles of Attraction Program (P4/21), Belgian State, Federal Office for Scientific, Technical and Cultural Affairs. P.G and J.C.J. are Maître de Recherches and Chercheur Qualifié, respectively, from the Fonds National de la Recherche Scientifique, Brussels.

#### REFERENCES

- Bergsten P: Pathophysiology of impaired pulsatile insulin release. *Diabetes Met Res Rev* 16:179–191, 2000
- Gylfe E, Ahmed M, Bergsten P, Dansk H, Dyachok O, Eberhardson M,

- Grapengiesser E, Hellman B, Lin JM, Sundsten T, Tengholm A, Vieira E, Westerlund J: Signaling underlying pulsatile insulin secretion. *Ups J Med Sci* 105:35–51, 2000
- Goodner CJ, Walike BC, Koerker DJ, Ensinnck JW, Brown AC, Chideckel EW, Palmer J, Kalnasy L: Insulin, glucagon, and glucose exhibit synchronous, sustained oscillations in fasting monkeys. *Science* 195:177–179, 1977
- Weigle DS: Pulsatile secretion of fuel-regulatory hormones. *Diabetes* 36:764–775, 1987
- Lefebvre PJ, Paolisso G, Scheen AJ, Henquin JC: Pulsatility of insulin and glucagon release: physiological significance and pharmacological implications. *Diabetologia* 30:443–452, 1987
- Porksen N, Nyholm B, Veldhuis JD, Butler PC, Schmitz O: In humans at least 75% of insulin secretion arises from punctuated insulin secretory bursts. *Am J Physiol* 273:E908–E914, 1997
- Sturis J, Scheen AJ, Leproult R, Polonsky KS, Van Cauter E: 24-Hour glucose profiles during continuous or oscillatory insulin infusion: demonstration of the functional significance of ultradian insulin oscillations. *J Clin Invest* 95:1464–1471, 1995
- Polonsky KS, Sturis J, Van Cauter E: Temporal profiles and clinical significance of pulsatile insulin secretion. *Horm Res* 49:178–184, 1998
- Lang DA, Matthews DR, Peto J, Turner RC: Cyclic oscillations of basal plasma glucose and insulin concentrations in human beings. *N Engl J Med* 301:1023–1027, 1979
- Porksen N, Munn S, Steers J, Vore S, Veldhuis J, Butler P: Pulsatile insulin secretion accounts for 70% of total insulin secretion during fasting. *Am J Physiol* 269:E478–E488, 1995
- Song SH, McIntyre SS, Shah H, Veldhuis JD, Hayes PC, Butler PC: Direct measurement of pulsatile insulin secretion from the portal vein in human subjects. *J Clin Endocrinol Metab* 85:4491–4499, 2000
- Laedtke T, Kjems L, Porksen N, Schmitz O, Veldhuis J, Kao PC, Butler PC: Overnight inhibition of insulin secretion restores pulsatility and proinsulin/insulin ratio in type 2 diabetes. *Am J Physiol* 279:E520–E528, 2000
- Jaspan JB, Lever E, Polonsky KS, Van Cauter E: In vivo pulsatility of pancreatic islet peptides. *Am J Physiol* 251:E215–E226, 1986
- Koerker DJ, Goodner CJ, Hansen BW, Brown AC, Rubenstein AH: Synchronous, sustained oscillation of C-peptide and insulin in the plasma of fasting monkeys. *Endocrinology* 102:1649–1652, 1978
- Simon C, Follenius M, Brandenberger G: Postprandial oscillations of plasma glucose, insulin and C-peptide in man. *Diabetologia* 30:769–773, 1987
- Lang DA, Matthews DR, Burnett MA, Turner RC: Brief, irregular oscillations of basal plasma insulin and glucose concentrations in diabetic man. *Diabetes* 30:435–439, 1981
- Matthews DR, Lang DA, Burnett MA, Turner RC: Control of pulsatile insulin secretion in man. *Diabetologia* 24:231–237, 1983
- Stagner JI, Samols E, Weir GC: Sustained oscillations of insulin, glucagon and somatostatin from the isolated canine pancreas during exposure to a constant glucose concentration. *J Clin Invest* 65:939–942, 1980
- Matthews DR, Hermansen K, Connolly AA, Gray D, Schmitz O, Clark A, Orskov H, Turner RC: Greater in vivo than in vitro pulsatility of insulin secretion with synchronized insulin and somatostatin secretory pulses. *Endocrinology* 120:2272–2278, 1987
- Goodner CJ, Koerker DJ, Stagner JI, Samols E: In vitro pancreatic hormonal pulses are less regular and more frequent than in vivo. *Am J Physiol* 260:E422–E429, 1991
- Stagner JI, Samols E: The vascular order of islet cellular perfusion in the human pancreas. *Diabetes* 41:93–97, 1992
- Longo EA, Tornheim K, Deeney JT, Varnum BA, Tillotson D, Prentki M, Corkey BE: Oscillations in cytosolic free  $Ca^{2+}$ , oxygen consumption, and insulin secretion in glucose-stimulated rat pancreatic islets. *J Biol Chem* 266:9314–9319, 1991
- Marchetti P, Scharp DW, Mclear M, Gingerich R, Finke E, Olack B, Swanson C, Giannarelli R, Navalesi R, Lacy PE: Pulsatile insulin secretion from isolated human pancreatic islets. *Diabetes* 43:827–830, 1994
- Cunningham BA, Deeney JT, Bliss CR, Corkey BE, Tornheim K: Glucose-induced oscillatory insulin secretion in perfused rat pancreatic islets and clonal  $\beta$ -cells (HIT). *Am J Physiol* 271:E702–E710, 1996
- Bergstrom RW, Fujimoto WY, Teller DC, De Haen C: Oscillatory insulin secretion in perfused isolated rat islets. *Am J Physiol* 257:E479–E485, 1989
- Chou H-F, Ipp E: Pulsatile insulin secretion in isolated rat islets. *Diabetes* 39:112–117, 1990
- Chou H-F, Berman N, Ipp E: Oscillations of lactate released from islets of Langerhans: evidence for oscillatory glycolysis in  $\beta$ -cells. *Am J Physiol* 262:E800–E805, 1992

28. Yao NK, Chang LW, Lin BJ, Kuo TS: Dynamic aspects for interislet synchronization of oscillatory insulin secretions. *Am J Physiol* 272:E981–E988, 1997
29. Gilon P, Shepherd RM, Henquin JC: Oscillations of secretion driven by oscillations of cytoplasmic Ca<sup>2+</sup> as evidenced in single pancreatic islets. *J Biol Chem* 268:22265–22268, 1993
30. Bergsten P, Hellman B: Glucose-induced amplitude regulation of pulsatile insulin secretion from individual pancreatic islets. *Diabetes* 42:670–674, 1993
31. Barbosa RM, Silva AM, Tomé AR, Stamford JA, Santos RM, Rosario LM: Real time electrochemical detection of 5-HT/insulin secretion from single pancreatic islets: effect of glucose and K<sup>+</sup> depolarization. *Biochem Biophys Res Commun* 228:100–104, 1996
32. Hellman B, Gylfe E, Bergsten P, Grapengiesser E, Lund P-E, Berts A, Tengholm A, Pipeleers DG, Ling Z: Glucose induces oscillatory Ca<sup>2+</sup> signalling and insulin release in human pancreatic beta cells. *Diabetologia* 37 (Suppl. 2):S11–S20, 1994
33. Bergsten P: Glucose-induced pulsatile insulin release from single islets at stable and oscillatory cytoplasmic Ca<sup>2+</sup>. *Am J Physiol* 274:E796–E800, 1998
34. Bergsten P, Hellman B: Glucose-induced cycles of insulin release can be resolved into distinct periods of secretory activity. *Biochem Biophys Res Commun* 192:1182–1188, 1993
35. Barbosa RM, Silva AM, Tomé AR, Stamford JA, Santos RM, Rosario LM: Control of pulsatile 5-HT/insulin secretion from single mouse pancreatic islets by intracellular calcium dynamics. *J Physiol* 510:135–143, 1998
36. Gilon P, Henquin JC: Distinct effects of glucose on the synchronous oscillations of insulin release and cytoplasmic Ca<sup>2+</sup> concentration measured simultaneously in single mouse islets. *Endocrinology* 136:5725–5730, 1995
37. Smith PA, Duchon MR, Ashcroft FM: A fluorimetric and amperometric study of calcium and secretion in isolated mouse pancreatic β-cells. *Pflugers Arch* 430:808–818, 1995
38. Bokvist K, Holmqvist M, Gromada J, Rorsman P: Compound exocytosis in voltage-clamped mouse pancreatic β-cells revealed by carbon fibre amperometry. *Pflugers Arch* 439:634–645, 2000
39. Huang L, Shen H, Atkinson MA, Kennedy RT: Detection of exocytosis at individual pancreatic β cells by amperometry at a chemically modified microelectrode. *Proc Natl Acad Sci U S A* 92:9608–9612, 1995
40. Pralong WF, Bartley C, Wollheim CB: Single islet beta-cell stimulation by nutrients: relationship between pyridine nucleotides, cytosolic Ca<sup>2+</sup> and secretion. *EMBO J* 9:53–60, 1990
41. Qian WJ, Aspinwall CA, Battiste MA, Kennedy RT: Detection of secretion from single pancreatic β-cells using extracellular fluorogenic reactions and confocal fluorescence microscopy. *Anal Chem* 72:711–717, 2000
42. Henquin JC: Triggering and amplifying pathways of regulation of insulin secretion by glucose. *Diabetes* 49:1751–1760, 2000
43. Valdeolmillos M, Santos RM, Contreras D, Soria B, Rosario LM: Glucose-induced oscillations of intracellular Ca<sup>2+</sup> concentration resembling bursting electrical activity in single mouse islets of Langerhans. *FEBS Lett* 259:19–23, 1989
44. Henquin JC, Jonas JC, Gilon P: Functional significance of Ca<sup>2+</sup> oscillations in pancreatic beta cells. *Diabetes Metab* 24:30–36, 1998
45. Fernandez J, Valdeolmillos M: Synchronous glucose-dependent [Ca<sup>2+</sup>]<sub>i</sub> oscillations in mouse pancreatic islets of Langerhans recorded in vivo. *FEBS Lett* 477:33–36, 2000
46. Santos RM, Rosario LM, Nadal A, Garcia-Sancho J, Soria B, Valdeolmillos M: Widespread synchronous [Ca<sup>2+</sup>]<sub>i</sub> oscillations due to bursting electrical activity in single pancreatic islets. *Pflugers Arch* 418:417–422, 1991
47. Gilon P, Henquin JC: Influence of membrane potential changes on cytoplasmic Ca<sup>2+</sup> concentration in an electrically excitable cell, the insulin-secreting pancreatic B-cell. *J Biol Chem* 267:20713–20720, 1992
48. Grapengiesser E, Gylfe E, Hellman B: Glucose-induced oscillations of cytoplasmic Ca<sup>2+</sup> in the pancreatic beta-cell. *Biochem Biophys Res Commun* 151:1299–1304, 1988
49. Jonkers FC, Jonas JC, Gilon P, Henquin JC: Influence of cell number on the characteristics and synchrony of Ca<sup>2+</sup> oscillations in clusters of mouse pancreatic islet cells. *J Physiol* 520:839–849, 1999
50. Liu YJ, Tengholm A, Grapengiesser E, Hellman B, Gylfe E: Origin of slow and fast oscillations of Ca<sup>2+</sup> in mouse pancreatic islets. *J Physiol* 508:471–481, 1998
51. Dryselius S, Grapengiesser E, Hellman B, Gylfe E: Voltage-dependent entry and generation of slow Ca<sup>2+</sup> oscillations in glucose-stimulated pancreatic β-cells. *Am J Physiol* 276:E512–E518, 1999
52. Henquin JC, Meissner HP, Schmeer W: Cyclic variations of glucose-induced electrical activity in pancreatic B cells. *Pflugers Arch* 393:322–327, 1982
53. Cook DL: Isolated islets of Langerhans have slow oscillations of electrical activity. *Metabolism* 32:681–685, 1983
54. Meissner HP: Electrophysiological evidence for coupling between β cells of pancreatic islets. *Nature* 262:502–504, 1976
55. Gylfe E, Grapengiesser E, Hellman B: Propagation of cytoplasmic Ca<sup>2+</sup> oscillations in clusters of pancreatic β-cells exposed to glucose. *Cell Calcium* 12:229–240, 1991
56. Perez-Armentariz EM, Atwater I, Bennett MVL: Mechanisms for fast intercellular communication within a single islet of Langerhans. In *Pacemaker Activity and Intercellular Communication*. Huizinga JD, Ed. London, CRC Press, 1995, p. 305–321
57. Bertuzzi F, Davalli AM, Nano R, Socci C, Codazzi F, Fesce R, Di Carlo V, Pozza G, Grohovaz F: Mechanisms of coordination of Ca<sup>2+</sup> signals in pancreatic islet cells. *Diabetes* 48:1971–1978, 1999
58. Jonkers FC, Henquin JC: Measurements of cytoplasmic Ca<sup>2+</sup> in islet cell clusters show that glucose rapidly recruits beta-cells and gradually increases the individual cell response. *Diabetes* 50:540–550, 2001
59. Gilon P, Arredouani A, Gailly P, Gromada J, Henquin JC: Uptake and release of Ca<sup>2+</sup> by the endoplasmic reticulum contribute to the oscillations of the cytosolic Ca<sup>2+</sup> concentration triggered by Ca<sup>2+</sup> influx in the electrically excitable pancreatic B-cell. *J Biol Chem* 274:20197–20205, 1999
60. Takasawa S, Nata K, Yonekura H, Okamoto H: Cyclic ADP-ribose in insulin secretion from pancreatic β cells. *Science* 259:370–373, 1993
61. Dukes ID, Roe MW, Worley JF III, Philipson LH: Glucose-induced alterations in β-cell cytoplasmic Ca<sup>2+</sup> involving the coupling of intracellular Ca<sup>2+</sup> stores and plasma membrane ion channels. *Curr Opin Endocrinol Diab* 4:262–271, 1997
62. Holz GG, Leech CA, Heller RS, Castonguay M, Habener JF: cAMP-dependent mobilization of intracellular Ca<sup>2+</sup> stores by activation of ryanodine receptors in pancreatic β-cells: a Ca<sup>2+</sup> signaling system stimulated by the insulinotropic hormone glucagon-like peptide-1(7–37). *J Biol Chem* 274:14147–14156, 1999
63. Lemmens R, Larsson O, Berggren PO, Islam MS: Ca<sup>2+</sup>-induced Ca<sup>2+</sup> release from endoplasmic reticulum amplifies Ca<sup>2+</sup> signal mediated by activation of voltage-gated L-type Ca<sup>2+</sup> channels in pancreatic beta cells. *J Biol Chem* 276:9971–9977, 2001
64. Henquin JC, Jonas JC, Sato Y, Detimary P, Gilon P: Signal transduction: regulation of insulin secretion by changes in Ca<sup>2+</sup> concentration and action in pancreatic β-cells. In *Advances in Molecular and Cell Biology: The Biology of the Pancreatic β-Cell*. Bittar ED, Howell SL, Eds. Stamford, Connecticut, JAI Press, Vol. 29, 1999, p. 247–275
65. Bergsten P, Grapengiesser E, Gylfe E, Tengholm A, Hellman B: Synchronous oscillations of cytoplasmic Ca<sup>2+</sup> and insulin release in glucose-stimulated pancreatic islets. *J Biol Chem* 269:8749–8753, 1994
66. Martin F, Sanchez-Andres JV, Soria B: Slow [Ca<sup>2+</sup>]<sub>i</sub> oscillations induced by ketoisocaproate in single mouse pancreatic islets. *Diabetes* 44:300–305, 1995
67. Jonas JC, Gilon P, Henquin JC: Temporal and quantitative correlations between insulin secretion and stably elevated or oscillatory cytoplasmic Ca<sup>2+</sup> in mouse pancreatic β-cells. *Diabetes* 47:1266–1273, 1998
68. Westerlund J, Hellman B, Bergsten P: Pulsatile insulin release from mouse islets occurs in the absence of stimulated entry of Ca<sup>2+</sup>. *J Clin Invest* 97:1860–1863, 1996
69. Westerlund J, Gylfe E, Bergsten P: Pulsatile insulin release from pancreatic islets with nonoscillatory elevation of cytoplasmic Ca<sup>2+</sup>. *J Clin Invest* 100:2547–2551, 1997
70. Kjems LL, Ravier MA, Jonas JC, Henquin JC: Do oscillations of insulin secretion occur in the absence of cytoplasmic Ca<sup>2+</sup> oscillations in β-cells? *Diabetes* 51 (Suppl. 1):S177–182, 2002
71. Corkey BE, Tornheim K, Deeney JT, Glennon MC, Parker JC, Matschinsky FM, Ruderman NB, Prentki M: Linked oscillations of free Ca<sup>2+</sup> and the ATP/ADP ratio in permeabilized RINm5F insulinoma cells supplemented with a glycolyzing cell-free muscle extract. *J Biol Chem* 263:4254–4258, 1988
72. Nilsson T, Schultz V, Berggren PO, Corkey BE, Tornheim K: Temporal patterns of changes in ATP/ADP ratio, glucose 6-phosphate and cytoplasmic free Ca<sup>2+</sup> in glucose-stimulated pancreatic β-cells. *Biochem J* 314: 91–94, 1996
73. Civelek VN, Deeney JT, Fusonie GE, Corkey BE, Tornheim K: Oscillations in oxygen consumption by permeabilized clonal pancreatic β-cells (HIT) incubated in an oscillatory glycolyzing muscle extract: roles of free Ca<sup>2+</sup>, substrates, and the ATP/ADP ratio. *Diabetes* 46:51–56, 1997
74. Jung SK, Kauri LM, Qian WJ, Kennedy RT: Correlated oscillations in

- glucose consumption, oxygen consumption, and intracellular free  $\text{Ca}^{2+}$  in single islets of Langerhans. *J Biol Chem* 275:6642–6650, 2000
75. Jung SK, Aspinwall CA, Kennedy RT: Detection of multiple patterns of oscillatory oxygen consumption in single mouse islets of Langerhans. *Biochem Biophys Res Commun* 259:331–335, 1999
  76. Porterfield DM, Corkey RF, Sanger RH, Tornheim K, Smith PJS, Corkey BE: Oxygen consumption oscillates in single clonal pancreatic  $\beta$ -cells (HIT). *Diabetes* 49:1511–1516, 2000
  77. Ortsäter H, Liss P, Lund PE, Åkerman KEO, Bergsten P: Oscillations in oxygen tension and insulin release of individual pancreatic *ob/ob* mouse islets. *Diabetologia* 43:1313–1318, 2000
  78. Tornheim K: Are metabolic oscillations responsible for normal oscillatory insulin secretion? *Diabetes* 46:1375–1380, 1997
  79. Detimary P, Gilon P, Henquin JC: Interplay between cytoplasmic  $\text{Ca}^{2+}$  and the ATP/ADP ratio: a feedback control mechanism in mouse pancreatic islets. *Biochem J* 333:269–274, 1998
  80. Magnus G, Keizer J: Model of  $\beta$ -cell mitochondrial calcium handling and electrical activity. I. Cytoplasmic variables. *Am J Physiol* 274:C1158–C1173, 1998
  81. Kennedy HJ, Pouli AE, Ainscow EK, Jouaville LS, Rizzuto R, Rutter GA: Glucose generates sub-plasma membrane ATP microdomains in single islet  $\beta$ -cells: potential role for strategically located mitochondria. *J Biol Chem* 274:13281–13291, 1999
  82. Sener A, Blachier F, Malaisse WJ: Crabtree effect in tumoral pancreatic islet cells. *J Biol Chem* 263:1904–1909, 1988
  83. Rolland JF, Henquin JC, Gilon P: Modulation of the  $\text{K}^+$ ATP current by  $\text{Ca}^{2+}$  may contribute to oscillations of the membrane potential in B-cells. *Diabetologia* 43 (Suppl. 1):A116, 2000
  84. Ainscow EK, Rutter GA: Mitochondrial priming modifies  $\text{Ca}^{2+}$  oscillations and insulin secretion in pancreatic islets. *Biochem J* 353:175–180, 2001
  85. Panten U, Christians J, Kriegstein EV, Poser W, Hasselblatt A: Effect of carbohydrates upon fluorescence of reduced pyridine nucleotides from perfused isolated pancreatic islets. *Diabetologia* 9:477–482, 1973
  86. Jonas JC, Li G, Palmer M, Weller U, Wollheim CB: Dynamics of  $\text{Ca}^{2+}$  and guanosine 5'-[gamma-thio]triphosphate action on insulin secretion from  $\alpha$ -toxin-permeabilized HIT-T15 cells. *Biochem J* 301:523–529, 1994
  87. Hom FG, Koerker DJ, Goodner CJ: Lack of effect of morphine, reserpine, and halothane on oscillation of plasma insulin in *M. mulatta*. *Am J Physiol* 240:E1–E4, 1981
  88. Lang DA, Matthews DR, Burnett M, Ward GM, Turner RC: Pulsatile synchronous basal insulin and glucagon secretion in man. *Diabetes* 31:22–26, 1982
  89. Goodner CJ, Hom FG, Koerker DJ: Hepatic glucose production oscillates in synchrony with the islet secretory cycle in fasting rhesus monkeys. *Science* 215:1257–1260, 1982
  90. Hansen BC, Jen KLC, Pek SB, Wolfe RA: Rapid oscillations in plasma insulin, glucagon, and glucose in obese and normal weight humans. *J Clin Endocrinol Metab* 54:785–792, 1982
  91. Marsh BD, Marsh DJ, Bergman RN: Oscillations enhance the efficiency and stability of glucose disposal. *Am J Physiol* 250:E576–E582, 1986
  92. Stagner JJ, Samols E: Role of intrapancreatic ganglia in regulation of periodic insular secretions. *Am J Physiol* 248:E522–E530, 1985
  93. King BF, Love JA, Szurszewski JH: Intracellular recordings from pancreatic ganglia of the cat. *J Physiol* 419:379–403, 1989
  94. Stagner JJ, Samols E: Perturbation of insulin oscillations by nerve blockade in the in vitro canine pancreas. *Am J Physiol* 248:E516–E521, 1985
  95. Hansen BC, Pek S, Koerker DJ, Goodner CJ, Wolfe RA, Schielke GP: Neural influences on oscillations in basal plasma levels of insulin in monkeys. *Am J Physiol* 240:E5–E11, 1981
  96. Kakizaki K, Basadonna G, Merrell RC: Neural regulation of heterotopic islets of Langerhans. *Surgery* 100:997–1002, 1986
  97. Shapiro ET, Strasser S, Polonsky KS, Alejandro R, Mintz DH: Persistence of oscillatory insulin secretion in denervated islet cell autografts. *Transplantation* 52:574–576, 1991
  98. Porksen N, Munn S, Ferguson D, O'Brien T, Veldhuis J, Butler P: Coordinate pulsatile insulin secretion by chronic intraportally transplanted islets in the isolated perfused rat liver. *J Clin Invest* 94:219–227, 1994
  99. Sonnenberg GE, Hoffmann RG, Johnson CP, Kissebah AH: Low- and high-frequency insulin secretion pulses in normal subjects and pancreas transplant recipients: role of extrinsic innervation. *J Clin Invest* 90:545–553, 1992
  100. O'Meara NM, Sturis J, Blackman JD, Byrne MM, Jaspan JB, Roland DC, Thistlethwaite JR, Polonsky KS: Oscillatory insulin secretion after pancreas transplant. *Diabetes* 42:855–861, 1993
  101. Sha L, Westerlund J, Szurszewski JH, Bergsten P: Amplitude modulation of pulsatile insulin secretion by intrapancreatic ganglion neurons. *Diabetes* 50:51–55, 2001
  102. Valdeolmillos M, Gomis A, Sánchez-Andrés JV: In vivo synchronous membrane potential oscillations in mouse pancreatic  $\beta$ -cells: lack of co-ordination between islets. *J Physiol* 493:9–18, 1996
  103. Kass GEN, Orrenius S: Calcium signaling and cytotoxicity. *Environ Health Perspect* 107 (Suppl. 1):25–35, 1999
  104. Dolmetsch RE, Xu K, Lewis RS: Calcium oscillations increase the efficiency and specificity of gene expression. *Nature* 392:933–936, 1998
  105. Ravier MA, Henquin JC: Functional roles of  $\text{Ca}^{2+}$  oscillations in  $\beta$ -cells. *Diabetologia* 43 (Suppl. 1):A117, 2000
  106. Goodner CJ, Sweet IR, Harrison HC: Rapid reduction and return of surface insulin receptors after exposure to brief pulses of insulin in perfused rat hepatocytes. *Diabetes* 37:1316–1323, 1988
  107. Peiris AN, Stagner JJ, Vogel RL, Nakagawa A, Samols E: Body fat distribution and peripheral insulin sensitivity in healthy men: role of insulin pulsatility. *J Clin Endocrinol Metab* 75:290–294, 1992
  108. Matthews DR, Naylor BA, Jones RG, Ward GM, Turner RC: Pulsatile insulin has greater hypoglycemic effect than continuous delivery. *Diabetes* 32:617–621, 1983
  109. Paolisso G, Sgambato S, Passariello N, Scheen A, D'Onofrio F, Lefebvre PJ: Greater efficacy of pulsatile insulin in type I diabetics critically dependent on plasma glucagon levels. *Diabetes* 36:566–570, 1987
  110. Paolisso G, Sgambato S, Gentile S, Memoli P, Giugliano D, Varricchio M, D'Onofrio F: Advantageous metabolic effects of pulsatile insulin delivery in noninsulin-dependent diabetic patients. *J Clin Endocrinol Metab* 67:1005–1010, 1988
  111. Koopmans SJ, Sips HC, Krans HM, Radder JK: Pulsatile intravenous insulin replacement in streptozotocin diabetic rats is more efficient than continuous delivery: effects on glycaemic control, insulin-mediated glucose metabolism and lipolysis. *Diabetologia* 39:391–400, 1996
  112. Schmitz O, Arnfred J, Nielsen OH, Beck-Nielsen H, Orskov H: Glucose uptake and pulsatile insulin infusion: euglycaemic clamp and [ $^3\text{H}$ ]glucose studies in healthy subjects. *Acta Endocrinol* 113:559–563, 1986
  113. Bratusch-Marrain PR, Komjati M, Waldhausl WK: Efficacy of pulsatile versus continuous insulin administration on hepatic glucose production and glucose utilization in type I diabetic humans. *Diabetes* 35:922–926, 1986
  114. Komjati M, Bratusch-Marrain P, Waldhausl W: Superior efficacy of pulsatile versus continuous hormone exposure on hepatic glucose production in vitro. *Endocrinology* 118:312–319, 1986
  115. Paolisso G, Scheen AJ, Albert A, Lefebvre PJ: Effects of pulsatile delivery of insulin and glucagon in humans. *Am J Physiol* 257:E686–E696, 1989
  116. Verdin E, Castillo M, Luyckx AS, Lefebvre PJ: Similar metabolic effects of pulsatile versus continuous human insulin delivery during euglycemic, hyperinsulinemic glucose clamp in normal man. *Diabetes* 33:1169–1174, 1984
  117. Paolisso G, Scheen AJ, Verdin EM, Luyckx AS, Lefebvre PJ: Insulin oscillations per se do not affect glucose turnover parameters in normal man. *J Clin Endocrinol Metab* 63:520–525, 1986
  118. Kerner W, Bruckel J, Zier H, Arias P, Thun C, Moncayo R, Pfeiffer EF: Similar effects of pulsatile and constant intravenous insulin delivery. *Diabetes Res Clin Pract* 4:269–274, 1988
  119. Paolisso G, Scheen AJ, Giugliano D, Sgambato S, Albert A, Varricchio M, D'Onofrio F, Lefebvre PJ: Pulsatile insulin delivery has greater metabolic effects than continuous hormone administration in man: importance of pulse frequency. *J Clin Endocrinol Metab* 72:607–615, 1991
  120. O'Rahilly S, Turner RC, Matthews DR: Impaired pulsatile secretion of insulin in relatives of patients with non-insulin-dependent diabetes. *N Engl J Med* 318:1225–1230, 1988
  121. Mao CS, Berman N, Roberts K, Ipp E: Glucose entrainment of high-frequency plasma insulin oscillations in control and type 2 diabetic subjects. *Diabetes* 48:714–721, 1999
  122. Nyholm B, Porksen N, Juhl CB, Gravholt CH, Butler PC, Weeke J, Veldhuis JD, Pincus S, Schmitz O: Assessment of insulin secretion in relatives of patients with type 2 (non-insulin-dependent) diabetes mellitus: evidence of early  $\beta$ -cell dysfunction. *Metabolism* 49:896–905, 2000
  123. Zarkovic M, Ciric J, Stojanovic M, Penezic Z, Trbojevic B, Drezgic M, Nesovic M: Effect of insulin sensitivity on pulsatile insulin secretion. *Eur J Endocrinol* 141:494–501, 1999
  124. Ravier MA, Gilon P, Henquin JC: Oscillations of insulin secretion can be triggered by imposed oscillations of cytoplasmic  $\text{Ca}^{2+}$  or metabolism in normal mouse islets. *Diabetes* 48:2374–2382, 1999