

# Linkage for BMI at 3q27 Region Confirmed in an African-American Population

Amy Luke,<sup>1</sup> Xiaodong Wu,<sup>1</sup> Xiaofeng Zhu,<sup>1</sup> Donghui Kan,<sup>1</sup> Yan Su,<sup>2</sup> and Richard Cooper<sup>1</sup>

Significant and suggestive linkage for BMI on 3q27 has been reported by several groups, including our own study in African Americans. To further establish the linkage evidence on 3q27, we recruited an independent African-American sample comprising 545 individuals in 128 families. We genotyped 15 short tandem-repeat markers evenly spaced in the 112 cM region around the peak on 3q27 identified in our earlier study. Multipoint linkage analysis by GENEHUNTER2 gave the maximum logarithm of odds (LOD) score 2.4 at map position 188 cM in this sample. When we combined the two samples, linkage evidence was increased to a maximum LOD score (MLS) of 4.3 (point-wise  $P = 4.34 \times 10^{-6}$ ) at 188 cM, with a 7 cM 1-LOD-drop interval around the peak. The multiple replications of linkage evidence in the region on 3q27 strongly confirm its potential importance as a candidate region in the search for obesity-related genes. *Diabetes* 52:1284–1287, 2003

Genome-wide linkage analyses for obesity have now been reported for European descendants (1–8), Mexican Americans (9,10), Pima Indians (11–13), and African Americans (14). Evidence of linkage has emerged for several chromosomal locations (15,16), with at least six regions significantly linked to an obesity phenotype (maximum LOD score [MLS] > 3.3) and replicated in at least two studies based on the recent Human Obesity Gene Map (16). Despite these positive results, the literature contains inconsistent and contradictory reports. Consistent replication is the logical escape from this dilemma, although this is difficult to achieve as linkage is generally underpowered, and even quantitative trait loci (QTL) with strong effect are not expected to generate significant evidence of linkage consistently (17).

We previously reported suggestive evidence that BMI was linked to D3S2427 on chromosome 3q27 (map location = 187 cM, LOD score = 1.8) in an African-American sample (14). At the time, our findings provided replication

of linkage at this location for several obesity phenotypes as reported by Kissebah et al. (1). Linkage with early-onset diabetes was also found on chromosome 3q27 (18). Our earlier report was the result of a genome-wide scan study for obesity and hypertension as part of the Family Blood Pressure Program (19). To confirm our findings at the 3q27 region, we conducted a linkage analysis in a separate African-American sample of families and again by combining the new sample with that used in our original report on 3q27.

## RESEARCH DESIGN AND METHODS

**Recruitment of participants.** Two samples of African Americans were included in this analysis. The total sample is composed of 1,163 individuals from 330 families. Sample 1 included 618 individuals from 202 families, as reported previously (14). The second independent sample (sample 2) included 545 individuals from 128 families. While sample 1 was recruited based on a proband between the ages of 25 and 45 years and a blood pressure in the upper 25th percentile of the age-sex-specific distribution for this community, sample 2 was recruited with only the age restriction for the proband. After identification of the proband, all available nuclear family members were asked to participate. No restrictions were imposed in terms of weight or diet history in either sample. Our previous genome-wide scan for obesity was performed in sample 1 (14).

A clinical examination was performed that included medical history, anthropometric measurements, and blood collection. Weight was measured in light clothing on a calibrated electronic scale, and height was measured using a stadiometer.

**Genotyping.** The genotyping procedure for sample 1 was described in detail elsewhere (14). Briefly, 387 microsatellite markers (average 1 marker/10 cM) were genotyped at the National Heart, Lung, and Blood Institute (NHLBI) Mammalian Genotyping Service, Marshfield, Wisconsin, in sample 1. To further establish the evidence for linkage at 3q27 region from sample 1, we identified and genotyped 15 microsatellite markers in sample 2 in a 112-cM region around the linkage peak at this region. The mean heterozygosity for these 15 markers was 0.78 with an average sex-equal distance of 7.5 cM between two flanking markers. Of these 15 markers, nine markers were located on a 43-cM region around the region showing the strongest evidence of linkage in our original genome scan analysis, with an average distance of 4.8 cM (14). There were a total of 21 markers in the combined samples; 8 markers were genotyped in both samples. Because the samples were genotyped in two different labs, systematic allele shifting was adjusted for these markers according to the allele frequency distributions in the samples. We further calculated the identity-by-descent (IBD) sharing in the two samples separately without adjusting for the allele shifting. This IBD matrix was then used in the linkage analysis for comparison to the original results. In addition, we tested for linkage after setting the six markers with shifted alleles to missing in both populations. RELCHECK (20) was used to check the biological relationships within the pedigrees based on the available markers; it classified the biological relationship using the likelihood method. Because of the limited number of markers, the rates of misclassification could be high and large numbers of true sibpairs could be discarded (21). Therefore, the reported family relationships were conserved unless extreme statistical significance was identified. PEDCHECK (22) was used to check for Mendelian inconsistency. Errors identified by PEDCHECK that were not based on misclassification of pedigrees were

From the <sup>1</sup>Department of Preventive Medicine and Epidemiology, Loyola University Stritch School of Medicine, Maywood, Illinois; and the <sup>2</sup>Department of Pathology, Loyola University Stritch School of Medicine, Maywood, Illinois.

Address correspondence and reprint requests to Amy Luke, Department of Preventive Medicine & Epidemiology, Loyola University Stritch School of Medicine, 2160 South First Ave., Maywood, IL 60153. E-mail: aluke@lumc.edu.

Received for publication 13 December 2002 and accepted in revised form 29 January 2003.

IBD, identity by descent; LOD, logarithm of odds; MLS, maximum LOD score; QTL, quantitative trait loci.

© 2003 by the American Diabetes Association.

TABLE 1  
Descriptive characteristics of participants

	Sample 1		Sample 2		Combined sample	
	Men	Women	Men	Women	Men	Women
Individuals ( <i>n</i> )	250	368	193	352	443	720
Families ( <i>n</i> )	201		128		329	
Sibpairs ( <i>n</i> )	311		271		582	
Half-sibpairs ( <i>n</i> )	212		84		296	
Age (years)	38.6 ± 10.1	41.8 ± 12.4	35.3 ± 14.8	36.7 ± 5.1	37.1 ± 12.5	39.2 ± 14.0
Height (cm)	176.6 ± 7.2	164.0 ± 7.2	175.9 ± 9.0	164.3 ± 7.1	176.3 ± 8.0	164.1 ± 7.2
Weight (kg)	84.4 ± 21.4	87.2 ± 24.0	82.7 ± 21.7	83.9 ± 23.2	83.7 ± 21.5	85.6 ± 23.7
BMI (kg/m <sup>2</sup> )	27.0 ± 6.6	32.5 ± 8.8	26.5 ± 6.0	31.0 ± 8.3	26.8 ± 6.3	31.7 ± 8.5
BMI ≥30 (%)	25.2	54.0	24.6	48.3	24.9	51.2

Data are means ± SD unless otherwise indicated.

assumed to have occurred in the genotyping process, and the associated markers were set to missing among the appropriate family members.

**Statistical analysis.** Linkage analysis of BMI was performed using the multipoint variance component method implemented in GENEHUNTER2 (23). The variance component method specifies the expected genetic covariances between relatives as a function of their IBD relationships at a marker locus. The IBD probabilities were estimated from all available genotyped marker loci. The likelihood ratio test was applied to test the null hypothesis of no additive genetic variance due to a QTL. Sex, age, and age squared were incorporated as covariates, and their effects were simultaneously estimated by the maximum-likelihood method. Allele frequencies were estimated from the marker data using all of the individuals. Marshfield map distances were used in the linkage analysis. Linkage analysis was performed first in sample 2 and also repeated in sample 1. We then performed linkage analysis using the combined data of the two samples. To make the allele calling consistent in the two samples, we checked the allele frequency distribution for both samples and adjusted any allele shifting before combining them for further analyses.

To assess the region-wide empirical significance of our result, we performed a simulation study in the combined sample. We retained the pedigree and phenotype data and simulated the marker genotype data based on the observed marker allele frequencies. One thousand replications were generated and analyzed by GENEHUNTER2.

## RESULTS

The descriptive characteristics of the participants are presented in Table 1. Women were on average slightly older than men and more likely to be obese in both samples. The participants in sample 1 were somewhat older and heavier than those in sample 2. The heritability for BMI was 53% in the combined sample after adjusting for sex, age, and age squared calculated by GENEHUNTER2.

The multipoint linkage results for BMI from GENEHUNTER2 in sample 1, sample 2, and the combined sample at 3q27 region are presented in the figure. The MLS = 2.4 for new sample 2 was found at 188.29 cM, at marker D3S2427, with 1-LOD drop interval from 180 to 196 cM. The maximum LOD score (MLS = 2.0) was found at 187.51 cM, between marker D3S3676 and marker D3S2427 in sample 1. The 1-LOD drop interval was from 172 to 192 cM. The MLS is little higher than the original result (MLS = 1.8) (14) in sample 1 because more genotype cleaning was performed and a new version of GENEHUNTER software was used in this analysis. Theoretical analysis and a simulation study show that a 1-LOD-drop support interval corresponds roughly to a 90% confidence region in the case of a dense map of markers and provides even greater probability than a support region of coverage for sparser maps (24). In the combined sample, the MLS = 4.3 was found at 187.65 cM between marker D3S3676 and D3S2427 (unadjusted point-wise  $P = 4.3 \times 10^{-6}$ ), with 1-LOD drop

interval from 185 to 192 cM. The distance between MLS peaks between samples 1 and 2 was <1 cM, and most of the 1-LOD drop intervals overlapped the other. The MLS for the combined sample was close to the sum of the MLS for the two samples at the same location. When we used the IBD matrix calculated in the two samples separately without adjustment for allele shifting, the results were the same. In addition, there was only a modest effect on the results of linkage analysis after setting the six markers with shifted alleles to missing: the MLS was decreased from 4.3 to 4.1 at 187.65 cM. The variance accounted for by this QTL was estimated at ~40%. We recognize, however, that the strength of the QTL-specific effect is almost certainly overestimated by this method of analysis (25).

The region-wide empirical *P* value was calculated based on 1,000 replicate data sets for the combined sample. We found only two replicates having MLS >4.3 across the 100-cM region, which gave a region-wide empirical *P* value of 0.002.

## DISCUSSION

In a combined analysis of two samples of African-American families, we found a highly significant result for linkage with BMI at a region on 3q27 (185–192 cM, MLS = 4.3). In each of the two samples separately, suggestive linkages (MLS = 2.0 and MLS = 2.4) were observed at the same region. Encouragingly, at least another four studies have reported significant linkage at the same region.




FIG. 1. Multipoint linkage results for BMI at the chromosome 3 region (112.42–224.8 cM) in sample 1 (---), sample 2 (—), and the combined sample (—).

Kissebah et al. (1) reported strong evidence for linkage for several obesity-related phenotypes (LOD from 2.4 to 3.5) at the same region (182–200 cM) in 2,209 individuals from 507 nuclear Caucasian families. Vionnet et al. (18) also report significant linkage with diabetes or glucose intolerance in French persons aged 45 years and older. Not only have we previously reported suggestive linkage on 3q27 in sample 1 (14), we also reported the results of a combined analysis of genome scans for obesity using the interim data from the Family Blood Pressure Program (26). Using data from 6,849 individuals from four ethnic groups (European American, African American, Mexican American, and Asian), we found strong linkage evidence for a QTL at 3q27 for BMI (marker D3S2427, LOD = 3.4) (26). This large study further demonstrates that the linkage evidence on 3q27 is unlikely to be a false positive.

Several potential candidate genes have been identified around this region, including adiponectin (APM1), which is ~10 cM away from our peak, glucose transporters (GLUT2), and apolipoprotein D (Apo-D) (15). Among these genes, adiponectin gene (3q27) seems the most interesting. APM1 is the most abundant gene transcript specific to adipose tissue (27). Although the precise function of the adiponectin in energy expenditure and fat partitioning remains unclear, it is known that its circulating level is inversely correlated with BMI (28) and its mRNA level is suppressed in adipose tissue of obese animals and humans (29). Several studies have recently shown that polymorphisms in APM1 are associated with type 2 diabetes and other features of the insulin resistance syndrome (30–33). However, it is still not clear that the linkage peak identified in this region is due to the effect of APM1, and further studies should be warranted.

#### ACKNOWLEDGMENTS

This research was supported by grants from the National Heart, Lung, and Blood Institute (HL45508 and HL53353). Some of the genotyping was conducted by the NHLBI Mammalian Genotyping Service, Marshfield, Wisconsin.

#### REFERENCES

- Kissebah AH, Sonnenberg GE, Myklebust J, Goldstein M, Broman K, James RG, Marks JA, Krakower GR, Jacob HJ, Weber J, Martin L, Blangero J, Comuzzie AG: Quantitative trait loci on chromosomes 3 and 17 influence phenotypes of the metabolic syndrome. *Proc Natl Acad Sci* 97:14478–14483, 2000
- Hager J, Dina C, Franck S, Dubois S, Houari M, Vatin V, Vaillant E, Lorentz N, Basdevant A, Clement K, Guy-Grand B, Froguel P: A genome-wide scan for human obesity genes reveals a major susceptibility locus on chromosome 10. *Nat Genet* 20:304–308, 1998
- Lee JH, Reed DR, Li WD, Xu W, Joo EJ, Kilker RL, Nanthakumar E, North M, Sakul H, Bell C, Price RA: Genome scan for obesity and linkage to markers in 20q13. *Am J Hum Genet* 64:196–209, 1999
- Van der Kallen CJH, Cantor RM, van Greevenbroek MMJ, Geurts JMW, Bouwman FG, Aouixerat BE, Allayee H, Buurman WA, Lusi AJ, Rotter JJ, de Bruin TWA: Genome scan for adiposity in Dutch dyslipidemic families reveals novel quantitative trait loci for leptin, body mass index and soluble tumor necrosis factor receptor super family 1A. *Int J Obes Rel Metab Disord* 24:1381–1391, 2000
- Watanabe RM, Ghosh S, Langefeld CD, Valle TT, Hauser ER, Magnuson VL, Mohlke KL, Silander K, Ally DS, Chines P, Blaschak-Harvan J, Douglas JA, Duren WL, Epstein MP, Fingerlin TE, Kaleta HS, Lange EM, Li C, McEachin RC, Stringham HM, Trager E, White PP, Balow Jr J, Birznieks G, Chang J, Eldridge W: The Finland-United States Investigation of Non-Insulin-Dependent Diabetes Mellitus Genetics (FUSION) Study. II. An autosomal genome scan for diabetes-related quantitative trait loci. *Am J Hum Genet* 67:1186–1200, 2000
- Chagnon YC, Perusse L, Weisnagel SJ, Rankinen T, Bouchard C: The human obesity gene map: the 1999 update. *Obes Res* 8:89–117, 2000
- Ohman M, Oksanen L, Kaprio J, Koskenvuo M, Mustajoki P, Rissanen A, Salmi J, Kontula K, Peltonen L: Genome-wide scan of obesity in Finnish sibpairs reveals linkage to chromosome Xq24. *J Clin Endocrinol Metab* 85:3183–3190, 2000
- Perola M, Ohman M, Hiekkalinna T, Leppavuori J, Pajukanta P, Wessman M, Koskenvuo M, Palotie A, Lange K, Kaprio J, Peltonen L: Quantitative trait locus analysis of body-mass index and of stature, by combined analysis of genome scans of five Finnish study groups. *Am J Hum Genet* 69:117–123, 2001
- Comuzzie AG, Hixson JE, Almasy L, Mitchell BD, Mahaney MC, Dyer TD, Stern MP, MacCluer JW, Blangero J: A major quantitative trait locus determining serum leptin levels and fat mass is located on human chromosome 2. *Nat Genet* 15:273–276, 1997
- Mitchell BD, Cole SA, Comuzzie AG, Almasy L, Blangero J, MacCluer JW, Hixson JE: A quantitative trait locus influencing BMI maps to the region of the  $\beta$ -3 adrenergic receptor. *Diabetes* 48:1863–1867, 1999
- Norman RA, Tataranni PA, Pratley R, Thompson DB, Hanson RL, Prochazka M, Baier L, Ehm MG, Sakul H, Foroud T, Garvey WT, Burns D, Knowler WC, Bennett PH, Bogardus C, Ravussin E: Autosomal genome scan for loci linked to obesity and energy metabolism in Pima Indians. *Am J Hum Genet* 62:659–668, 1998
- Hanson RL, Ehm MG, Pettitt DJ, Prochazka M, Thompson DB, Timberlake D, Foroud T, Kobes S, Baier L, Burns D, Almasy L, Blangero J, Garvey WT, Bennett PH, Knowler WC: An autosomal genomic scan for loci linked to type II diabetes mellitus and body mass index in Pima Indians. *Am J Hum Genet* 63:1130–1138, 1998
- Walder K, Hanson RL, Kobes S, Knowler WC, Ravussin E: An autosomal genomic scan for loci linked to leptin concentrations in Pima Indians. *Int J Obes Rel Metab Disord* 200:24:559–565
- Zhu X, Cooper RS, Luke A, Chen G, Wu X, Kan D, Chakravarti A, Weder A: A genome-wide scan for obesity in African Americans. *Diabetes* 51:541–544, 2002
- Comuzzie AG, Williams JT, Martin LJ, Blangero J: Searching for genes underlying normal variation in human adiposity. *J Mol Med* 79:57–70, 2001
- Rankinen T, Perusse L, Weisnagel SJ, Snyder EE, Chagnon YC, Bouchard C: The human obesity gene map: the 2001 update. *Obes Res* 10:196–243, 2002
- Hirschhorn JN, Lindgren CM, Daly MJ, Kirby A, Schaffner SF, Burt NP, Altshuler D, Parker A, Rioux JD, Platko J, Gaudet D, Hudson TJ, Groop LC, Lander ES: Genomewide linkage analysis of stature in multiple populations reveals several regions with evidence of linkage to adult height. *Am J Hum Genet* 69:106–116, 2001
- Vionnet N, Hani EH, Dupont S, Gallina S, Francke S, Dotte S, Matos FD, Durand E, Lepretre F, Lecoeur C, Gallina P, Zekiri L, Dina C, Froguel P: Genomewide search for type 2 diabetes susceptibility genes in French whites: evidence for a novel susceptibility for early-onset diabetes on chromosome 3q27-pter and independent replication of a type 2 diabetes loci on chromosome 1q21q24. *Am J Hum Genet* 67:1470–1480, 2000
- Province MA, Boerwinkle E, Chakravarti A, Cooper R, Fornage M, Leppert M, Risch N, Ranade K: Lack of association of the angiotensinogen-6 polymorphism with blood pressure levels in the comprehensive NHLBI Family Blood Pressure Program. *J Hypertens* 18:867–875, 2000
- Broman KW, Weber JL: Estimation of pairwise relationships in the presence of genotyping errors. *Am J Hum Genet* 63:1563–1564, 1998
- Olson JM: Relationship estimation by Markov-process models in a sib-pair linkage study. *Am J Hum Genet* 64:1464–1472, 1999
- O'Connell JR, Weeks DE: PedCheck: A program for identification of genotype incompatibilities in linkage analysis. *Am J Hum Genet* 63:259–266, 1998
- Kruglyak L, Daly MJ, Reeve-Daly MP, Lander ES: Parametric and non-parametric linkage analysis: a unified multipoint approach. *Am J Hum Genet* 8:1347–1363, 1996
- Dupuis J, Siegmund D: Statistical methods for mapping quantitative trait loci from a dense set of markers. *Genetics* 151:373–386, 1999
- Goring HH, Terwilliger JD, Blangero J: Larger upward bias in estimation of locus-specific effects from genomewide scans. *Am J Hum Genet* 69:1357–1369, 2001
- Wu X, Cooper RS, Borecki I, Hanis C, Bray M, Lewis CE, Zhu X, Kan D, Luke A, Curb D: A combined analysis of genomewide linkage scans for body mass index from the National Heart, Lung, and Blood Institute Family Blood Pressure Program. *Am J Hum Genet* 70:1247–1256, 2002
- Maeda K, Okubo K, Shimomura I, Funahashi T, Matsuzawa Y, Matsubara K:

- cDNA cloning and expression of a novel adipose specific collagen-like factor apMI (adipose most abundant gene transcript 1). *Biochem Biophys Res Comm* 221:286–289, 1996
28. Comuzzie AG, Funahashi T, Sonnenberg G, Martin LJ, Jacob HJ, Kwitek Black AE, Maas D, Takahashi M, Kihara S, Tanaka S, Matsuzawa Y, Blangero J, Cohen D, Kissebah A: The genetic basis of plasma variation in adiponectin, a global endophenotype for obesity and the metabolic syndrome. *J Clin Endocrinol Metab* 86:4321–4325, 2001
  29. Matsubara M, Maruoka S, Katayose S: Inverse relationship between plasma adiponectin and leptin concentrations in normal-weight and obese women. *Eur J Endocrinol* 147:173–180, 2002
  30. Stumvoll M, Tschrirter O, Fritsche A, Staiger H, Renn W, Weisser M, Machicao F, Haring H: Association of the T-G polymorphism in adiponectin (exon 2) with obesity and insulin sensitivity: interaction with family history of type 2 diabetes. *Diabetes* 51:37–41, 2002
  31. Menzaghi C, Ercolino T, Di Paola R, Berg AH, Warram JH, Scherer PE, Trischitta V, Doria A: A haplotype at the adiponectin locus is associated with obesity and other features of the insulin resistance syndrome. *Diabetes* 51:2306–2312, 2002
  32. Kondo H, Shimomura I, Matsukawa Y, Kumada M, Takahashi M, Matsuda M, Ouchi N, Kihara S, Kawamoto T, Sumitsuji S, Funahashi T, Matsuzawa Y: Association of adiponectin mutation with type 2 diabetes: a candidate gene for the insulin resistance syndrome. *Diabetes* 51:2325–2328, 2002
  33. Kazuo H, Boutin P, Mori Y, Tobe K, Dina C, Yasuda K, Yamauchi T, Otabe S, Okada T, Eto K, Kadowaki H, Hagura R, Akanuma Y, Yazaki Y, Nagai R, Taniyama M, Matsubara K, Yoda M, Nakano Y, Kimura S, Tomita M, Ito C, Froguel P, Kadowaki T: Genetic variation in the gene encoding adiponectin is associated with an increased risk of type 2 diabetes in the Japanese population. *Diabetes* 51:536–540, 2002