

# Diacylglycerol Production and Protein Kinase C Activity Are Increased in a Mouse Model of Diabetic Embryopathy

Yuji Hiramatsu,<sup>1,2,4</sup> Naotaka Sekiguchi,<sup>2,3</sup> Michio Hayashi,<sup>2,3</sup> Keiji Isshiki,<sup>2,3</sup> Tamotsu Yokota,<sup>2,3</sup> George L. King,<sup>2,3</sup> and Mary R. Loeken<sup>1,3</sup>

**Activation of the diacylglycerol–protein kinase C (DAG-PKC) cascade by excess glucose has been implicated in vascular complications of diabetes. Its involvement in diabetic embryopathy has not been established. We examined DAG production and PKC activities in embryos and decidua of streptozotocin (STZ)-diabetic or transiently hyperglycemic mice during neural tube formation. STZ diabetes significantly increased DAG and total PKC activity in decidua (1.5- and 1.4-fold, respectively) and embryos (1.7- and 1.3-fold, respectively) on day 9.5. Membrane-associated PKC  $\alpha$ ,  $\beta$ II,  $\delta$ , and  $\zeta$  were increased in decidua by 1.25- to 2.8-fold. Maternal hyperglycemia induced by glucose injection on day 7.5, the day before the onset of neural tube formation, also increased DAG, PKC activity, and PKC isoforms (1.1-, 1.6-, and 1.5-fold, respectively) in the embryo on day 9.5. Notably, membrane-associated PKC activity was increased 24-fold in embryos of diabetic mice with structural defects. These data indicate that hyperglycemia just before organogenesis activates the DAG-PKC cascade and is correlated with congenital defects. *Diabetes* 51:2804–2810, 2002**

**S**ince the advent of insulin therapy, the survival of mothers with diabetes and their offspring has greatly improved. Most of the complications associated with diabetic pregnancy, such as spontaneous abortion, stillbirth, macrosomia, intrauterine growth restriction, and respiratory distress syndrome, can be significantly reduced by tight glycemic control before and throughout pregnancy (1,2). However, congenital malformations, which occur within the first 8 weeks of pregnancy, have been particularly difficult to avoid. The frequency of congenital malformations in the offspring of diabetic mothers is estimated to be 6–13%, a two- to fivefold increase over that in the nondiabetic population

From the <sup>1</sup>Section on Cellular and Molecular Physiology, Joslin Diabetes Center, Boston, Massachusetts; the <sup>2</sup>Section on Vascular Cell Biology, Joslin Diabetes Center, Boston, Massachusetts; the <sup>3</sup>Department of Medicine, Harvard Medical School, Boston, Massachusetts; and the <sup>4</sup>Department of Obstetrics and Gynecology, Okayama University Graduate School of Medicine and Dentistry, Okayama, Japan.

Address correspondence and reprint requests to Mary R. Loeken, Section on Cellular and Molecular Physiology, Joslin Diabetes Center, One Joslin Place, Boston, MA 02215. E-mail: mary.loeken@joslin.harvard.edu.

Received for publication 28 September 2001 and accepted in revised form 22 May 2002.

DAG, diacylglycerol; NTD, neural tube defect; PKC, protein kinase C; STZ, streptozotocin.

(3–7). The malformations associated with diabetic pregnancy affect many major organs, including the central nervous system, heart, skeleton, kidney, gastrointestinal system, and genitalia (5,6).

Studies using rodent models have attempted to provide biochemical and molecular explanations for diabetes-induced congenital defects. In vitro culture of rodent embryos has demonstrated that media containing elevated glucose or  $\beta$ -hydroxybutyrate can induce malformations like those caused by diabetic pregnancy (8,9). Several biochemical abnormalities have been observed in yolk sacs and embryos of diabetic rodents or after culture in high glucose, such as decreased uptake or membrane release of *myo*-inositol, arachidonic acid, and prostaglandin; increased lipid peroxidation; and decreased flux through the pentose shunt pathway (10–13). Moreover, the effects of antioxidants or overexpression of  $\text{Cu}^{2+}/\text{Zn}^{2+}$  superoxide dismutase to prevent diabetes-induced defects suggest that oxidative stress is an important factor in hyperglycemia-induced embryopathy (14–17).

Previously, we demonstrated using a mouse model of diabetic pregnancy that expression of *Pax-3*, a gene required for neural tube development, is inhibited on day 8.5 and that increased neural tube defects (NTDs) are observed on day 10.5 (18). This suggests that disturbed expression of genes that regulate morphogenetic processes may be central to diabetes-induced congenital defects. Antioxidants can prevent the increase in oxidative stress as well as decreased *Pax-3* expression and increased NTDs (Chang et al., submitted manuscript), supporting a role of oxidative stress in mediating the molecular and developmental abnormalities associated with diabetic pregnancy. Transient hyperglycemia caused by glucose injection only during day 7.5 is sufficient to inhibit *Pax-3* expression and to increase NTDs (19), indicating that there is a critical period before organogenesis in which the molecular program that leads to neural tube formation can be disturbed. This observation is consistent with earlier findings, in which withdrawal of insulin treatment from pregnant diabetic rats just before organogenesis increased diabetes-induced malformations (20).

Excess glucose activates the diacylglycerol–protein kinase C (DAG-PKC) cascade and is associated with several diabetic complications that affect retinal, renal, neural, and cardiovascular systems in patients with diabetes and diabetic animal models (21–29). PKC signaling is pivotal

for many cellular functions, including exocytosis, ion-exchange regulation, hormone and neurotransmitter release, and some processes that are particularly relevant to embryogenesis, such as cell proliferation, differentiation, and programmed cell death (21,30). Thus, if DAG-PKC signal transduction pathways are stimulated in embryonic tissues exposed to excess glucose, as they are in adult tissues affected by diabetic complications, then normal embryonic development may be disturbed. It is surprising that the ontogeny of DAG-PKC signaling during normal embryonic and fetal development has not been characterized extensively. In this report, we investigated whether DAG-PKC signaling was disturbed by diabetic pregnancy or transient hyperglycemia during organogenesis in the mouse embryo and whether this was associated with defective embryonic neural tube development.

## RESEARCH DESIGN AND METHODS

**Animals procedures.** Female ICR mice (Taconic, Germantown, NY) were obtained at 6–8 weeks of age, and diabetes was induced and treated as described previously (18). Briefly, mice received three daily injections of doses of 75 mg/kg streptozotocin (STZ; Sigma, St. Louis, MO). Tail-vein glucose concentrations were measured daily using a Glucometer Elite (Miles, Elhart, IN) starting at 1 week after STZ injection. Diabetes was controlled with subcutaneously implanted insulin pellets (Linshin, ON, Canada). Three weeks after implantation of insulin pellets, mice were mated along with age-matched nondiabetic mice. Noon on the day in which a copulation plug was found was determined to be day 0.5 of gestation. As reported previously (18), mice that had received STZ injections and been treated with insulin pellet, which had been euglycemic before pregnancy, became hyperglycemic on day 4.5 of pregnancy until the time they were sacrificed.

For testing the effects of transient hyperglycemia just before the beginning of organogenesis, nondiabetic mice were made hyperglycemic by subcutaneous injection of 1 ml of 25% glucose in PBS during an 8-h period on day 7.5 as described previously (19). Control mice received injections in parallel with saline. Blood glucose was assayed hourly, and additional injections were administered whenever blood glucose fell below 11.1 mmol/l (approximately every 1–2 h).

Diabetic mice or mice that received glucose injections were sacrificed by cervical dislocation on day 9.5 or 11.5 of gestation, uteri were excised, and embryos were recovered under a dissecting microscope in cold PBS. Extraembryonic membranes consisting of decidua, ectoplacental cones, and visceral yolk sac membranes (referred to collectively as decidua) were separated from embryos. Decidua and embryos were immediately frozen in liquid nitrogen, except for day 9.5 embryos, which were transferred to buffer as described below for in situ PKC assay. Staging of day 9.5 and 11.5 embryos was by comparison to Kaufman (31), and presence of externally detectable malformations in day 11.5 embryos was determined by an individual who was blinded to the treatment of the pregnancy. All procedures performed on mice were approved by the Institutional Animal Care and Use Committee of the Joslin Diabetes Center.

**DAG and PKC assays.** DAG and PKC were assayed as described previously (24,25,27) with the following modifications for whole embryo or decidua samples. Total DAG content of embryos and decidua or placentas were determined by radioenzymatic assay (Amersham, Arlington Heights, IL), using DAG kinase (Calbiochem, San Diego, CA). The values of total DAG content were normalized for protein by the Bradford method (32).

PKC activity in decidua and day 11.5 embryos was measured as previously described (20). Briefly, PKC activity in membrane and cytosolic fractions was measured by the ability to transfer  $^{32}\text{P}$  from ( $\gamma\text{-}^{32}\text{P}$ )-ATP (New England Nuclear, Boston, MA) into specific substrate octapeptide ( $\text{NH}_2\text{-RKRTLRLR-OH}$ ; Bio Synthesis, Lewisville, TX) in the presence of  $\text{Ca}^{2+}$ , phosphatidylserine, and DAG. PKC activity was calculated by subtracting the nonspecific kinase activity obtained in the absence of  $\text{Ca}^{2+}$ , phosphatidylserine, and DAG.

Because of the small size of day 9.5 embryos, from which sufficient amounts of cytosol and membrane to assay PKC activity could not be recovered, the sensitive in situ assay was performed as described previously (32,33) with modifications for whole embryos. Briefly, after dissection, embryos were transferred individually into a microfuge tube containing a buffered salt solution (137 mmol/l NaCl, 5.4 mmol/l KCl, 10 mmol/l magnesium chloride, 0.3 mmol/l sodium phosphate, 0.4 mmol/l potassium phosphate, 25 mmol/l  $\beta$ -glycerophosphate, 5.5 mmol/l D-glucose, 5 mmol/l EGTA, 1 mmol/l

TABLE 1  
Effects of diabetes on maternal blood glucose (mmol/l)

Gestational day	Nondiabetic	Diabetic
0.5	5.95 $\pm$ 0.79 (25)	5.36 $\pm$ 1.69 (26)
9.5	7.16 $\pm$ 0.39 (25)	23.61 $\pm$ 3.85* (15)

Data are means  $\pm$  SD (*n*). \**P* < 0.0001 vs. control.

$\text{CaCl}_2$ , and 20 mmol/l HEPES [pH 7.2]). The reaction was started by adding a mixture of 100  $\mu\text{mol/l}$  ( $\gamma\text{-}^{32}\text{P}$ )-ATP, 50  $\mu\text{g/ml}$  digitonin, and 100  $\mu\text{mol/l}$  octapeptide. The basal activity was calculated by adding the mixture without octapeptide. The kinase reaction was terminated after 15 min at 25°C by adding 40  $\mu\text{l}$  of 25% (wt/vol) TCA. Fifty microliters of the acidified reaction mixture was spotted onto phosphocellulose filters (Whatman P-21) and washed three times with 75 mmol/l sodium phosphate (pH 7.5). The PKC-dependent phosphorylation of the peptide substrate bound to the filter was quantified by scintillation counting with a LC6500 multipurpose scintillation counter (Beckman). Embryos were solubilized in 0.1% SDS and 1 N of NaOH and neutralized with 1 N of HCl for protein quantitation by the Bradford method.

**Immunoblot analysis.** A total of 30–60  $\mu\text{g}$  of protein extracts from whole embryos or membrane and cytosolic fractions of decidua were separated by SDS/PAGE (8% PAGE) using a XCell II system (Novex, San Diego, CA) and transferred to nitrocellulose membranes. After transfer, membranes were blocked and then incubated with rabbit polyclonal antibodies raised against cPKC- $\alpha$ , PKC- $\beta$  II, nPKC- $\delta$ , or nPKC- $\zeta$  (Santa Cruz Biotechnology, Santa Cruz, CA). The blots were washed and then incubated with horseradish peroxidase-coupled donkey anti-rabbit IgG (Amersham Pharmacia Biotech, Piscataway, NJ). Bands were detected by ECL Western blotting detection reagents (Amersham Pharmacia Biotech) and were quantified by densitometric scanning of the X-ray film and analysis by Image Quant software.


**Statistical analysis.** Statistical analyses were performed using Student's unpaired *t* test. Analyses were performed using the Statview statistical package for Macintosh (Hulinks, Cary, NC). *P* < 0.05 was defined as statistically significant.

## RESULTS

**Maternal diabetes increases DAG production and PKC activity in embryos and decidua on day 9.5.** In preliminary studies, a profile for DAG production and PKC activity during gestation was performed using decidua and placenta to determine whether DAG-PKC signaling was detectable during prenatal development. Decidua or placentas were obtained from normal, nondiabetic pregnant mice beginning on day 9.5 and every 3 days until day 18.5, the day before parturition. DAG was detectable (200  $\pm$  86 pmol/mg wet wt) on day 9.5 and was increased by 1.7-fold on day 18.5 (data not shown). Similarly, membrane-associated PKC activity was detectable (25.6  $\pm$  4.5 pmol  $\cdot$  mg protein $^{-1}$   $\cdot$  min $^{-1}$ ) on day 9.5 and increased twofold by day 15.5 (data not shown). Immunoblot analysis demonstrated that PKC  $\alpha$ ,  $\beta$ II,  $\delta$ , and  $\zeta$  were expressed at these time points and increased on days 15.5 and 18.5.

Because DAG and PKC were detectable during early organogenesis (day 9.5) in decidua, the effects of maternal diabetes on DAG and PKC in the embryo and decidua were investigated. Similar to previous findings (18), diabetic mice were euglycemic on day 0.5 of pregnancy but were significantly hyperglycemic on day 9.5 (Table 1). Mice were sacrificed on day 9.5, and DAG and PKC were assayed in embryos and decidua.


DAG concentrations and PKC activity were significantly increased (1.7- and 1.3-fold, respectively) in embryos of diabetic mice (Fig. 1A and B). Similarly, decidual DAG was increased 1.5-fold, and PKC activity in membrane and cytosolic fractions was significantly increased (1.2- and



**FIG. 1.** DAG and PKC in day 9.5 embryos and decidua of nondiabetic (control) or diabetic (DM) mice. *A*: DAG concentrations in embryos; \* $P < 0.001$  vs. control. *B*: PKC activity in embryos measured by in situ assay; \* $P < 0.05$  vs. control. *C*: DAG concentrations in decidua; \* $P < 0.05$  vs. control. *D*: PKC activity of decidua measured in membrane and cytosolic fractions; \* $P < 0.05$  vs. control. □, control (nondiabetic) results; ■, diabetic results.

1.3-fold, respectively) in samples from diabetic mice (Fig. 1C and D). Immunoblot analysis demonstrated that steady-state concentrations of PKC isoforms  $\alpha$ ,  $\beta$ II,  $\delta$ , and  $\zeta$  in whole embryo were not significantly affected by maternal diabetes (Fig. 2A), suggesting that increased PKC activity in cell extracts (Fig. 1B) resulted from activation of the enzymes, not from new mRNA or protein synthesis. In decidua, membrane association of all PKC isoforms examined (PKC  $\alpha$ ,  $\beta$ II,  $\delta$ ,  $\zeta$ ) were increased in samples from diabetic pregnancies but not in the cytosolic fraction (Fig. 2B). This suggests that diabetes activated PKC located in the cytoplasmic compartment, causing it to be translocated to the membranous compartment.

**Transient hyperglycemia before organogenesis activates DAG and PKC during organogenesis.** We next investigated whether transient hyperglycemia occurring only on day 7.5 is sufficient to increase DAG production and PKC activity on day 9.5. As shown in Table 2, injection of glucose at approximately hourly intervals during an 8-h time period on day 7.5 significantly increased maternal blood glucose, and blood glucose returned to normal by day 9.5. Nevertheless, even though maternal glucose concentrations were normal on day 9.5, DAG concentrations and PKC activity were significantly increased in embryos of mice that received glucose injections (Fig. 3A and B). Similarly, PKC activity in membrane fractions of decidua and decidua of mice that received glucose injections was significantly increased, and decidual DAG was also increased, although the increase in DAG was not statistically significant (Fig. 3C and D). Western blot analysis demonstrated that the steady-state concentrations of PKC isozymes in whole embryos or in membrane or cytosolic fractions of decidua were variable but were not significantly increased by glucose injection (Fig. 4A and B). This suggests that, like maternal diabetes, effects of hyperglycemia on PKC activity were due to stimulation of enzymatic activity rather than to de novo mRNA or protein synthesis.



**FIG. 2.** Immunoblot analysis of PKC isoforms  $\alpha$ ,  $\beta$ II,  $\delta$ , and  $\zeta$  from day 9.5 nondiabetic or diabetic pregnancies. *A*: Representative immunoblot of two embryos of nondiabetic control mice and two embryos of diabetic mice. Quantitation of bands in the upper panel is shown in the lower panel, with band intensity from embryos of diabetic mice expressed as a percentage of bands from embryos of nondiabetic mice. *B*: Representative immunoblot of two decidua from nondiabetic mice and two from diabetic mice. Quantitation of bands from six replicate decidua from nondiabetic pregnancies and six replicate decidua from diabetic pregnancies are shown in the lower panel. Quantitation of membrane bands are shown in solid bars and of cytosol bands are shown in shaded bars. \* $P < 0.05$ ; \*\* $P < 0.002$  vs. nondiabetic control.

**Increased PKC in defective embryos of diabetic mice.** By day 10.5, formation of the neural tube is complete and embryos with defective neural tubes can be distinguished from normal embryos. Examples of a normal embryo and an embryo with an NTD caused by maternal diabetes are shown in Fig. 5.

To test whether PKC activity was affected in malformed embryos of diabetic mice, we obtained embryos on day 11.5 of gestation from diabetic and nondiabetic pregnancies. As on day 9.5, diabetic mice were significantly more hyperglycemic than nondiabetic mice ( $25.53 \pm 2.33$  and  $7.58 \pm 0.19$  mmol/L, respectively;  $P < 0.0005$ ). Like day 10.5 embryos, normal and defective day 11.5 embryos can be

TABLE 2

Effects of glucose injection on day 7.5 on maternal blood glucose (mmol/l)

Gestational day	Saline injection	Glucose injection
0.5	5.97 ± 0.47 (12)	
7.5	7.23 ± 0.44 (12)	12.47 ± 2.02* (31)
9.5	7.17 ± 0.46 (12)	7.48 ± 0.78 (22)

Data are means ± SD (*n*). \**P* < 0.0001 vs. control.

distinguished from each other, but they are significantly larger so that PKC in membrane and cytosol fractions can readily be assayed. All of the embryos of nondiabetic mice collected for this assay were normal. In contrast, 25% of the embryos of diabetic mice had defects affecting the neural tube (exencephaly, cerebral hemorrhage, midbrain/hindbrain underdevelopment), sometimes also including heart or gut defects.

As shown in Fig. 6, membrane-associated PKC activity in defective embryos of diabetic mice was increased 24-fold compared with embryos of nondiabetic mice and threefold compared with morphologically normal embryos from diabetic pregnancies. Although mean PKC activity was higher in normal embryos of diabetic mice compared with embryos from nondiabetic pregnancies, the increase was not statistically significant. There was no significant difference in PKC activity of cytosol fractions from any treatment group. In decidua, there was a slight increase in membrane and cytosol PKC activity in both normal and abnormal embryos of diabetic mice, although these increases were not significant.

## DISCUSSION

The PKC family of serine/threonine kinases plays a critical role in the regulation of cellular differentiation and proliferation (34). Therefore, dysregulation of this pathway would be expected to have adverse consequences on




FIG. 3. Effects of glucose injections on day 7.5 on DAG and PKC in embryos and decidua on day 9.5 assayed as in Fig. 2. **A**: DAG concentrations in embryos; \**P* < 0.05 vs. control. **B**: PKC activity in embryos; \**P* < 0.001 vs. control. **C**: DAG concentrations in decidua. **D**: PKC activity in cytosolic and membrane fractions of decidua; \**P* < 0.001 vs. control. □, control (saline injection) results; ■, glucose injection results.





FIG. 4. **A**: Representative immunoblot analysis of PKC isoforms in two day 9.5 embryos of mice that received saline injections and in three embryos of mice that received glucose injections. Quantitation of bands in the upper panel after scanning densitometry is shown in the lower panel with band intensity from embryos of diabetic mice expressed as a percentage of bands from embryos of nondiabetic mice. **B**: Representative immunoblot analysis of two day 9.5 decidua of mice that received saline injections and in two decidua of mice that received glucose injections. Quantitation of bands from four replicate decidua of saline-injected pregnancies and four glucose-injected pregnancies is shown in solid bars and of cytosol bands are shown in shaded bars.

embryonic development. Increases in DAG-PKC signaling have been noted in retina, aorta, heart, and glomeruli of patients with diabetes and diabetic animals and seem to be



**FIG. 5. A:** Normal day 10.5 embryo from a nondiabetic pregnancy. **B:** Embryo from a diabetic mouse displaying exencephaly of hindbrain and midbrain structures.

directly related to elevated glucose concentrations (22,24–29). Activation of the DAG-PKC pathway associated with abnormal growth and differentiation during diabetic embryopathy has not previously been demonstrated.


Several PKC isotypes have been detected in the placenta and decidua of the human and the rat (35–39). In the human placenta, there are significant increases in PKC  $\beta$ II and  $\epsilon$  on the microvillus membrane and in PKC  $\gamma$  and  $\epsilon$  on the basal membrane between 16 and 40 weeks of gestation (40). PKC is involved in the regulation of human chorionic gonadotropin, estrogen, and progesterone production by trophoblasts. Downregulation of PKC activity by chronic exposure to the phorbol ester 12-myristate 13-acetate or 12,13-dibutyrate stimulates secretion of human chorionic gonadotropin  $\beta$  (37,39). Similarly, a general PKC inhibitor, isoquinolinesulfonamide (H7) attenuates DNA synthesis in proliferating and differentiating trophoblast cells and accelerates the acquisition of progesterone biosynthetic capabilities (38). In the experiments reported here, we investigated the activities and protein concentrations of PKC  $\alpha$ ,  $\beta$ II,  $\delta$ , and  $\zeta$  in embryos and extraembryonic tissues during early organogenesis. We selected these PKC isoforms for study because of their involvement in other diabetic complications (41–43). Notably, DAG production and PKC activity were increased in the extraembryonic structures (decidua, ectoplacental cone, and visceral yolk sac) by diabetes or hyperglycemia. Because these membranes contribute to implantation, development of the placenta, hormone and growth factor production, and nutrient and gas exchange, increased DAG-PKC signaling

could interfere with any of these processes. Thus, increased PKC signaling may be of consequence to the embryo during organogenesis as well as impair later placental function, which could disturb fetal development.

In the embryo, inhibition of PKC activity in rat embryos causes NTDs, indicating that PKC activity is required for normal neural tube development (44,45). However, our results suggest that excess PKC signaling is also associated with defective development. It should be noted that increased DAG-PKC activity occurred on day 9.5, while the embryo is undergoing dramatic morphogenesis, particularly the establishment of the neural tube and the heart. There are three obvious interpretations of the correlation between increased DAG-PKC in embryos of diabetic dams and defective organogenesis: 1) that altered PKC signaling is responsible for abnormal morphogenesis, 2) that DAG-PKC signaling is increased in embryos of diabetic dams as a consequence of an already disturbed developmental program, or 3) that sustained increased DAG-PKC signaling disturbs the formation of structures that form at successively later times in gestation. Because the increase in DAG and PKC on day 9.5 occurs after the onset of gene expression that induces neural tube formation (day 8.5), this suggests that the increase in DAG and PKC on day 9.5 could not be a cause but could be a consequence of defective development of the neural tube. The increase in PKC activity in embryos of diabetic mice, which was even greater in defective embryos than in embryos that were morphologically normal, supports this interpretation. However, there are many structures that continue to form throughout the embryonic period and that depend on inductive influences from previously formed structures. Therefore, increased PKC activity in defective day 11.5 embryos could be due to abnormal signaling pathways in structures undergoing morphogenesis at that stage of development and that are adversely affected by previously existing structural defects. Because we did not cellularly localize the sites of increased PKC activity, it is not possible to distinguish these possibilities at this time. Future research will be necessary to determine the relationship between abnormal DAG-PKC signaling and defective development.

Another interpretation is that increased DAG-PKC signaling originates from vascular stem cells rather than from neuroepithelial or other primordial cell types. Because neovascularization is important for establishment of early organ structures, diabetic embryopathy secondary to defective neovascularization may be a novel type of diabetic vascular complication. In this regard, it should be noted that vitamin E prevents diabetes-induced vascular complications by inhibiting glucose activation of DAG-PKC signaling (46–48). Several studies have demonstrated that antioxidants can prevent diabetic embryopathy in animal models (14–17), and we have shown that hyperglycemia-induced oxidative stress inhibits expression of *Pax-3* (Chang et al., submitted manuscript). Thus, it will be important in the future to determine whether antioxidants normalize DAG and PKC in the embryo and decidua, as it does in retinal and renal tissues (48,49).

It is interesting that induction of hyperglycemia during day 7.5 alone was sufficient to cause a sustained activation of this pathway that could still be measured on day 9.5.



**FIG. 6. PKC activity in membrane and cytosolic fractions of day 11.5 embryos and decidua from nondiabetic (control) mice or in normal or abnormal embryos of diabetic mice. □, results from control (nondiabetic) embryos; ■, results from diabetic normal (without malformations) embryos; ▨, results from diabetic abnormal (with structural defects) embryos; n = 3 replicate samples of pooled embryos; \*P < 0.01 vs. control and diabetic, normal.**

This indicates that, just as there is a sensitive period of development before organogenesis in which withholding insulin treatment from diabetic rats will cause NTDs or inducing hyperglycemia will disturb gene expression and cause NTDs (19,20), this same period of development is susceptible to sustained hyperglycemia-induced DAG production and PKC signaling. Because there was no consistent increase in the steady-state amount of any PKC isozyme by immunoblot analysis, this suggests that there is a stable activation of PKC activity, most likely due to increased production of DAG rather than to increased expression of PKC mRNA or protein. For example, maternal hyperglycemia may induce expression of genes that encode the regulatory enzymes in this pathway, such as phospholipase C. A sustained effect of hyperglycemia on DAG and PKC is consistent with our previous observations that DAG and PKC remain elevated even years after the onset of diabetes in patients, but the increase in activity may take several days of exposure to hyperglycemia to appear (24). Indeed, increased activity of PKC, rather than increased synthesis, may explain the failure of others to see an increase in mRNA for PKC  $\alpha$ ,  $\beta$ , or  $\gamma$  in rat embryos of diabetic mothers or cultured in high glucose (45).

This is the first report that PKC activity is increased by experimental procedures (maternal diabetes or transient hyperglycemia) that cause diabetic embryopathy and, furthermore, that PKC activity is even more increased in defective embryos than in normal embryos of diabetic mice after completion of neural tube fusion. Certainly, further investigation will be necessary to localize cellularly the sites of increased PKC activity, both in embryonic and in extraembryonic tissues, at critical stages during organogenesis and to determine whether increased PKC activity contributes to abnormal development.

#### ACKNOWLEDGMENTS

This research was supported by grants R01-DK52865 to M.R.L. and R01-EY5110 to G.K.

We are grateful to Melissa Horal and Rakhi Patel for valuable technical assistance.

#### REFERENCES

- Cunningham FG, MacDonald PC, Gant NF, Leveno KJ, Gilstrap LCI, Hankins GDV, Clark SL: Diabetes. In *Williams Obstetrics*. 20th ed. London, Prentice-Hall International, 1997, p. 1203-1221
- The Diabetes Control and Complications Trial Study Group: Pregnancy outcomes in the Diabetes Control and Complications Trial. *Am J Obstet Gynecol* 174:1343-1353, 1996
- Mills JL, Baker L, Goldman AS: Malformations in infants of diabetic mothers occur before the seventh gestational week: implications for treatment. *Diabetes* 28:292-293, 1979
- Miller E, Hare JW, Cloherty JP, Dunn PJ, Gleason RE, Soeldner JS, Kitzmiller JL: Elevated maternal hemoglobin A1c in early pregnancy and major congenital anomalies in infants of diabetic mothers. *N Engl J Med* 304:1331-1334, 1981
- Miodovnik M, Mimouni F, Dignan PSJ, Berk MA, Ballard JL, Siddiqi TA, Khoury J, Tsang RC: Major malformations in infants of IDDM women: vasculopathy and early first-trimester poor glycemic control. *Diabetes Care* 11:713-718, 1988
- Becerra JE, Khoury MJ, Cordero JF, Erickson JD: Diabetes mellitus during pregnancy and the risks for the specific birth defects: a population-based case-control study. *Pediatrics* 85:1-9, 1990
- Hawthorne G, Robson S, Ryall EA, Sen D, Roberts SH, Ward Platt MP: Prospective population based survey of outcome of pregnancy in diabetic women: results of the Northern Diabetic Pregnancy Audit, 1994. *Br Med J* 315:279-281, 1997
- Cockroft DL, Coppola PT: Teratogenic effects of excess glucose on head-fold rat embryos in culture. *Teratology* 16:141-146, 1977
- Horton WE Jr, Sadler TW: Effects of maternal diabetes on early embryogenesis: alterations in morphogenesis produced by the ketone body, B-hydroxybutyrate. *Diabetes* 32:610-616, 1983
- Shum L, Sadler TW: Biochemical basis for D,L-beta-hydroxybutyrate-induced teratogenesis. *Teratology* 42:553-563, 1990
- Reece E, Homko C, Wu U, Wiznitzer A: The role of free radicals and membrane lipids in diabetes-induced congenital malformations. *J Soc Gynecol Investig* 5:178-187, 1998
- Wentzel P, Welsh N, Eriksson UJ: Developmental damage, increased lipid peroxidation, diminished cyclooxygenase-2 gene expression, and lowered prostaglandin E<sub>2</sub> levels in rat embryos exposed to a diabetic environment. *Diabetes* 48:813-820, 1999
- Engstrom E, Haglund A, Eriksson UJ: Effects of maternal diabetes or in vitro hyperglycemia on uptake of palmitic and arachidonic acid by rat embryos. *Pediatr Res* 30:150-153, 1991
- Siman CM, Eriksson UJ: Vitamin E decreases the occurrence of malformations in the offspring of diabetic rats. *Diabetes* 46:1054-1061, 1997
- Sivan E, Reece EA, Wu Y-K, Homko CJ, Polansky M, Borenstein M: Dietary vitamin E prophylaxis and diabetic embryopathy: morphologic and biochemical analysis. *Am J Obstet Gynecol* 175:793-799, 1996
- Viana M, Herrera E, Bonet B: Teratogenic effects of diabetes mellitus in the rat. Prevention by vitamin E. *Diabetologia* 39:1041-1046, 1996
- Wentzel P, Eriksson UJ: Antioxidants diminish developmental damage induced by high glucose and cyclooxygenase inhibitors in rat embryos in vitro. *Diabetes* 47:677-684, 1998
- Phelan SA, Ito M, Loeken MR: Neural tube defects in embryos of diabetic mice: role of the Pax-3 gene and apoptosis. *Diabetes* 46:1189-1197, 1997
- Fine E, Horal M, Chang T, Fortin G, Loeken M: Hyperglycemia is responsible for altered gene expression, apoptosis, and neural tube defects associated with diabetic pregnancy. *Diabetes* 48:2454-2462, 1999
- Eriksson UJ, Bone AJ, Turnbull DM, Baird JD: Timed interruption of insulin therapy in diabetic BB/E rat pregnancy: effect on maternal metabolism and fetal outcome. *Acta Endocrinol (Copenh)* 120:800-810, 1989
- Meire M, King GL: Protein kinase C. In *Diabetes Mellitus, A Fundamental and CLINICAL Text*. 2nd ed. LeRoith D, Taylor SI, Olefsky JM, Eds. Philadelphia, Lippincott Williams & Wilkins, 2000, p. 1016-1027
- King GL, Wakasaki H: Theoretical mechanisms by which hyperglycemia and insulin resistance could cause cardiovascular diseases in diabetes. *Diabetes Care* 22 (Suppl. 3):C31-C37, 1999
- Jack AM, Cameron NE, Cotter MA: Effects of the diacylglycerol complexing agent, cremophor, on nerve-conduction velocity and perfusion in diabetic rats. *J Diabetes Complications* 13:2-9, 1999
- Ishii H, Koya D, King GL: Protein kinase C activation and its role in the development of vascular complications in diabetes mellitus. *J Mol Med* 76:21-31, 1998
- Inoguchi T, Battan R, Handler E, Sportsman JR, Heath W, King GL: Preferential elevation of protein kinase C isoform beta II and diacylglycerol levels in the aorta and heart of diabetic rats: differential reversibility to glycemic control by islet cell transplantation. *Proc Natl Acad Sci U S A* 89:11059-11063, 1992
- Haller H, Baur E, Quass P, Behrend M, Lindschau C, Distler A, Luft FC: High glucose concentrations and protein kinase C isoforms in vascular smooth muscle cells. *Kidney Int* 47:1057-1067, 1995
- Xia P, Inoguchi T, Kern TS, Engerman RL, Oates PJ, King GL: Characterization of the mechanism for the chronic activation of diacylglycerol-protein kinase C pathway in diabetes and hypergalactosemia. *Diabetes* 43:1122-1129, 1994
- Wolf BA, Williamson JR, Easom RA, Chang K, Sherman WR, Turk J: Diacylglycerol accumulation and microvascular abnormalities induced by elevated glucose levels. *J Clin Invest* 87:31-38, 1991
- Shiba T, Inoguchi T, Sportsman JR, Heath WF, Bursell S, King GL: Correlation of diacylglycerol level and protein kinase C activity in rat retina to retinal circulation. *Am J Physiol* 265:E783-E793, 1993
- Buchner K: The role of protein kinase C in the regulation of cell growth and in signalling to the cell nucleus. *J Cancer Res Clin Oncol* 126:1-11, 2000
- Kaufman MH: *The Atlas of Mouse Development*. San Diego, CA, Academic Press, 1992
- Ishii H, Hirousek MR, Koya D, Takagi C, Xia P, Clermont A, Bursell SE, Kern TS, Ballas LM, Heath WF, Stramm LE, Feener EP, King GL: Amelioration of vascular dysfunctions in diabetic rats by an oral PKC beta inhibitor. *Science* 272:728-731, 1996
- Williams B, Schrier RW: Characterization of glucose-induced in situ protein kinase C activity in cultured vascular smooth muscle cells. *Diabetes* 41:1464-1472, 1992

34. Nishizuka Y: The molecular heterogeneity of protein kinase C and its implications for cellular regulation. *Nature* 334:661–665, 1988
35. Nomura S, Tokumitsu H, Mizutani S, Narita O, Tomada Y, Hidaka H: Identification of two subtypes of protein kinase C in human placenta. *Placenta* 12:605–613, 1991
36. Tertrin-Clary C, Chenut MC, de la Llosa P: Isolation of protein kinase C isoforms from human full-term placenta. *Ann Endocrinol* 52:327–330, 1991
37. Iwashita M, Watanabe M, Setoyama T, Mimuro T, Nakayama S, Adachi T, Takeda Y, Sakamoto S: Effects of diacylglycerol and gonadotropin-releasing hormone on human chorionic gonadotropin release by cultured trophoblast cells. *Placenta* 13:213–221, 1992
38. Yamamoto T, Chapman BM, Soares MJ: Protein kinase C dependent and independent mechanisms controlling rat trophoblast cell DNA synthesis and differentiation. *J Reprod Fertil* 111:15–20, 1997
39. Baker VL, Murai JT, Taylor RN: Downregulation of protein kinase C by phorbol ester increases expression of epidermal growth factor receptors in transformed trophoblasts and amplifies human chorionic gonadotropin production. *Placenta* 19:475–482, 1998
40. Ruzycy AL, Jansson T, Illsley N: Differential expression of protein kinase C isoforms in the human placenta. *Placenta* 17:461–469, 1996
41. Kang N, Alexander G, Park JK, Maasch C, Buchwalow I, Luft FC, Haller H: Differential expression of protein kinase C isoforms in streptozotocin-induced diabetic rats. *Kidney Int* 56:1737–1750, 1999
42. Liu X, Wang J, Takeda N, Binaglia L, Panagia V, Dhalla NS: Changes in cardiac protein kinase C activities and isozymes in streptozotocin-induced diabetes. *Am J Physiol* 277:E798–E804, 1999
43. Babazono T, Kapor-Drezgic J, Dlugosz JA, Whiteside C: Altered expression and subcellular localization of diacylglycerol-sensitive protein kinase C isoforms in diabetic rat glomerular cells. *Diabetes* 47:668–676, 1998
44. Ward KW, Rogers EH, Hunter ES 3rd: Dymorphogenic effects of a specific protein kinase C inhibitor during neurulation. *Reprod Toxicol* 12:525–534, 1998
45. Wentzel P, Wentzel CR, Gareskog MB, Eriksson UJ: Induction of embryonic dymorphogenesis by high glucose concentration, disturbed inositol metabolism, and inhibited protein kinase C activity. *Teratology* 63:193–201, 2001
46. Kunisaki M, Fumio U, Nawata H, King GL: Vitamin E normalizes diacylglycerol-protein kinase C activation induced by hyperglycemia in rat vascular tissues. *Diabetes* 45 (Suppl. 3):S117–S119, 1996
47. Keegan A, Walbank H, Cotter MA, Cameron NE: Chronic vitamin E treatment prevents defective endothelium-dependent relaxation in diabetic rat aorta. *Diabetologia* 38:1475–1478, 1995
48. Kunisaki M, Bursell S-E, Clermont AC, Ishii H, Ballas LM, Jirousek MR, Umeda F, Naata H, King GL: Vitamin E prevents diabetes-induced abnormal retinal blood flow via the diacylglycerol-protein kinase C pathway. *Am J Physiol (Endocrinol Metab)* 269:E239–E246, 1995
49. Koya D, Lee I-K, Ishii H, Kanoh H, King GL: Prevention of glomerular dysfunctions in diabetic rats by treatment of d-a-tocopherol. *J Am Soc Nephrol* 8:426–435, 1997