

Coordinated Regulation of Fat-Specific and Liver-Specific Glycerol Channels, Aquaporin Adipose and Aquaporin 9

Hiroshi Kuriyama, Iichiro Shimomura, Ken Kishida, Hidehiko Kondo, Naoki Furuyama, Hitoshi Nishizawa, Norikazu Maeda, Morihiro Matsuda, Hiroyuki Nagaretani, Shinji Kihara, Tadashi Nakamura, Yoshihiro Tochino, Tohru Funahashi, and Yuji Matsuzawa

Plasma glycerol is a major substrate for hepatic gluconeogenesis. Aquaporin adipose (AQPap/7), an adipose-specific glycerol channel, provides fat-derived glycerol into plasma. In the present study, we cloned the coding and promoter regions of mouse aquaporin 9 (AQP9), a liver-specific glycerol channel. Fasting and refeeding of mice increased and decreased hepatic AQP9 mRNA levels, respectively. Insulin deficiency induced by streptozotocin resulted in increased hepatic AQP9 mRNA. These changes in hepatic AQP9 mRNA were accompanied by those of hepatic gluconeogenic mRNAs and plasma glycerol levels. In cultured hepatocytes, insulin downregulated AQP9 mRNA. The AQP9 promoter contained the negative insulin response element TGTTTTC at $-496/-502$, similar to the promoter of the AQPap/7 gene. In contrast, in insulin-resistant *db+/db+* mice, AQPap/7 mRNA in fat and AQP9 mRNA in liver were increased, despite hyperinsulinemia, with high plasma glycerol and glucose levels. Glycerol infusion in the *db+/db+* mice augmented hepatic glucose output. Our results indicate that coordinated regulations of fat-specific AQPap/7 and liver-specific AQP9 should be crucial to determine glucose metabolism in physiology and insulin resistance. *Diabetes* 51: 2915–2921, 2002

The precise mechanism for the common association between obesity and diabetes remains unresolved. Adipose tissue supplies lipolysis-derived free fatty acid (FFA) and glycerol to other organs in various physiological conditions, such as diet restriction and physical exercise (1–4). We and others have reported that accumulation of intra-abdominal visceral fat is closely associated with increased incidence of metabolic complications in obesity (5–9). Higher influx of

FFA into the liver, especially from accumulated visceral fat, drives the hepatic synthesis of triglyceride, leading to fatty liver and hypertriglyceridemia (10–12).

Glycerol, another product of lipolysis, is an important substrate for hepatic gluconeogenesis (13–15). As the substrate for hepatic glucose production, glycerol accounts for 90% in the prolonged fasting state and ~50% in the postabsorptive state in rodents (16,17). In humans, it was estimated that 20% of the gluconeogenesis was mediated by glycerol after 60 h of fasting (13,18). Higher influx of glycerol into the liver from the accumulated fat may be relevant to the development of diabetes in obesity. The molecular basis for the secretion and uptake of glycerol has been characterized recently. Aquaporins (AQPs), which are channel-forming integral proteins, function as water channels (19). To date, at least 10 AQPs have been identified (20). Functional studies have distinguished the members of the AQP family into two subgroups: AQPs that are selective water channels and aquaglyceroporins that transport glycerol as well as water (21). We recently cloned a novel adipose-specific glycerol channel, designated aquaporin adipose (AQPap/7) (22), and reported that this protein functions as a glycerol channel from fat into blood (23–25). Recently, a liver-specific aquaglyceroporin was also identified and named AQP9 (26). The significance and regulation of AQP9 in physiological and pathophysiological conditions have not been elucidated.

In the current study, we demonstrate the inhibitory effect of insulin on the expression of the AQP9 gene in normal liver through the negative insulin response element (IRE) in its promoter region (27), similar to the regulation of AQPap/7 in adipose tissue (23,24). These coordinated regulations of AQPap/7 and AQP9 should account for the physiological suppression of glycerol release from fat and gluconeogenesis in liver by feeding. We also show a dysregulated increase of AQP9 in the liver and AQPap/7 in the adipose tissue of insulin-resistant hyperinsulinemic mice. Higher amounts of glycerol, released by increased function of AQPap/7, should be taken up into hepatocytes more efficiently via enhanced AQP9, leading to augmentation of hepatic gluconeogenesis. The coordinated changes in two tissue-specific glycerol channels, adipose AQPap/7 and hepatic AQP9, may determine, at least in part, plasma glucose levels during nutritional alterations as well as in insulin-resistant conditions.

From the Department of Internal Medicine and Molecular Science, Graduate School of Medicine, Osaka University, Osaka, Japan.

Address correspondence and reprint requests to Iichiro Shimomura Department of Organismal Biosystems, Graduate School of Frontier Bioscience, Department of Medicine and Pathophysiology, Graduate School of Medicine, Osaka University, 2-2 Yamadaoka, Suita, Osaka 565-0871, Japan. E-mail: ichi@imed2.med.osaka-u.ac.jp.

Received for publication 15 March 2002 and accepted in revised form 26 June 2002.

AQP, aquaporin; AQPap/7, aquaporin adipose; DMEM, Dulbecco's modified Eagle's medium; FFA, free fatty acid; GlyK, glycerokinase; HSL, hormone-sensitive lipase; IRE, insulin response element; IRS, insulin receptor substrate; PEPCK, phosphoenolpyruvate carboxykinase; RACE, rapid amplification of cDNA ends; STZ, streptozotocin.

RESEARCH DESIGN AND METHODS

Materials and general methods. Plasma glycerol was measured by a fluorometric/colorimetric enzyme method (22–25). FFA was determined using Nescauto NEFA kit (Azwel, Osaka, Japan). Glucose and insulin were measured by a Blood Sugar Glucose Oxidase Perid Method kit (Roche Diagnostic, Tokyo, Japan) and a double-antibody enzyme immunoassay using a Glazyme Insulin EIA kit (Sanyo Chemical Industries, Osaka, Japan) with rat insulin as a standard, respectively.

Animals and cells. All experimental protocols described in this article were approved by the Ethics Review Committee for Animal Experimentation of Osaka University School of Medicine. Eight-week-old male ICR (MCH), C57BL/6J, C57BL/KsJ (*db+/+m*), and C57BL/KsJ (*db+/db+*) mice were purchased from Clea Japan (Osaka, Japan). The animals were housed in an air-conditioned room with a 12/12-h dark-light cycle (8:00 A.M.–8:00 P.M.) and acclimated to the new environment for 1 week before the experiment. For experiments on insulin deficiency, streptozotocin (STZ; Sigma Aldrich, Tokyo, Japan) or PBS was administered intraperitoneally (100 mg/kg in 0.05 mol/l citrate buffer, pH 4.5) into 9-week-old male ICR (MCH) mice. Three days later, both groups of mice were anesthetized with 5 mg/ml pentobarbital sodium salt before being killed.

For experiments designed to study the effects of fasting and refeeding, male C57BL/6J mice (each group, $n = 4$) were allowed free access to standard laboratory diet and tap water for 12 h after a 24-h fast. The fasted group was fasted for 12 or 18 h before being killed. The refeed group had free access to standard laboratory diet for 12 h after an 18-h fast. For comparison between the obese and lean controls, C57BL/KsJ (*db+/db+*) and C57BL/KsJ (*db+/+m*) mice on 3 h fasting were used. All mice were phlebotomized from the tail and portal vein.

A rat hepatoma H4IIE cell line (passage no. 6) was obtained from Dainippon Pharmaceutical (Osaka, Japan). H4IIE rat hepatoma cells were grown as monolayer cultures in low-glucose Dulbecco's minimal essential medium (1.0 g/l) supplemented with 10% FCS and incubated in a humidified 95% air–5% CO₂ atmosphere at 37°C until they reached confluence (60–70%), when they were collected for analysis (28). H4IIE hepatoma cells (preconfluent state) were washed twice with calcium- and magnesium-free PBS and incubated with Dulbecco's modified Eagle's medium (DMEM) containing 0.5% fatty acid free BSA (Sigma) for 12 h. After cells were washed twice with PBS, they were incubated in DMEM with 10 nmol/l insulin for 0, 6, or 24 h or with 0, 10, or 100 nmol/l insulin for 24 h. Total RNA was isolated and subjected to Northern blotting for AQP9.

Isolation of mouse AQP9 gene. The primer was designed to amplify the coding region of mouse AQP9, according to the sequence of rat AQP9 (AF016406) (26), which was cloned from rat. The following forward and reverse primers were used for specific PCR amplification: 5'-GGCAGAAACCCCAA GATGCTTC-3' and 5'-CTGCAGATCTGGCCATCCCACTACATGATG-3'. RT-PCR was performed by using the template of mouse liver mRNA.

Determination of 5' and 3' end of mouse AQP9 by 5' and 3' rapid amplification of cDNA ends PCR. For determining the 5' and 3' ends of mouse AQP9, the 5' and 3' end rapid amplifications of the cDNA ends (5' and 3' RACE) were conducted according to the protocol provided by the manufacturer (Marathon cDNA amplification kit; BD Biosciences Clontech Japan, Tokyo, Japan), using the following primers: first primer 5'-TTCTTCTT GGCTCGGTCCTTCTCAGAAGGCATCTT-3' (italicized translation start codon ATG), and nested primer, 5'-CTTGGGGTTTCTGCCAGAGGTAAGCTTTC CACCA-3' for 5' RACE; and first primer 5'-GAACAGCCGAGAACAATCTA GAAAAACAC-3' and nested primer 5'-AGAAAAACACGAACTCAGCGTC ATCATGTAG-3' (italicized translation stop codon TAG), for 3' RACE, respectively.

Isolation of mouse AQP9 promoter. Cloning of the 5' flanking region and promoter of the mouse AQP9 gene was performed using the GenomeWalker kit (BD Biosciences Clontech Japan) according to the protocol recommended by the manufacturer, with the following primers: primer 5'-TGCACGGGT TCTTCTTGGCTCGGTCCTT-3', 5'-ACGAGGTTCTTCTTGGCTCGGTCCT TCTC-3', 5'-TTCTTCTTGGCTCGGTCCTTCTCAGAAGGCATCTT-3' (italicized translation start codon ATG).

Luciferase assay. The mouse AQP9 promoter regions (–667/–1) were amplified from mouse genomic DNA using a Mlu I site-added 5' primer and Xho I site-added 3' primers. The mouse AQP9 promoter-luciferase reporter plasmids were constructed by excising the amplified promoter fragment of AQP9 and inserting it into the Mlu I and Xho I site of the control pGL3 basic luciferase expression vector (Promega, Madison, WI). Partial deletion mutant of pGL3-AQP9 luciferase plasmid was constructed using the QuickChange Site-Directed Mutagenesis kit (Stratagene, La Jolla, CA). IRE-deleted constructs (ΔIRE) were designed from the wild-type construct by deleting the IRE region (–502/–496). The plasmids for transfection were purified using the Endfree Plasmid kit (Qiagen, Valencia, CA).

Primary hepatocytes were isolated from nonfasting male Sprague-Dawley rats (approximate body weight 250 g), as described previously (29). Briefly, the prepared hepatocytes were attached to a 12-well plate in DMEM supplemented with 10% FCS and 100 nmol/l triiodothyronine. Four hours later, the medium was replaced with Medium 199 (M199) supplemented with 100 nmol/l dexamethasone, 100 nmol/l triiodothyronine, and 1 nmol/l insulin. For each 12-well culture plate, 50 ng of firefly (*Photinus pyralis*) luciferase plasmid, constructed from pGL3-basic luciferase expression vector and 10 ng of a sea pansy (*Renilla reniformis*) luciferase pRL-SV40 plasmid (Promega), were complexed with LipofectAMINE2000 (Life Technologies, Tokyo, Japan) following the protocol provided by the manufacturer and then used for transfection. The transfection mixture was removed 24 h after transfection, and the cells were maintained in M199 containing the indicated concentration of insulin. The cells were harvested with passive lysis buffer (Promega), and luciferase activity was measured with the Dual Luciferase Reporter Assay System (Promega) using the procedure described by the manufacturer.

Construction of cDNA probes and RNA analysis. Mouse AQP9, AQP9, glycerokinase (GlyK), phosphoenolpyruvate carboxykinase (PEPCK), and hormone-sensitive lipase (HSL) cDNAs were synthesized by RT-PCR as reported previously (22,30). Insulin receptor substrate (IRS)-1 or -2 cDNA probe was described previously (30). Total cellular RNA was extracted using TRIZOL reagent kit (Invitrogen, Tokyo, Japan). Northern blot analysis was performed as described previously (23). Abundance of mRNAs was determined by densitometric analysis using FAST SCAN Scanning Imager (Molecular Dynamics, Buckinghamshire, U.K.) and expressed in arbitrary units.

Immunoblot of AQP9. Immunoblot analysis of AQP9 protein was performed as reported previously (22). Briefly, the membrane fractions of adipose tissues were obtained. The proteins (40 μg) were loaded and separated on a 12.5% SDS-polyacrylamide gel and transferred onto nitrocellulose transfer membrane (Schleicher & Schuell, Keene, NH). Western blot analysis was performed using antiserum to AQP9 at a dilution of 1:500. Horseradish peroxidase-coupled donkey anti-rabbit immunoglobulins were used at a dilution of 1:3,000. Detection by chemiluminescence was performed using the ECL system (Amersham Pharmacia Biotech, Tokyo, Japan).

Mouse liver perfusion studies. Mouse liver perfusion studies were performed as described previously, with a minor modification (31,32). After 18 h of fasting, the male C57BL/KsJ (*db+/db+*) and (*db+/+m*) mice ($n = 4$, each) were anesthetized with an intraperitoneal injection of 5 mg/ml pentobarbital sodium. The abdominal cavity was opened, and the superior mesenteric artery and bilateral renal arteries and veins were ligated. Then, 0.25 ml (25 units) of heparin was injected into the inferior vena cava and the liver was perfused via the portal vein at a constant flow rate of 4 ml/min. The perfusate was albumin-free Krebs buffer (in mmol/l: NaCl 120, KH₂PO₄ 1.2, MgSO₄ · 7H₂O 1.2, KCl 4.8, CaCl₂ · 2H₂O 1.72, and NaHCO₃) at 37°C, bubbled with 95% O₂–5% CO₂ gas mixture. The concentration of glycerol infused into the portal vein was increased every 20 min from 200 to 400, 600, and 800 μmol/l. A recovery catheter was inserted into the inferior vena cava, and samples were collected every 10 min after reperfusion for 30 min and stored at –80°C for later analysis. At the end of perfusion, the whole liver was removed and weighed. Glucose concentration in the recovery samples was measured using the Blood Sugar Glucose Oxidase Perid Method kit.

Statistical analysis. All data were expressed as mean ± SE. Differences between groups were examined for statistical significance using the Student's *t* test. $P < 0.05$ denoted the presence of a statistically significant difference.

RESULTS

Cloning of the mouse AQP9 gene. To investigate the regulation of AQP9, we cloned mouse AQP9 cDNA from the mouse liver. Figure 1 illustrates the amino acid alignments of AQP9 among mouse, rat, and human. The mouse AQP9 cDNA encodes a 295 amino acid (GenBank accession no. AB 037180) and shows 90% homology to rat AQP9 (GenBank accession no. AF 016406) (26) and 76% homology to human AQP9 (GenBank accession no. XM 012421) (33,34). Hydrophathy analysis showed that mouse AQP9 has six putative transmembrane domains, similar to human and rat AQP9. Mouse AQP9 has two Asn-Pro-Ala consensus motifs, characteristic of members of the major intrinsic protein family (20).

Effect of fasting/refeeding on AQP9, GlyK, and PEPCK mRNA expressions. Fasting activates lipolysis and glycerol release from adipose tissue (13). We recently

	1	30
m AQP9	MPSEKDRAKKNLVQRLALKSCLAKETLSEF	
r AQP9	MPSEKDGAKKSLMQRLALKSRIAKETLSEF	
h AQP9	MQPEGAEKGKSFQRLVLKSSLAKETLSEF	
m AQP9	LGTFIMIVLGCSSIAQAVLSREKAGGIITI	
r AQP9	LGTFIMIVLGCSSIAQAVLSRERFGGIITI	
h AQP9	LGTFILIVLGCSCVAQAILSRGRFGGVITI	
	①	
m AQP9	NIGFATAVVMALYATFGVSGGHINPAVSFA	
r AQP9	NIGFASAVVMALYVTFGISGGHINPAVSFA	
h AQP9	NVGFSAVAMAIYVAGGVSGGHINPAVSLA	
	②	
m AQP9	MCTFGRMEWFKFPFYVGAQLLGAFLVGAATV	
r AQP9	MCAFGRMEWFKFPFYVGAQFLGAFLVGAATV	
h AQP9	MCLFGRMKWFKLPFYVGAQFLGAFLVGAATV	
m AQP9	FGIYYDGLMAFADGKLLITGENGTAFIFAT	
r AQP9	FGIYYDGLMAFAGGKLLVGENATAFIFAT	
h AQP9	FGIYYDGLMSFAGGKLLIVGENATAHIFAT	
	③	
m AQP9	YKPFVSVPGAFVDQVVSTMFLLLIVFAIF	
r AQP9	YPAPFISTPGAFVDQVVSTMFLLLIVFAMF	
h AQP9	YPAPYLSLANAFADQVVATMILLIIVFAIF	
	④	
m AQP9	DSRNLGVPRGLEPVIIGLLIIVISCSLGLN	
r AQP9	DSRNLGVPRGLEPVVIGLLIIVLSCSLGLN	
h AQP9	DSRNLGAPRGLEPIAIGLLIIVIASLGLN	
	⑤	
m AQP9	SGCAMNPARDLSPRLFTALAGWGFEVFTFG	
r AQP9	SGCAMNPARDLSPRLFTALAGWGFEVFTVG	
h AQP9	SGCAMNPARDLSPRLFTALAGWGFEVFRAG	
m AQP9	NNFWWIPVVGPMIGAVLGGLIYVLFIQMHH	
r AQP9	NNFWWIPVVGPMIGAVLGGLIYVLFIQMHH	
h AQP9	NNFWWIPVVGPLVGAIVIGGLIYVLFVIEIHH	
	⑥	295
m AQP9	SNPDPEVKAEPAAENNLEKHELSVIM	
r AQP9	SKLDPDMKAEPSENNLEKHELSVIM	
h AQP9	PEPDSVFKTEQSEDKPEKYELSVIM	

FIG. 1. Deduced amino acid sequence of mouse AQP9. Amino acid sequence alignments of mouse AQP9 (m; GenBank accession no. AB 037180) isolated from the mouse liver, rat AQP9 (r; GenBank accession no. AF 016406) (24), and human AQP9 (h; GenBank accession no. XM 012421) (31). Asn-Pro-Ala (NPA) motifs are boxed, and putative membrane-spanning regions are underlined and numbered 1–6 inside a circle. The conserved residues are indicated by bold letters.

reported that AQPap mRNA expression in white adipose tissue was enhanced by fasting and suppressed by refeeding and that the changes in AQPap mRNA showed a pattern similar to that of plasma glycerol (23). Glycerol becomes a substrate for gluconeogenesis in the liver that expresses GlyK (35). Figure 2 shows the effects of fasting and refeeding on the regulation of hepatic mRNAs of AQP9 and GlyK. Portal plasma levels of glycerol and FFA increased during fasting and decreased after refeeding (Fig. 2A). In contrast, portal plasma insulin levels decreased during fasting and increased after refeeding (Fig. 2A). The mRNA levels of AQP9 and GlyK increased significantly by fasting and decreased by refeeding, similar to the change in PEPCK mRNA (Fig. 2B) (36), which is known to be downregulated by insulin. The coordinated enhancements of AQPs (22), GlyK, and PEPCK are effective to supply plasma glycerol for hepatic gluconeogenesis in fasting. Similar effects of fasting on these hepatic mRNAs were also seen when fasting started at 0:00 h.




FIG. 2. Effect of fasting/refeeding on expression of AQP9, GlyK, and PEPCK mRNAs. Mice were killed after 0, 12, and 18 h of fasting or 12 h of refeeding after 18 h of fasting and then the livers were removed for analysis. A: Plasma FFA, glycerol, and insulin in the portal vein were measured and the data are mean \pm SE. B: Northern blots using pooled RNA samples (10 μ g/lane). Abundance of mRNAs was determined by densitometric analysis using FAST SCAN Scanning Imager (Molecular Dynamics, Buckinghamshire, U.K.) and is presented in arbitrary units. Autoradiographs of the blot were inserted.

Effect of insulin on AQP9 mRNA expression in mouse liver and H4IIE cells. Next, we examined the regulation of hepatic AQP9 mRNA by insulin (Fig. 3). Hepatic mRNA levels of AQP9 and GlyK were high in insulin-deficient STZ-treated mice, similar to that of PEPCK mRNA (Fig. 3A). Furthermore, insulin downregulated the expression levels of AQP9 mRNA in H4IIE hepatoma cells in dose- and time-dependent manners (Fig. 3B). These results suggested that the AQP9 gene expression is directly downregulated by insulin, similar to PEPCK mRNA.

Promoter of mouse AQP9 gene. The 5' flanking region of the mouse AQP9 gene was sequenced (GenBank accession no. AB 073723), and transcriptional initiation site of the AQPap/7 gene was determined by 5' RACE using total RNA isolated from mouse liver (Fig. 4A). A search of the promoter region of the AQP9 gene for canonical consensus sequences revealed the presence of several putative binding sites for transcription factors (Fig. 4A). Several binding sites for CCAAT enhancer binding protein and hepatic nuclear factor-3 β were identified in the promoter (23).

Negative IRE in the mouse AQP9 gene promoter. In the mouse AQP9 gene promoter, we identified a region similar to the core negative IRE, T(G/A)TTTT(G/T) (27,37), which is also found in promoters of various genes, including PEPCK (38) and AQPap/7 (23,24) (Fig. 4B). For




FIG. 5. Metabolic profiles and AQPap/7 and AQP9 mRNA expression in *db/db+* mice. **A**: Plasma concentrations of insulin, glucose, and glycerol in *db+/+m* (lean) and *db/db+* (obese) mice ($n = 4$, each) after 3 h of fasting. Data are mean \pm SE, and comparisons are between lean and obese (Student's *t* test). **B**: Northern blotting of AQPap/7 and HSL and Western blotting of AQPap/7 from the mesenteric fat of *db+/+m* and *db/db+* mice. Northern and Western blotting were performed as described in RESEARCH DESIGN AND METHODS. **C**: Northern blotting of AQP9, IRS-1, and IRS-2 from the livers of *db+/+m* and *db/db+* mice. For **B** and **C**, the fold changes in obese mice, compared with lean mice, were calculated from the results of individual animals in each group.

insulin-mediated suppression on the transcription of the mouse AQP9 gene.

Metabolic profile and expression of AQPap/7 and AQP9 mRNAs in *db/db+* mice. The *db/db+* mice were significantly insulin resistant. Plasma glycerol levels in the portal vein were higher in *db/db+* mice. They had a higher concentration of plasma glucose despite severe hyperinsulinemia (Fig. 5A). Figure 5B and C demonstrates the results of Northern blotting of mesenteric fat and liver, respectively. The mRNA and protein amounts of AQPap/7 and HSL mRNA were higher in the mesenteric fat of *db/db+* mice (Fig. 5B), although both are negatively regulated by insulin. The hepatic AQP9 mRNA was also induced despite hyperinsulinemia (Fig. 5C). Hepatic insulin resistance was accompanied by markedly reduced IRS-2 (Fig. 5C), as reported previously (30).

Hepatic glucose production from glycerol in control and *db/db+* mice. Glycerol is the main gluconeogenic substrate in prolonged fasting (15–17). Hepatic glucose production from glycerol was compared between *db+/+* and *db/db+* mice in the 18-h fasted state using a previously described hepatic perfusion procedure (31,32). Glucose production increased in parallel with the amount of glycerol perfused into the portal vein in *db+/+m* and *db/db+* mice (Fig. 6). Glucose output was significantly higher in the liver of *db/db+* mice than *db+/+m* mice. These results indicated that the glycerol-mediated gluconeogenic activity was augmented in the liver of *db/db+* mice (39), where the mRNAs of AQP9 and gluconeogenic enzymes were augmented.




FIG. 6. Hepatic glucose production from glycerol in control and *db/db+* mice. Hepatic glucose production from glycerol in *db/db+* (obese) and *db+/+m* (lean) mice ($n = 4$, each) after 18 h of overnight fasting was measured using the hepatic perfusion procedure as described in RESEARCH DESIGN AND METHODS. Glycerol was infused from the portal vein at the indicated concentration, and the concentration was increased every 20 min. The recovery samples were collected from the inferior vena cava every 10 min, and glucose concentration was measured. Data are mean \pm SE; * $P < 0.05$, between lean and obese mice (Student's *t* test).

DISCUSSION

Glycerol is one of the major factors that determine plasma glucose (40). During fasting, hepatic glucose output is the main source of plasma glucose, and glycerol taken up from plasma becomes a major substrate for hepatic gluconeogenesis (14). However, adipose tissue is the major source of plasma glycerol (13). Therefore, effective systems are required during fasting that enhance glycerol release from adipose tissue and glycerol uptake into the liver. With regard to the glycerol transport proteins, three aquaglyceroporins are known: AQPap/7, AQP3, and AQP9 (20,22). To date, AQPap/7 is the only aquaglyceroporin expressed in adipose tissues, and AQP9 is the only aquaglyceroporin expressed in the liver (22). It is conceivable that during fasting, AQPap/7 in fat is essential for the supply of glycerol to the liver and that AQP9 is critical for hepatic uptake of glycerol for glucose production.

Insulin suppresses the amount of AQPap/7 mRNA in adipose tissue (23) and AQP9 in liver (current study). We also identified similar negative IREs in the promoter regions of AQPap/7 (23,24) and AQP9 genes (current study). We identified the potential IRE sequence (–591/–597; TGTTTTC) in the promoter region of the human AQP9 gene (33). These results indicate that the gene expressions of AQPap/7 and AQP9 are regulated coordinately by the plasma concentrations of insulin in accordance with the nutritional condition, such as fasting and refeeding. These regulations should increase or decrease the supply of glycerol and glucose in plasma by fasting or refeeding, respectively.

In hepatic gluconeogenesis, AQP9, which is localized at the sinusoidal plasma membrane (41), serves as a channel for glycerol uptake. GlyK converts the incorporated glycerol into the phosphorylated form to be used for gluconeogenesis, and PEPCK is one of the limiting enzymes to drive the glycerol-independent gluconeogenic pathway (2). In the normal liver, the mRNAs for all of these proteins

are downregulated by insulin (27,37). During fasting, the low concentration of plasma insulin coordinately and effectively enhances all of these mRNAs to drive hepatic gluconeogenesis. However, under pathological conditions of insulin resistance in the *db/db* mice, high mRNA expression levels of hepatic AQP9, GlyK, and PEPCK mRNAs and of adipose AQPap/7 were found despite the high concentration of plasma insulin. Dysregulated increase of AQPap/7 and hepatic AQP9 may underlie the pathophysiology of hyperglycemia in severe insulin resistance. Our recent studies indicated that the dysregulated increase of AQPap/7 in adipose tissues causes higher plasma glycerol in insulin-resistant rodents (22). The liver of insulin-resistant *db+/db+* mice exhibited a high gluconeogenic activity by using glycerol as a substrate (Fig. 6). Hepatic insulin resistance may be accounted for by the significantly low levels of IRS-2. We previously demonstrated that chronic hyperinsulinemia in lipodystrophy and *ob/ob* mice was associated with low hepatic IRS-2, which resulted in increased mRNAs of gluconeogenesis, such as glucose-6-phosphatase and PEPCK (30). There were no apparent changes in adipose mRNAs for IRS-1 and IRS-2 between obese and lean mice (data not shown). In the previous articles, IRS-1-mediated insulin signaling was shown to be deteriorated in the adipose tissue of insulin-resistant animals, with relatively normal IRS-2-mediated pathway (42,43). We previously showed that insulin's inhibition of AQPap/7 was mediated by a phosphatidylinositol 3-kinase-dependent manner (23). Taken together, the increase of AQPap under insulin resistance may be caused by the decreased IRS-1-mediated insulin signaling in the adipose cells. In the present study, it is conceivable that reduced hepatic IRS-2 in *db+/db+* mice resulted in disturbance of insulin signaling, which led to an increase in AQP9 and hepatic glycerol uptake. In the insulin-resistant state, both the level of plasma glycerol and the glucose production associated with glycerol input were increased, which might lead to the more severe hyperglycemia. Coordinated augmentation of AQPap/7 and AQP9 in insulin resistance should increase the supply of fat-derived glycerol as a substrate for hepatic gluconeogenesis, which aggravates hyperglycemia.

Figure 7 represents a summary of the working model based on the results of our previous (21–25) and present studies. In the physiological feeding state, high plasma insulin coordinately suppresses the AQPap/7 mRNA for glycerol release from fat and AQP9 mRNA for glycerol uptake into the liver, through the negative IRE in AQPap/7 and AQP9 gene promoters. In contrast, in the insulin-resistant state, reciprocal increases of AQPap/7 and AQP9 despite hyperinsulinemia lead to more utilization of the fat-derived glycerol for hepatic glucose production and release. Identifying the responsible transcriptional factors and associated proteins involved in the transcriptional regulation of AQPap/7 and AQP9 genes and defining the precise mechanism that modifies the amounts and activities of these proteins by insulin should provide pivotal insight for the design of new pharmaceutical strategies to combat insulin resistance syndrome.

ACKNOWLEDGMENTS

This work was supported in part by the fund from the "Research for the Future" Program from the Japan Society for the Promotion of Science (JSPS-RFTF97L00801) and




FIG. 7. Physiological and pathophysiological involvement of AQPap/7 and AQP9. In the physiological feeding state, high plasma insulin suppresses AQPap/7 mRNA in the fat and AQP9 mRNA in the liver through the negative IRE in AQPap/7 and AQP9 gene promoters. These coordinated regulations efficiently reduce the supply of glycerol from fat to liver for hepatic glucose production. In contrast, in the pathological state of insulin resistance, increased AQPap/7 and AQP9 mRNAs, despite hyperinsulinemia, augment the utilization of glycerol for hepatic glucose production. These coordinated dysregulations of AQPap/7 and AQP9 aggravate the hyperglycemia in the insulin-resistant state.

Grants-in-Aid from the Ministry of Education, Science, Sports and Culture of Japan (09307019, 10557100, 10557101, 10671035, and 13671189).

We thank Yuko Matsukawa and Sachiyo Tanaka of our laboratory for technical assistance.

REFERENCES

- Ramsay TG: Fat cells. *Endocrinol Metab Clin North Am* 25:847–870, 1996
- Coppack SW, Jensen MD, Miles JM: In vivo regulation of lipolysis in humans. *J Lipid Res* 35:177–193, 1994
- Shimomura I, Tokunaga K, Kotani K, Keno Y, Yanase-Fujiwara M, Kanosue K, Jiao S, Funahashi T, Kobatake T, Yamamoto T, Matsuzawa Y: Marked reduction of acyl-CoA synthetase activity and mRNA in intra-abdominal visceral fat by physical exercise. *Am J Physiol* 265:E44–E50, 1993
- Amer P, Kriegholm E, Engfeldt P, Bolinder J: Adrenergic regulation of lipolysis in situ at rest and during exercise. *J Clin Invest* 85:893–898, 1990
- Reaven GM: Role of insulin resistance in human disease. *Diabetes* 37:1595–1607, 1988
- Kaplan NM: How bad are diuretic-induced hypokalemia and hypercholesterolemia? *Arch Intern Med* 149:1514–1520, 1989
- Kissebah AH, Vydelingum N, Murray R, Evans DJ, Hartz AJ, Kalkhoff RK, Adams PW: Relation of body fat distribution to metabolic complications of obesity. *J Clin Endocrinol Metab* 54:254–260, 1982
- DeFronzo RA, Ferrannini E: Insulin resistance: a multifaceted syndrome responsible for NIDDM, obesity, hypertension, dyslipidemia, and atherosclerotic cardiovascular disease. *Diabetes Care* 14:173–194, 1991
- Matsuzawa Y, Shimomura I, Nakamura T, Keno Y, Tokunaga K: Pathophysiology and pathogenesis of visceral fat obesity. *Ann N Y Acad Sci* 676:270–278, 1993
- Baynes C, Henderson AD, Hughes CL, Richmond W, Johnston DG, Ellekes RS: Determinants of mild fasting hypertriglyceridaemia in non-insulin-dependent diabetes. *J Intern Med* 229:267–273, 1991
- Large V, Arner P: Regulation of lipolysis in humans: pathophysiological modulation in obesity, diabetes, and hyperlipidaemia. *Diabetes Metab* 24:409–418, 1998

12. Kuriyama H, Yamashita S, Shimomura I, Funahashi T, Ishigami M, Aragane K, Miyaoka K, Nakamura T, Takemura K, Man Z, Toide K, Nakayama N, Fukuda Y, Lin MC, Wetterau JR, Matsuzawa Y: Enhanced expression of hepatic acyl-coenzyme A synthetase and microsomal triglyceride transfer protein messenger RNAs in the obese and hypertriglyceridemic rat with visceral fat accumulation. *Hepatology* 27:557–562, 1998
13. Baba H, Zhang XJ, Wolfe RR: Glycerol gluconeogenesis in fasting humans. *Nutrition* 11:149–153, 1995
14. Hetenyi G Jr: Gluconeogenesis in vivo. *Am J Physiol* 249:R792–R793, 1985
15. Landau BR, Wahren J, Chandramouli V, Schumann WC, Ekberg K, Kalhan SC: Contributions of gluconeogenesis to glucose production in the fasted state. *J Clin Invest* 98:378–385, 1996
16. Peroni O, Large V, Beylot M: Measuring gluconeogenesis with [2-¹³C]glycerol and mass isotopomer distribution analysis of glucose. *Am J Physiol* 269:E516–E523, 1995
17. Peroni O, Large V, Odeon M, Beylot M: Measuring glycerol turnover, gluconeogenesis from glycerol, and total gluconeogenesis with [2-¹³C] glycerol: role of the infusion-sampling mode. *Metabolism* 45:897–901, 1996
18. Jensen MD, Chandramouli V, Schumann WC, Ekberg K, Previs SF, Gupta S, Landau BR: Sources of blood glycerol during fasting. *Am J Physiol Endocrinol Metab* 281:E998–E1004, 2001
19. Engel A, Fujiyoshi Y, Agre P: The importance of aquaporin water channel protein structures. *EMBO J* 19:800–806, 2000
20. Verkman AS, Mitra AK: Structure and function of aquaporin water channels. *Am J Physiol* 278:F13–F28, 2000
21. Kuriyama H, Kawamoto S, Ishida N, Ohno I, Mita S, Matsuzawa Y, Matsubara K, Okubo K: Molecular cloning and expression of a novel human aquaporin from adipose tissue with glycerol permeability. *Biochem Biophys Res Commun* 241:53–55, 1997
22. Kishida K, Kuriyama H, Funahashi T, Shimomura I, Kihara S, Ouchi N, Nishida M, Nishizawa H, Matsuda M, Takahashi M, Hotta K, Nakamura T, Yamashita S, Tochino Y, Matsuzawa Y: Aquaporin adipose, a putative glycerol channel in adipocytes. *J Biol Chem* 275:20896–20902, 2000
23. Kishida K, Shimomura I, Kondo H, Kuriyama H, Makino Y, Nishizawa H, Maeda N, Matsuda M, Ouchi N, Kihara S, Kurachi Y, Funahashi T, Matsuzawa Y: Genomic structure and insulin-mediated repression of the aquaporin adipose (AQPap), adipose-specific glycerol channel. *J Biol Chem* 276:36251–36260, 2001
24. Kondo H, Shimomura I, Kishida K, Kuriyama H, Makino Y, Nishizawa H, Matsuda M, Maeda N, Nagaretani H, Kihara S, Kurachi Y, Nakamura T, Funahashi T, Matsuzawa Y: Human aquaporin adipose (AQPap) gene. *Eur J Biochem* 269:1814–1826, 2002
25. Kishida K, Shimomura I, Nishizawa H, Maeda N, Kuriyama H, Kondo H, Matsud M, Nagaretani H, Ouchi N, Hotta K, Kihara S, Kadowaki T, Funahashi T, Matsuzawa Y: Enhancement of the aquaporin adipose gene expression by a peroxisome proliferator-activated receptor gamma. *J Biol Chem* 276:48572–48579, 2001
26. Tsukaguchi H, Shayakul C, Berger UV, Mackenzie B, Devidas S, Guggino WB, van Hoek AN, Hediger MA: Molecular characterization of a broad selectivity neutral solute channel. *J Biol Chem* 273:24737–24743, 1998
27. O'Brien RM, Streep RS, Ayala JE, Stadelmaier BT, Hornbuckle LA: Insulin-regulated gene expression. *Biochem Soc Trans* 29:552–558, 2001
28. Tomizawa M, Kumar A, Perrot V, Nakae J, Accili D, Rechler MM, Kumaro A: Insulin inhibits the activation of transcription by a C-terminal fragment of the forkhead transcription factor FKHR: a mechanism for insulin inhibition of insulin-like growth factor-binding protein-1 transcription. *J Biol Chem* 275:7289–7295, 2000
29. Shimomura I, Bashmakov Y, Ikemoto S, Horton JD, Brown MS, Goldstein JL: Insulin selectively increases SREBP-1c mRNA in the livers of rats with streptozotocin-induced diabetes. *Proc Natl Acad Sci U S A* 96:13656–13661, 1999
30. Shimomura I, Matsuda M, Hammer RE, Bashmakov Y, Brown MS, Goldstein JL: Decreased IRS-2 and increased SREBP-1c lead to mixed insulin resistance and sensitivity in livers of lipodystrophic and ob/ob mice. *Mol Cell* 6:77–86, 2000
31. Arai K, Lee K, Berthiaume F, Tompkins RG, Yarmush ML: Intrahepatic amino acid and glucose metabolism in a D-galactosamine-induced rat liver failure model. *Hepatology* 34:360–371, 2001
32. Burns SP, Murphy HC, Iles RA, Cohen RD: Lactate supply as a determinant of the distribution of intracellular pH within the hepatic lobule. *Biochem J* 358:569–571, 2001
33. Tsukaguchi H, Weremowicz S, Morton CC, Hediger MA: Functional and molecular characterization of the human neutral solute channel aquaporin-9. *Am J Physiol* 277:F685–F696, 1999
34. Ishibashi K, Kuwahara M, Gu Y, Tanaka Y, Marumo F, Sasaki S: Cloning and functional expression of a new aquaporin (AQP9) abundantly expressed in the peripheral leukocytes permeable to water and urea, but not to glycerol. *Biochem Biophys Res Commun* 244:268–274, 1998
35. Westergaard N, Madsen P, Lundgren K: Characterization of glycerol uptake and glycerol kinase activity in rat hepatocytes cultured under different hormonal conditions. *Biochim Biophys Acta* 1402:261–268, 1998
36. Nandan SD, Beale EG: Regulation of phosphoenolpyruvate carboxykinase mRNA in mouse liver, kidney, and fat tissues by fasting, diabetes, and insulin. *Lab Anim Sci* 42:473–477, 1992
37. O'Brien RM, Granner DK: Regulation of gene expression by insulin. *Physiol Rev* 76:1109–1161, 1996
38. Hall RK, Yamasaki T, Kucera T, Waltner-Law M, O'Brien R, Granner DK: Regulation of phosphoenolpyruvate carboxykinase and insulin-like growth factor-binding protein-1 gene expression by insulin: the role of winged helix/forkhead proteins. *J Biol Chem* 275:30169–30175, 2000
39. Andrikopoulos S, Proietto J: The biochemical basis of increased hepatic glucose production in a mouse model of type 2 (non-insulin-dependent) diabetes mellitus. *Diabetologia* 38:1389–1396, 1995
40. Previs SF, Brunengraber H: Methods for measuring gluconeogenesis in vivo. *Curr Opin Clin Nutr Metab Care* 1:461–465, 1998
41. Elkjaer M, Vajda Z, Nejsum LN, Kwon T, Jensen UB, Amiry-Moghaddam M, Frokiaer J, Nielsen S: Immunolocalization of AQP9 in liver, epididymis, testis, spleen, and brain. *Biochem Biophys Res Commun* 276:1118–1128, 2000
42. Di Guglielmo GM, Drake PG, Baass PC, Authier F, Posner BI, Bergeron JJ: Insulin receptor internalization and signalling. *Mol Cell Biochem* 182:59–63, 1998
43. Esposito DL, Li Y, Cama A, Quon MJ: Tyr(612) and Tyr(632) in human insulin receptor substrate-1 are important for full activation of insulin-stimulated phosphatidylinositol 3-kinase activity and translocation of GLUT4 in adipose cells. *Endocrinology* 142:2833–2840, 2001