



Marta Y. Pepino,<sup>1</sup> Sarah A. Eisenstein,<sup>2,3</sup> Allison N. Bischoff,<sup>2</sup> Samuel Klein,<sup>1</sup> Stephen M. Moerlein,<sup>3,4</sup> Joel S. Perlmutter,<sup>3,5,6,7,8</sup> Kevin J. Black,<sup>2,3,5,6</sup> and Tamara Hershey<sup>2,3,5</sup>

## Sweet Dopamine: Sucrose Preferences Relate Differentially to Striatal D<sub>2</sub> Receptor Binding and Age in Obesity

*Diabetes* 2016;65:2618–2623 | DOI: 10.2337/db16-0407

**Alterations in dopaminergic circuitry play a critical role in food reward and may contribute to susceptibility to obesity. Ingestion of sweets releases dopamine in striatum, and both sweet preferences and striatal D<sub>2</sub> receptors (D2R) decline with age and may be altered in obesity. Understanding the relationships between these variables and the impact of obesity on these relationships may reveal insight into the neurobiological basis of sweet preferences. We evaluated sucrose preferences, perception of sweetness intensity, and striatal D2R binding potential (D2R BP<sub>ND</sub>) using positron emission tomography with a D2R-selective radioligand insensitive to endogenous dopamine, (*N*-[<sup>11</sup>C] methyl)benperidol, in 20 subjects without obesity (BMI 22.5 ± 2.4 kg/m<sup>2</sup>; age 28.3 ± 5.4 years) and 24 subjects with obesity (BMI 40.3 ± 5.0 kg/m<sup>2</sup>; age 31.2 ± 6.3 years). The groups had similar sucrose preferences, sweetness intensity perception, striatal D2R BP<sub>ND</sub>, and age-related D2R BP<sub>ND</sub> declines. However, both striatal D2R BP<sub>ND</sub> and age correlated with sucrose preferences in subjects without obesity, explaining 52% of their variance in sucrose preference. In contrast, these associations were absent in the obese group. In conclusion, the age-related decline in D2R was not linked to the age-related decline in sweetness preferences, suggesting that other, as-yet-unknown mechanisms play a role and that these mechanisms are disrupted in obesity.**

Obesity is caused by ingesting more energy than expended over a long period of time. However, the mechanisms

involved in regulating food intake are complex and not completely understood. A growing body of evidence suggests that neural circuits involved in reward and motivation pathways, typically studied in the context of addiction, interact with classic homeostatic regulatory brain areas to cause hedonic hyperphagia and contribute to the development of obesity (1). Data from several studies that involved blood oxygen level-dependent (BOLD) functional MRI have shown that food consumption elicits a smaller BOLD response in the dorsal striatum in subjects with obesity than those who are lean (2,3), presumably mediated by reduced striatal dopaminergic functioning.

The dopamine system plays a critical role in mediating the rewarding effects of food ingestion and thereby influences ingestive behavior. For example, dopamine-deficient rodents stop eating and die of starvation, but these animals can be rescued by restoring dopamine signaling either systemically (4) or within the dorsal, but not ventral, striatum (5). The evolutionarily conserved organization of the striatum serves, in part, to differentially encode and regulate the hedonic value of simply tasting sweetness in the ventral striatum and the postingestive rewarding effects of sugar in the dorsal striatum (6).

A heightened preference for intense sweetness is innate and a hallmark of youth (7). From late adolescence to adulthood, however, sweetness preference declines in both rodents (8) and people (7,9–11). In addition, data from studies conducted in rodent models demonstrate

<sup>1</sup>Atkins Center of Excellence in Obesity Medicine, Center for Human Nutrition, Washington University School of Medicine in St. Louis, St. Louis, MO

<sup>2</sup>Department of Psychiatry, Washington University School of Medicine in St. Louis, St. Louis, MO

<sup>3</sup>Department of Radiology, Washington University School of Medicine in St. Louis, St. Louis, MO

<sup>4</sup>Department of Biochemistry and Molecular Biophysics, Washington University School of Medicine in St. Louis, St. Louis, MO

<sup>5</sup>Department of Neurology, Washington University School of Medicine in St. Louis, St. Louis, MO

<sup>6</sup>Department of Neuroscience, Washington University School of Medicine in St. Louis, St. Louis, MO

<sup>7</sup>Department of Occupational Therapy, Washington University School of Medicine in St. Louis, St. Louis, MO

<sup>8</sup>Department of Physical Therapy, Washington University School of Medicine in St. Louis, St. Louis, MO

Corresponding author: Marta Y. Pepino, pepinodegruevmy@wustl.edu.

Received 28 March 2016 and accepted 31 May 2016.

© 2016 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <http://diabetesjournals.org/site/license>.

that the decline in sweetness preference is attenuated by obesity (8). The mechanisms responsible for the age-related decline in sweetness preference and the alteration in sweetness preference associated with obesity are not known (12). The main purpose of this study was to test the hypothesis that the age-related decline in sweet preferences is directly related to the age-related decline in striatal D<sub>2</sub> receptors (D2R) and that this relationship is altered in people with obesity.

## RESEARCH DESIGN AND METHODS

### Participants

Twenty-four subjects with obesity (BMI  $\geq 30.0$  kg/m<sup>2</sup>) and 20 subjects without obesity (BMI  $< 26.0$  kg/m<sup>2</sup>) between the ages of 18 and 40 years old (Table 1) provided written informed consent and participated in this study, which was approved by the institutional review board of the Washington University School of Medicine in St. Louis. Selected data from these subjects have been reported previously (13–15). Potential subjects completed a comprehensive medical evaluation that included an oral glucose tolerance test (OGTT); those who had diabetes, smoked tobacco, were taking medications that affect dopamine function, had an IQ  $< 80$ , or had neurological conditions including parkinsonism, lifetime psychosis, mania, substance dependence, major depression, social phobia, or eating disorders and panic disorders determined by neurological examination and psychiatric interview were excluded.

### Taste Testing

The following tests were performed  $\sim 1$  h after subjects completed the OGTT and consumed a standardized light snack.

### Sucrose Intensity Perception

To ensure differences in preferences were not confounded by differences in perception of stimuli strength, we determined subjects' perception of sweet taste intensity across a range of sucrose concentrations by using the general label magnitude scale (16), as previously described (17).

### Sucrose Preferences

To determine the subjects' most preferred intensity of sweetness, we used the Monell two-series, forced-choice tracking procedure (11). Subjects were presented with pairs of solutions that differed in sucrose concentration (3%–36% w/v), and preferences were determined as previously described (11,17).

### Positron Emission Tomography and MRI Acquisition and Analyses

On a separate visit, an average of 13.9 days (SD 17.1) after taste testing, subjects underwent functional MRI and 2-h positron emission tomography (PET) scans. Methods for (*N*-[<sup>11</sup>C] methyl)benperidol ([<sup>11</sup>C]NMB) synthesis, MRI, and PET scan acquisitions and preprocessing were described previously (13,14). Each participant received an intravenous infusion of a D2R-selective radioligand not sensitive to endogenous dopamine, [<sup>11</sup>C]NMB (6.4–18.1 mCi) over 20 s. [<sup>11</sup>C]NMB purity was  $\geq 96\%$  and specific activity was  $\geq 1,066$  Ci/mmol (39 TBq/mmol). PET scans were done with the Siemens/CTI ECAT EXACT HR+ tomograph (18). Emission data were collected in 3D mode for 2 h with a total of 30 frames: 3  $\times$  1 min, 4  $\times$  2 min, 3  $\times$  3 min, and 20  $\times$  5 min. Transaxial and axial spatial resolution at slice center are 4.3 mm and 4.1 mm full width at half maximum in 3D mode.

The methods for our region of interest–based analyses are described previously (13,14). We used FreeSurfer for segmentation of striatal regions (19). To limit multiple comparisons, we averaged D2R specific binding (nondisplaceable binding potential [BP<sub>ND</sub>]) across left and right hemispheres for each region of interest. Putamen and caudate D2R BP<sub>ND</sub> values were averaged to obtain a composite dorsal striatal BP<sub>ND</sub>, and nucleus accumbens D2R BP<sub>ND</sub> was used to determine ventral striatal BP<sub>ND</sub>. The midbrain region of each individual's magnetization-prepared rapid gradient-echo was traced as previously described (13).

### Statistical Analyses

The statistical significance of values between groups was evaluated by using Pearson's  $\chi^2$  or unpaired *t* test as appropriate, and differences in sweetness intensity perception between groups was evaluated by using a mixed ANOVA. We used separate linear regression models for dorsal and ventral striatum to analyze the ability of age and D2R BP<sub>ND</sub> to predict sucrose preferences. Age  $\times$  group and D2R BP<sub>ND</sub>  $\times$  group interactions were included in the model to test the hypothesis that group affected the strength of relationships between main factors. Analyses were repeated using race and sex as covariates

**Table 1—Subject characteristics**

	Without obesity ( <i>n</i> = 19)	With obesity ( <i>n</i> = 22)	<i>P</i>
Age, years	28.3 $\pm$ 5.4	31.2 $\pm$ 6.3	0.12
Sex, <i>n</i> (men/women)	4/15	3/19	0.41
Height, cm	170.0 $\pm$ 8.4	166.8 $\pm$ 8.4	0.23
Weight, kg	64.5 $\pm$ 9.9	111.9 $\pm$ 17.9	$< 0.001$
BMI, kg/m <sup>2</sup>	22.5 $\pm$ 2.4	40.3 $\pm$ 5.0	$< 0.001$
Glucose, mg/dL	89.2 $\pm$ 6.3	97.5 $\pm$ 7.0	$< 0.001$
Insulin, $\mu$ IU/mL	5.5 $\pm$ 3.1	17.0 $\pm$ 8.8	$< 0.001$
HOMA of insulin resistance	1.2 $\pm$ 0.7	4.1 $\pm$ 2.2	$< 0.001$
D2R BP <sub>ND</sub>			
Dorsal striatum	8.13 $\pm$ 1.0	8.11 $\pm$ 1.1	0.95
Nucleus accumbens	2.11 $\pm$ 0.2	1.99 $\pm$ 0.4	0.21
Race (%)			0.08
White	84	54	
Black	11	41	
Other/mixed	5	5	
Education, years	16.1 $\pm$ 1.6	15.1 $\pm$ 1.8	0.08

Data are mean  $\pm$  SD, unless otherwise indicated.

in step 1 of hierarchical multiple linear regression models. Data in the tables and figures are presented as mean  $\pm$  SD, unless otherwise indicated.

## RESULTS

### Subject Characteristics

Subjects with and without obesity did not significantly differ in age, height, years of education, sex, or race distribution, but subjects with obesity had higher plasma glucose and insulin concentrations and were more insulin resistant, as indicated by the HOMA of insulin resistance score. As previously reported (15), both groups had similar striatal BP<sub>ND</sub> (Table 1). With the exception of two participants who were overweight (BMI 25.9 and 25.1 kg/m<sup>2</sup>), all other subjects in the group without obesity were normal weight (BMI 18.6–24.6 kg/m<sup>2</sup>). Two of the 24 subjects with obesity were excluded from the analyses because their sucrose preferences testing was unreliable, and one subject without obesity was excluded because his sucrose preference was more than 2 SD above the group mean and his sweetness perception was markedly blunted.

### Sweet Taste Testing

Sucrose preferences ( $P = 0.41$ ) (Fig. 1A) and perception of sweetness intensity were similar between groups (group  $\times$  sucrose concentration interaction  $P = 0.37$ ) (Fig. 1B).

### Relationships Among Age, Central D2Rs, and Sucrose Preferences


Consistent with our previous finding in a subset of these subjects (13), there was an age-related decline in dorsal ( $r = -0.42$ ;  $P < 0.01$ ), but not ventral ( $r = -0.23$ ;  $P = 0.14$ ), D2R BP<sub>ND</sub> in both groups. However, the relationship between age, striatal D2R BP<sub>ND</sub>, and sucrose preferences were markedly different in groups with and without obesity (model for dorsal striatum:  $R^2 = 0.32$ ,  $F_{4,36} = 4.32$ ,  $P < 0.01$ ; both interactions: dorsal D2R BP<sub>ND</sub>  $\times$  group and age  $\times$  group had a  $P \leq 0.01$ ) (model for ventral

striatum:  $R^2 = 0.28$ ;  $F_{4,36} = 3.53$ ;  $P = 0.02$ ; ventral D2R BP<sub>ND</sub>  $\times$  group interaction had a  $P = 0.03$ ; age  $\times$  group interaction had a  $P = 0.06$ ). There was a negative linear relationship between age and sucrose preferences and between both dorsal and ventral D2R BP<sub>ND</sub> and sucrose preferences in subjects without obesity; age and dorsal striatal D2R BP<sub>ND</sub> accounted for 52% (Fig. 2A and B) and age and ventral striatal D2R BP<sub>ND</sub> accounted for 40% (Fig. 3A and B) of the individual differences in sucrose preferences. These associations remained significant when controlling for race and sex in a hierarchical regression model. In contrast, there were no significant relationships between age and sucrose preferences (Figs. 2C and 3C) or between D2R BP<sub>ND</sub> and sucrose preferences (Figs. 2D and 3D) in subjects with obesity.


## DISCUSSION

The primary findings of this study are that subjects with obesity lack the normal associations between age and sucrose preferences and between striatal D2R BP<sub>ND</sub> and sucrose preferences. Additionally, age-related decline in D2R BP<sub>ND</sub> is not directly related to the age-related decline in sucrose preferences. Instead, lower sucrose preference in subjects without obesity related both to increased striatal D2R BP<sub>ND</sub> and older age, explaining up to 52% of individual variance in sucrose preference.

Our findings in subjects without obesity contribute to the understanding of the important role of dopamine in feeding behavior and food reward (20,21). Previous neuroimaging studies in healthy lean people report that dopamine release in dorsal striatum correlates positively with subjective ratings of wanting a food (20) or pleasure experienced from eating a favorite food (21). Here we found that lower striatal D2R BP<sub>ND</sub> relates to preferences for more intense sweetness, which supports observations that patients with Parkinson disease, a neurodegenerative disorder characterized by striatal presynaptic dopamine



**Figure 1**—Sucrose preferences (A) and perceived sweetness of increasing sucrose concentrations (0%, 3%, 12%, and 36% w/v) (B) do not differ between subjects without and with obesity. Both groups similarly perceived progressively increasing sweetness intensity with increasing sucrose concentrations ( $P < 0.0001$ ). Data are presented as geometric mean  $\pm$  geometric SE for sucrose preferences and as mean  $\pm$  SEM for sweetness intensity. sqrt gLMS, square root of ratings made on the general label magnitude scale.




**Figure 2**—Preferences for higher sucrose concentrations were related to younger age (A) and lower dorsal striatal D2R BP<sub>ND</sub> (B) in subjects without obesity (open symbols). Together, age and dorsal striatal D2R BP<sub>ND</sub> accounted for 52% of the individual differences in sucrose preferences. In contrast, sucrose preferences did not associate with age (C) or with dorsal striatal D2R BP<sub>ND</sub> (D) in subjects with obesity (closed symbols). Data points are standardized residuals of partial correlations after controlling for D2R BP<sub>ND</sub> in panels A and C and after controlling for age in panels B and D.

deficiency, exhibit greater liking for higher sucrose concentrations (22) and correlates with data from rodent models that show that treatment with dopamine receptor antagonists cause a shift in sweet preferences toward more concentrated sucrose solutions (23). By contrast, sucrose preference did not relate to striatal D2R BP<sub>ND</sub> in subjects with obesity. The mechanism underlying this lack of association in subjects with obesity is unknown. Our data suggest one more type of dopaminergic disruption associated with obesity. However, whether this dopaminergic disruption truly relates to having a BMI  $>30 \text{ kg/m}^2$  or to other characteristics that may be clustered with obesity, such as having a specific type of diet (e.g., high-fat diet) or markers of metabolic syndrome (e.g., insulin resistance or hyperleptinemia), is unknown.

Several research studies tried to determine whether subjects with obesity respond to sweetness differently from subjects without obesity, with conflicting results

(16). The disruption in subjects with obesity of the well-known age-related decline in sucrose preference suggests that age could be an important factor contributing to inconsistency in the literature of sweetness preferences in obesity. That is, differences may be negligible when sucrose preferences are measured in young adulthood, as we found, but may become more apparent in older populations. The age consideration may also apply to D2R differences; however, in this same population, we reported no D2R differences between groups with and without obesity (13).

Our findings in subjects without obesity support the known age-related decline in sweet preferences (7,9,11) but do not help explain the mechanism(s) underlying such a developmental shift. A previous study in 11- to 15-year-old children found that preferences for high versus low sucrose concentrations related to differential amounts of a biomarker of bone growth in urine, suggesting that biochemical changes controlling linear growth mediate developmental changes in sugar preferences (10). The



**Figure 3**—Preferences for higher sucrose concentrations were related to younger age (A) and lower nucleus accumbens (ventral striatum) D2R BP<sub>ND</sub> (B) in subjects without obesity (open symbols). Together, age and ventral striatal D2R BP<sub>ND</sub> accounted for 40% of the individual differences in sucrose preferences. In contrast, sucrose preferences did not associate with age (C) or with ventral striatal D2R BP<sub>ND</sub> (D) in subjects with obesity (closed symbols). Data points are standardized residuals of partial correlations after controlling for D2R BP<sub>ND</sub> in panels A and C and after controlling for age in panels B and D.

strong negative association between age and sucrose preference found in our nonobese subjects, who were between 20 and 40 years old and have presumably completed their linear growth, suggests that some other, as-yet-unknown mechanism contributes to age-related changes in sucrose preferences and that this mechanism is disrupted in obesity.

Our study has several important limitations. It is possible that our results were influenced by differences in race between groups and by the majority of women in both groups, as people of African American descent generally prefer higher concentrations of sucrose than people of Caucasian descent (9) and there could be sex-related variations in taste perception (24). However, the statistical significance of the relationships between any of our outcome measures do not change when using statistical analyses that control for race and sex. Therefore, it is unlikely that race or sex significantly influenced our conclusions. In addition, we cannot exclude the possibility

that gonadal hormone-related variations in taste perception (24) affected our results because we did not measure plasma hormone concentrations and did not control for the menstrual cycle phase. Finally, these results will require replication because of the novel nature of our findings and the small sample on which they are based.

In summary, both striatal D2R BP<sub>ND</sub> and age are negatively correlated with sucrose preferences in subjects without obesity, but these associations are absent in subjects with obesity. Our data suggest that the age-related decline in D2R is not directly linked to the age-related decline in sweetness preferences, suggesting that other, as-yet-unknown mechanisms are involved and that these mechanisms are disrupted in obesity.

**Acknowledgments.** The authors thank Johanna Sonnenschein for technical assistance with the sensory test; Samantha Ranck and Emily Bihun for assistance in study recruitment and data collection; Heather Lugar, Jerrell Rutlin, Jonathan Koller, and Johanna Hartlein for help in scanning participants

and data processing (all of the Washington University School of Medicine in St. Louis); and the study subjects for their participation.

**Funding.** This work was supported by the National Institutes of Health (R01 DK085575, P30 DK056341, and the Clinical and Translational Science Award UL1 TR000448).

**Duality of Interest.** No potential conflicts of interest relevant to this article were reported.

**Author Contributions.** M.Y.P., J.S.P., K.J.B., and T.H. conceived and designed the experiments. M.Y.P., S.A.E., and A.N.B. performed the experiments. M.Y.P., S.A.E., and T.H. analyzed the data. M.Y.P. wrote the manuscript. S.A.E., A.N.B., S.K., S.M.M., J.S.P., K.J.B., and T.H. contributed to the discussion and reviewed and edited the manuscript. S.M.M. contributed reagents, materials, and analysis tools. M.Y.P. and T.H. are the guarantors of this work and, as such, had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

## References

- Volkow ND, Wang GJ, Tomasi D, Baler RD. The addictive dimensionality of obesity. *Biol Psychiatry* 2013;73:811–818
- Stice E, Spoor S, Bohon C, Small DM. Relation between obesity and blunted striatal response to food is moderated by TaqIA A1 allele. *Science* 2008;322:449–452
- Green E, Jacobson A, Haase L, Murphy C. Reduced nucleus accumbens and caudate nucleus activation to a pleasant taste is associated with obesity in older adults. *Brain Res* 2011;1386:109–117
- Ljungberg T, Ungerstedt U. Reinstatement of eating by dopamine agonists in aphagic dopamine denervated rats. *Physiol Behav* 1976;16:277–283
- Szczypka MS, Kwok K, Brot MD, et al. Dopamine production in the caudate putamen restores feeding in dopamine-deficient mice. *Neuron* 2001;30:819–828
- Tellez LA, Han W, Zhang X, et al. Separate circuitries encode the hedonic and nutritional values of sugar. *Nat Neurosci* 2016;19:465–470
- Desor JA, Beauchamp GK. Longitudinal changes in sweet preferences in humans. *Physiol Behav* 1987;39:639–641
- Bertino M, Wehmer F. Dietary influences on the development of sucrose acceptability in rats. *Dev Psychobiol* 1981;14:19–28
- Pepino MY, Mennella JA. Factors contributing to individual differences in sucrose preference. *Chem Senses* 2005;30(Suppl. 1):i319–i320
- Coldwell SE, Oswald TK, Reed DR. A marker of growth differs between adolescents with high vs. low sugar preference. *Physiol Behav* 2009;96:574–580
- Mennella JA, Lukasewycz LD, Griffith JW, Beauchamp GK. Evaluation of the Monell forced-choice, paired-comparison tracking procedure for determining sweet taste preferences across the lifespan. *Chem Senses* 2011;36:345–355
- Antonini A, Leenders KL. Dopamine D2 receptors in normal human brain: effect of age measured by positron emission tomography (PET) and [<sup>11</sup>C]-raclopride. *Ann N Y Acad Sci* 1993;695:81–85
- Eisenstein SA, Antenor-Dorsey JA, Gredysa DM, et al. A comparison of D2 receptor specific binding in obese and normal-weight individuals using PET with (N-[(11)C]methyl)benperidol. *Synapse* 2013;67:748–756
- Eisenstein SA, Bischoff AN, Gredysa DM, et al. Emotional eating phenotype is associated with central dopamine D2 receptor binding independent of body mass index. *Sci Rep* 2015;5:11283
- Eisenstein SA, Gredysa DM, Antenor-Dorsey JA, et al. Insulin, central dopamine D2 receptors, and monetary reward discounting in obesity. *PLoS One* 2015;10:e0133621
- Bartoshuk LM, Duffy VB, Hayes JE, Moskowitz HR, Snyder DJ. Psychophysics of sweet and fat perception in obesity: problems, solutions and new perspectives. *Philos Trans R Soc Lond B Biol Sci* 2006;361:1137–1148
- Pepino MY, Finkbeiner S, Beauchamp GK, Mennella JA. Obese women have lower monosodium glutamate taste sensitivity and prefer higher concentrations than do normal-weight women. *Obesity (Silver Spring)* 2010;18:959–965
- Brix G, Zaers J, Adam LE, et al.; National Electrical Manufacturers Association. Performance evaluation of a whole-body PET scanner using the NEMA protocol. *J Nucl Med* 1997;38:1614–1623
- Fischl B, Salat DH, Busa E, et al. Whole brain segmentation: automated labeling of neuroanatomical structures in the human brain. *Neuron* 2002;33:341–355
- Volkow ND, Wang GJ, Fowler JS, et al. “Nonhedonic” food motivation in humans involves dopamine in the dorsal striatum and methylphenidate amplifies this effect. *Synapse* 2002;44:175–180
- Small DM, Jones-Gotman M, Dagher A. Feeding-induced dopamine release in dorsal striatum correlates with meal pleasantness ratings in healthy human volunteers. *Neuroimage* 2003;19:1709–1715
- Travers JB, Akey LR, Chen SC, Rosen S, Paulson G, Travers SP. Taste preferences in Parkinson's disease patients. *Chem Senses* 1993;18:47–55
- Xenakis S, Sclafani A. The dopaminergic mediation of a sweet reward in normal and VMH hyperphagic rats. *Pharmacol Biochem Behav* 1982;16:293–302
- Gandelman R. Gonadal hormones and sensory function. *Neurosci Biobehav Rev* 1983;7:1–17