



Direct Analysis of Insulin-Specific T Cells Provides New Insights

Eddie A. James

Diabetes 2017;66:2940–2941 | <https://doi.org/10.2337/dbi17-0041>

Type 1 diabetes (T1D) is an autoimmune disease in which destruction of insulin-producing β -cells leads to dependence on exogenous insulin (1). T1D progression involves progressive loss of tolerance to β -cell antigens, but this has been challenging to observe in human disease. Loss of self-tolerance is most evident through development of islet cell antibodies, among which insulin antibodies typically appear first (2). Given strong genetic association of T1D with HLA-DQ8, DQ8-restricted CD4⁺ T cells activated by self-peptides are thought to play a crucial role in disease. Key evidence from the nonobese diabetic (NOD) mouse model supports a primary role for recognition of an epitope within the insulin B chain amino acids 9–23 (InsB_{9–23}) peptide in disease initiation. The majority of CD4⁺ T cells isolated from pancreatic islets of NOD mice were shown to respond to this peptide (3). Furthermore, expression of an insulin transgene with a substitution within this peptide abrogated diabetes development (4). Efforts to directly visualize and study insulin-specific T cells in human subjects have been hampered by the biochemical instability of the HLA-DQ8 protein (5). However, recent reports highlight the development of HLA-DQ8 tetramer reagents and their application to study T1D, affirming that InsB_{11–23} peptide is analogously presented by IA^{g7} and HLA-DQ8 (6,7). In addition, two recent reports documented the presence of InsB_{9–23}-specific T cells within pancreatic infiltrates of cadaveric donors with T1D (8,9). Nevertheless, key attributes of InsB-specific responses have remained unstudied.

In this issue of *Diabetes*, Spanier et al. (10) report a characterization of DQ8/InsB_{10–23}-specific CD4⁺ T cells in subjects with T1D. Their results support the relevance of such T cells and demonstrate an intriguing correlation between their frequency and insulin antibody titers. After first validating the performance of their DQ8 tetramer by staining of T cells from peptide-immunized and control HLA-DQ8 transgenic mice and peptide-stimulated peripheral blood mononuclear cells (PBMCs), Spanier et al. determined

the ex vivo frequency of DQ8/InsB tetramer binding cells within PBMCs obtained from human subjects, detecting more than twofold higher frequencies in subjects with T1D than in control subjects. Although the insulin B chain is present in exogenous insulin, they observed higher frequencies in patients with a disease duration of <5 years. Furthermore, DQ8/InsB_{10–23}-specific T cells were present within 15 days of diagnosis, suggesting that T cells were not merely expanded in patients because of daily insulin injections. Based on direct surface marker analysis, DQ8/InsB_{10–23}-specific T cells were antigen experienced (CD45RO⁺) and exhibited markers indicative of T helper 1 (CXCR3) and T follicular helper (CXCR5) lineages. Finally, frequencies of DQ8/InsB_{10–23}-specific T cells were positively correlated with insulin antibody titers, implying a possible causal relationship.

Spanier et al. (10) observed that DQ8/InsB_{10–23}-specific T cells were detectable in approximately half of the T1D patients tested and were rarely detected in control subjects. Their results affirm disease relevance of these T cells but also bring into question the central role of this epitope in human T1D that had been postulated based on observations in NOD mice. As summarized in Table 1, the evidence supporting the preeminence of DQ8/InsB_{10–23}-specific T cells is somewhat disparate in mouse and man. Although DQ8/InsB_{10–23}-specific T cells were seen at higher frequencies in subjects with T1D, observed frequencies were 20-fold lower than DQ8/GAD65_{250–266}-specific T cell frequencies that were observed in another study (6). Spanier et al. did not characterize the cytokine profiles of DQ8/InsB_{10–23}-specific T cells. However, their observation that a significant proportion were CXCR3⁺ memory T cells is in agreement with a prior study that demonstrated that DQ8/InsB_{10–23}-specific T cell clones isolated from subjects with T1D predominantly produced IFN- γ , IL-2, and TNF- α (7).

Despite their novel observations, Spanier et al. (10) leave some compelling questions unanswered. In their in vitro tetramer studies, Yang et al. (7) noted that subjects who

Table 1—Insulin autoimmunity in mouse and man

| | Murine diabetes (reference) | Human diabetes (reference) |
|---|--------------------------------|----------------------------------|
| Insulin Ab appear early in disease | Yes (17) | Yes (2) |
| InsB ₉₋₂₃ has low affinity binding register | Yes (12) | Yes (7) |
| InsB ₉₋₂₃ -specific T cells present in islets | Yes (3) | Yes (8,9) |
| InsB ₉₋₂₃ -specific T cells predominant | Yes (3) | No (10) |
| InsB ₉₋₂₃ linked to insulin Ab | Yes (18) | Yes (10) |
| Insulin-specific T cells increase with duration | Unknown | Yes (10) |

Ab, antibody.

had detectable responses to InsB₁₁₋₂₃ were highly enriched for the susceptible VNTR genotype (7). Given the influence of this variant on levels of thymic insulin expression and T cell selection (11), it would be of interest to know whether the susceptible genotype is correlated with higher frequencies of DQ8/InsB₁₀₋₂₃-specific T cells. Recent studies verify that InsB₁₀₋₂₃ is presented by IA^{g7} and DQ8 in an unconventional weak binding register (7,12) such that truncation of the peptide results in an improved ligand. Given recent observations that hybrid insulin peptides are important antigens relevant in human disease (8,13), it could be speculated that a hybrid peptide could generate a more potent agonist. However, rigorous experimental work would be needed to corroborate that idea. Although interesting, the significance of the observed correlation between DQ8/InsB₁₀₋₂₃-specific T cells and higher titers of insulin autoantibodies remains unclear. In particular, it is not discernable whether insulin antibody titer was measured from serum samples drawn at the same time as PBMCs that were subjected to tetramer analysis or whether antibody data were obtained closer to onset. To date, relatively few studies have correlated autoreactive T cell and antibody responses. McLaughlin et al. (14) observed modest correlations between T cell responses to IA-2₈₄₁₋₈₆₀ and IA-2 juxtamembrane domain antibodies. Acevedo-Calado et al. (15) identified T cell and antibody responses to newly described antigenic determinants within the amino terminus of IA-2. However, responses were not assessed in the same subjects. Consequently, the new data provided by Spanier et al. (10) represents a useful contribution to the field.

Although confirming key findings in a larger patient cohort and/or interrogating a longitudinal sample set would be advisable, these results have the potential to refine our understanding of the contribution and role of DQ8/InsB₁₀₋₂₃-specific T cells in human disease. It should be noted that another recent study (16) used HLA-DQ8 tetramers to explore the potential of inducing HLA-DQ8-restricted insulin-specific Foxp3⁺ regulatory T cells. Their findings indicated that DQ8/insulin-specific regulatory T cells were associated

with delayed progression, suggesting that the DQ8/InsB₁₀₋₂₃ epitope would be applicable to elicit antigen-specific tolerance. Clearly, DQ8/InsB₁₀₋₂₃ tetramer assays will be crucial to facilitate such studies and to address remaining questions about the role of InsB-specific responses in T1D.

Duality of Interest. No potential conflicts of interest relevant to this article were reported.

References

- Chiang JL, Kirkman MS, Laffel LMB, Peters AL; Type 1 Diabetes Sourcebook Authors. Type 1 diabetes through the life span: a position statement of the American Diabetes Association. *Diabetes Care* 2014;37:2034–2054
- Yu L, Dong F, Miao D, Fouts AR, Wenzlau JM, Steck AK. Proinsulin/insulin autoantibodies measured with electrochemiluminescent assay are the earliest indicator of prediabetic islet autoimmunity. *Diabetes Care* 2013;36:2266–2270
- Wegmann DR, Norbury-Glaser M, Daniel D. Insulin-specific T cells are a predominant component of islet infiltrates in pre-diabetic NOD mice. *Eur J Immunol* 1994;24:1853–1857
- Nakayama M, Abiru N, Moriyama H, et al. Prime role for an insulin epitope in the development of type 1 diabetes in NOD mice. *Nature* 2005;435:220–223
- Reizis B, Altmann DM, Cohen IR. Biochemical characterization of the human diabetes-associated HLA-DQ8 allelic product: similarity to the major histocompatibility complex class II I-A(g)7 protein of non-obese diabetic mice. *Eur J Immunol* 1997;27:2478–2483
- Chow IT, Yang J, Gates TJ, et al. Assessment of CD4⁺ T cell responses to glutamic acid decarboxylase 65 using DQ8 tetramers reveals a pathogenic role of GAD65 121-140 and GAD65 250-266 in T1D development. *PLoS One* 2014;9:e112882
- Yang J, Chow IT, Sosinowski T, et al. Autoreactive T cells specific for insulin B:11-23 recognize a low-affinity peptide register in human subjects with autoimmune diabetes. *Proc Natl Acad Sci U S A* 2014;111:14840–14845
- Babon JA, DeNicola ME, Blodgett DM, et al. Analysis of self-antigen specificity of islet-infiltrating T cells from human donors with type 1 diabetes. *Nat Med* 2016;22:1482–1487
- Michels AW, Landry LG, McDaniel KA, et al. Islet-derived CD4 T cells targeting proinsulin in human autoimmune diabetes. *Diabetes* 2017;66:722–734
- Spanier JA, Sahli NL, Wilson JC, et al. Increased effector memory insulin-specific CD4⁺ T cells correlate with insulin autoantibodies in patients with recent-onset type 1 diabetes. *Diabetes* 2017;66:3051–3060
- Durinovic-Belló I, Wu RP, Gersuk VH, Sanda S, Shilling HG, Nepom GT. Insulin gene VNTR genotype associates with frequency and phenotype of the autoimmune response to proinsulin. *Genes Immun* 2010;11:188–193
- Stadinski BD, Zhang L, Crawford F, Marrack P, Eisenbarth GS, Kappler JW. Diabetogenic T cells recognize insulin bound to IAg7 in an unexpected, weakly binding register. *Proc Natl Acad Sci U S A* 2010;107:10978–10983
- Delong T, Wiles TA, Baker RL, et al. Pathogenic CD4 T cells in type 1 diabetes recognize epitopes formed by peptide fusion. *Science* 2016;351:711–714
- McLaughlin KA, Gulati K, Richardson CC, et al. HLA-DR4-associated T- and cell responses to specific determinants on the IA-2 autoantigen in type 1 diabetes. *J Immunol* 2014;193:4448–4456
- Acevedo-Calado M, James EA, Morran MP, et al. Identification of unique antigenic determinants in the amino terminus of IA-2 (ICA512) in childhood and adult autoimmune diabetes: new biomarker development. *Diabetes Care* 2017;40:561–568
- Serr I, Fürst RW, Achenbach P, et al. Type 1 diabetes vaccine candidates promote human Foxp3(+)Treg induction in humanized mice. *Nat Commun* 2016;7:10991
- Yu L, Robles DT, Abiru N, et al. Early expression of antiinsulin autoantibodies of humans and the NOD mouse: evidence for early determination of subsequent diabetes. *Proc Natl Acad Sci U S A* 2000;97:1701–1706
- Abiru N, Maniatis AK, Yu L, et al. Peptide and major histocompatibility complex-specific breaking of humoral tolerance to native insulin with the B9-23 peptide in diabetes-prone and normal mice. *Diabetes* 2001;50:1274–1281