

# Synthetic Peroxisome Proliferator-Activated Receptor- $\gamma$ Agonist, Rosiglitazone, Increases Plasma Levels of Adiponectin in Type 2 Diabetic Patients

WEI-SHIUNG YANG, MD, PHD<sup>1,2</sup>  
 CHI-YUAN JENG, MD, PHD<sup>3</sup>  
 TA-JEN WU, MD, PHD<sup>4</sup>  
 SACHIYO TANAKA, MD, PHD<sup>5</sup>  
 TOHRU FUNAHASHI, MD, PHD<sup>5</sup>

YUJI MATSUZAWA, MD, PHD<sup>5</sup>  
 JAO-PING WANG, BS<sup>1</sup>  
 CHI-LING CHEN, PHD<sup>6</sup>  
 TONG-YUAN TAI, MD, PHD<sup>1</sup>  
 LEE-MING CHUANG, MD, PHD<sup>1,2</sup>

**OBJECTIVE** — Adiponectin, a plasma protein exclusively synthesized and secreted by adipose tissue, has recently been shown to have anti-inflammatory, antiatherogenic properties in vitro and beneficial metabolic effects in animals. Lower plasma levels of adiponectin have been documented in human subjects with metabolic syndrome and coronary artery disease. We investigated whether the level of this putative protective adipocytokine could be increased by treatment with a peroxisome proliferator-activated receptor- $\gamma$  (PPAR- $\gamma$ ) agonist in diabetic patients.

**RESEARCH DESIGN AND METHODS** — Type 2 diabetic patients (30 in the treatment group and 34 in the placebo group) were recruited for a randomized double-blind placebo-controlled trial for 6 months with the PPAR- $\gamma$  agonist rosiglitazone. Blood samples were collected and metabolic variables and adiponectin levels were determined in all patients before initiation of the study.

**RESULTS** — In the rosiglitazone group, mean plasma adiponectin level was increased by more than twofold ( $P < 0.0005$ ), whereas no change was observed in the placebo group. Multivariate linear regression analysis showed that whether rosiglitazone was used was the single variable significantly related to the changes of plasma adiponectin. The amount of variance in changes of plasma adiponectin level explained by the treatment was  $\sim 24\%$  ( $r^2 = 0.24$ ) after adjusting for age, sex, and changes in fasting plasma glucose, HbA<sub>1c</sub>, insulin resistance index, and BMI.

**CONCLUSIONS** — Rosiglitazone increases plasma adiponectin levels in type 2 diabetic subjects. Whether this may contribute to the antihyperglycemic and putative antiatherogenic benefits of PPAR- $\gamma$  agonists in type 2 diabetic patients warrants further investigation.

*Diabetes Care* 25:376–380, 2002

From the <sup>1</sup>Department of Internal Medicine, National Taiwan University Hospital, Taipei, Taiwan; the <sup>2</sup>Graduate Institute of Clinical Medicine, College of Medicine, National Taiwan University, Taipei, Taiwan; the <sup>3</sup>Department of Internal Medicine, Tri-Service General Hospital, Taipei, Taiwan; the <sup>4</sup>Department of Internal Medicine, National Cheng-Kung University Hospital, Tainan, Taiwan; the <sup>5</sup>Department of Internal Medicine and Molecular Science, Graduate School of Medicine, Osaka University, Osaka, Japan; and the <sup>6</sup>Institute of Epidemiology, College of Public Health, National Taiwan University, Taipei, Taiwan.

Address correspondence and reprint requests to Lee-Ming Chuang, MD, PhD, Department of Internal Medicine, National Taiwan University Hospital, 7 Chung-Shan South Road, Taipei, Taiwan. E-mail: leeming@ha.mc.ntu.edu.tw.

Received for publication 20 June 2001 and accepted in revised form 24 October 2001.

L.M.C. has received funding from Smith-Kline Beecham.

Abbreviations: HOMA, homeostasis model assessment; TNF, tumor necrosis factor; PPAR- $\gamma$ , peroxisome proliferator-activated receptor- $\gamma$ .

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

Adiponectin is an adipose tissue-specific plasma glycoprotein with various homology to collagen, complement proteins, and hibernation-associated proteins (1–4). The human adiponectin (also called apM1 or GBP-28) gene and its mouse homolog (adipoQ or ACRP30) have been identified (1–4). This protein is strictly expressed in and secreted by adipocytes and adipose tissue (1–4). Prokaryote-expressed recombinant adiponectin was capable of reducing tumor necrosis factor (TNF)- $\alpha$ -induced monocyte adhesion, nuclear factor- $\kappa$ B signaling, and expression of intracellular adhesion molecule-1, endothelial cell adhesion molecule-1 and E-selectin in endothelial cells in vitro (5,6). It also attenuates production of TNF- $\alpha$  in lipopolysaccharide-treated macrophages (7). Recently, it was also demonstrated to reduce cholesterol ester accumulation and class A scavenger receptor gene expression in cultured human monocyte-derived macrophages (8). These studies indicate that adiponectin is an adipocyte and adipose tissue-specific anti-inflammatory and antiatherogenic molecule and may be a bridge between obesity and atherosclerosis. Furthermore, adiponectin was detectable only in the walls of catheter-injured vessels but not in the intact vascular walls in an animal model and in humans (8,9).

Lower plasma levels of adiponectin have been documented in human subjects with obesity, type 2 diabetes, or coronary artery disease (5,10,11). Plasma levels of adiponectin were shown to correlate negatively with plasma glucose, insulin, triglyceride levels, and BMI but positively with plasma levels of HDL cholesterol (11). Plasma adiponectin levels were also demonstrated to correlate positively with insulin-stimulated glucose disposal measured by hyperinsulinemic-euglycemic clamp (12). Taken together,

**Table 1—Baseline characteristics of 30 patients treated with rosiglitazone and 34 patients treated with placebo**

	Rosiglitazone	Placebo
Age (years)	58.9 ± 9.4	57.8 ± 8.9
Sex (M/F)	13/17	13/21
Body weight (kg)	64.9 ± 11.8	65.3 ± 11.2
BMI (kg/m <sup>2</sup> )	25.76 ± 2.87	25.84 ± 3.50
Systolic blood pressure (mmHg)	127.8 ± 13.8	134.8 ± 18.8
Diastolic blood pressure (mmHg)	79.5 ± 10.5	80.2 ± 10.0
Fasting plasma glucose (mmol/l)	10.7 ± 2.7	11.0 ± 2.9
Fasting plasma insulin (pmol/l)	69.6 ± 33.3	75.7 ± 45.2
Insulin resistance (by HOMA)	1.08 ± 0.61	1.16 ± 0.63
HbA <sub>1c</sub> (%)	9.5 ± 1.1	9.7 ± 1.4

Data are means ± SD.

these suggest that low plasma adiponectin according to levels may be a novel biomarker of insulin resistance syndrome.

Recent animal studies indicate that adiponectin has a wide array of metabolic effects. Injection of recombinant adiponectin in experimental animals was shown to reduce plasma fatty acid and glucose levels, increase fatty acid  $\beta$ -oxidation and decrease triglyceride content in skeletal muscles, improve insulin sensitivity, and reduce body weight (13–15). These studies further support that a low plasma adiponectin level may play a role in the pathophysiology of type 2 diabetes.

The expression of adiponectin is developmentally regulated and is activated by day 4 during adipocyte differentiation in cultured 3T3-L1 cells (1). Furthermore, we have tested whether proliferator-activated receptor- $\gamma$  (PPAR- $\gamma$ ), the master regulator of adipocyte differentiation, might upregulate its expression in fully differentiated 3T3-L1 adipocytes. We showed that the steady-state mRNA of adiponectin in differentiated 3T3-L1 adipocytes was increased by administration of rosiglitazone for 24 h (16). Rosiglitazone is a synthetic PPAR- $\gamma$  agonist and is now widely used to treat patients with type 2 diabetes through enhancing insulin sensitivity (17). Therefore, we asked in this study whether plasma adiponectin levels might increase in type 2 diabetes patients treated with rosiglitazone.

## RESEARCH DESIGN AND METHODS

In a phase IIIb, double-blind, placebo-controlled, parallel-group comparative study of rosiglitazone (BRL 49653C) and concurrent sulfonylurea

therapy, patients who met the inclusion criteria (aged 30–80 years, type 2 diabetic according to diagnostic criteria of the World Health Organization, fasting plasma glucose 7–15 mmol/l, and HbA<sub>1c</sub> >7.5%) and exclusion criteria (other severe medical problems and severe microvascular complications requiring immediate medical attention) and who had been stable on sulfonylurea therapy for at least 2 months before the screening visit were recruited for the study. During the screening visit, patients entered a single-blind placebo/sulfonylurea run-in period for 4 weeks to establish baseline characteristics. Patients were then randomized for the double-blind phase. Each patient received two tablets of either rosiglitazone (Avandia, 2 mg/tablet) or placebo in a dosing regimen of one tablet twice daily for 6 months. A total of eight follow-up visits were arranged in these 6 months. After evaluation of the safety and efficacy of rosiglitazone (manuscript in preparation), 30 patients in the rosiglitazone group and 34 patients in the placebo group (of 38 and 39 recruited, respectively) who were observed for 6 months were included in this follow-up study. The average age at onset of diabetes was ~50 years. The baseline characteristics of these subjects are shown in Table 1. This protocol has been approved by the Institutional Review Board and Department of Health of Taiwan. Written informed consent was obtained from each participant.

## Anthropometric and biochemical measurements

Body weight, height, blood pressure, and heart rate were measured at the time points specified. The concentrations of

plasma glucose, total cholesterol, and triglyceride were measured in fasting blood samples by an autoanalyzer (Hitachi 7250 Special; Hitachi, Tokyo, Japan). Serum insulin levels were determined by a microparticle enzyme immunoassay using the AxSYM system (Abbott Laboratories, Dainabot, Tokyo, Japan). HbA<sub>1c</sub> was measured with DCA2000 (Bayer Sankyo, Tokyo, Japan). The homeostasis model assessment was applied to estimate the degree of insulin resistance (homeostasis model assessment [HOMA]-IR = Insulin/22.5e<sup>-ln(Glucose)</sup>) and  $\beta$ -cell function (HOMA  $\beta$  = 20 × Insulin/[Glucose - 3.5]), where insulin is expressed in  $\mu$ U/ml and glucose is expressed in mmol/l (18). Plasma levels of adiponectin were determined by an enzyme-linked immunosorbent assay system as described previously (10). All data except those for adiponectin were presented in SI units, because the molecular weight of adiponectin has not been precisely determined.

## Statistical analyses

Data were presented as means and SDs. Statistical analyses, including Student's *t* test and multivariate linear regression analysis, were performed using Statistical Analysis System software (Edition 6.12; SAS Institute, Cary, NC). Differences in clinical characteristics between those treated with rosiglitazone and those treated with placebo were tested by Student's *t* test. Several multivariate linear regression models were performed that included age, sex, changes in fasting plasma glucose level, HbA<sub>1c</sub>, insulin resistance index by HOMA, BMI, and treatment status as independent variables and changes in plasma adiponectin levels as dependent variables. The amount of variance explained by treatment status was indicated by the *r*<sup>2</sup> or partial *r*<sup>2</sup> in the models.

**RESULTS** — At baseline, there was no difference in general characteristics between rosiglitazone and placebo groups (Table 1). Three months after treatment, there were significant differences in mean plasma adiponectin levels, fasting plasma glucose level, and HbA<sub>1c</sub> between the groups (data not shown). After 6 months of treatment, the subjects in the rosiglitazone group had significant weight gain as well as better glycemic control compared with those in the placebo group (Table 2).

**Table 2—Changes of characteristics from baseline to 6 months after treatment with either rosiglitazone in 30 subjects or placebo in 34 subjects**

	Rosiglitazone	Placebo
Body weight	3.0 ± 2.4 <sup>§</sup>	-0.4 ± 1.9
BMI	1.20 ± 1.00 <sup>§</sup>	-0.18 ± 0.79
Systolic blood pressure	-0.3 ± 15.7*	-8.1 ± 16.3
Diastolic blood pressure	-0.4 ± 8.0	-1.1 ± 7.4
Fasting plasma glucose	-10.6 ± 41.0†	17.8 ± 58.5
Fasting plasma insulin	-1.07 ± 4.87	-1.16 ± 4.72
Insulin resistance	-0.19 ± 0.47*	0.04 ± 0.60
Insulin secretion	-1.20 ± 12.80	-3.11 ± 8.31
HbA <sub>1c</sub>	-0.7 ± 1.0‡	0.4 ± 1.3

Data are means ± SD. \* $P < 0.1$ ; † $P < 0.05$ ; ‡ $P < 0.005$ ; § $P < 0.0005$  compared with the placebo group by student's *t* test.

Insulin resistance was reduced in the rosiglitazone group after 6 months. However, the change was not significant when compared with that of the placebo group.

The increase of plasma adiponectin was more than twofold in the rosiglitazone group, whereas in the placebo group, there was virtually no change of plasma adiponectin (Fig. 1). Treatment with rosiglitazone for 6 months did not further increase the plasma adiponectin level from that measured at 3 months, suggesting that the maximal effect of rosiglitazone on adiponectin levels had been attained by the 3rd month.

Using the changes in adiponectin as the dependent variable in multivariate linear regression models, we found that changes in HbA<sub>1c</sub>, BMI, and treatment status (use of rosiglitazone or placebo), but not fasting plasma glucose level and insulin resistance, as independent variables were significantly related to adiponectin after adjusting age and sex (data not shown). After further adjustment of treatment status in each analysis, none of these variables remained significantly related to adiponectin (data not shown). Stepwise multivariate linear regression analyses showed that treatment status was the single significant variable related to the changes in plasma adiponectin (Table 3), although both the changes in insulin resistance and BMI were significantly related to that of plasma adiponectin before treatment status was adjusted. Without adjusting other variables, treatment status alone contributed to ~38% of variance in the changes of plasma adiponectin ( $r^2 = 0.38$ ). After adjusting the other variables in model 5, treatment status was attributed to ~24% of variance ( $r^2 = 0.24$ ),


whereas the  $r^2$  of the model was 0.44. These results suggest that the increase in plasma adiponectin level may primarily be the result of rosiglitazone treatment.

**CONCLUSIONS**— Adiponectin is an adipose tissue-derived plasma protein, the functions of which have not been clear until recent years. In vitro studies have shown antiatherogenic, anti-inflammatory, and apoptotic effects of adiponectin (5–8). However, strong evidence linking adiponectin to these biological functions in vivo is still deficient. In many human studies, low plasma lev-

els of adiponectin were shown to be associated with insulin resistance and various clinical manifestations of metabolic syndrome (5,10–12). Whether low plasma adiponectin levels may directly contribute to accelerated atherosclerosis in patients with metabolic syndrome merits intensive investigation.

Recently, adiponectin also has been shown to have various metabolic effects. It was reported that the decrease in plasma adiponectin paralleled the development of insulin resistance and diabetes among monkeys in a longitudinal observation study (19). Plasma level of adiponectin was also related to the direct measurement of insulin sensitivity by clamp studies in humans (12). Furthermore, injection of recombinant adiponectin in experimental animals was shown to increase fatty acid oxidation in muscle, improve insulin sensitivity, and decrease hepatic glucose output (13–15). This indicates that low plasma adiponectin may contribute to the pathophysiology of insulin resistance among patients with type 2 diabetes.

The mechanism of adiponectin in improving insulin sensitivity has been elucidated, at least in part. Recombinant adiponectin injection in experimental animal models showed a decrease in plasma fatty acid levels (13,14). Fatty acid  $\beta$ -



**Figure 1—Mean and SEM of plasma adiponectin levels at baseline and at 3 and 6 months after treatment. White bars, placebo group; black bars, rosiglitazone group. \*  $P < 0.05$ ; \*\*  $P < 0.01$ .**

**Table 3—Multivariate linear regression models showing intercepts and regression coefficients  $\pm$  SEM, using changes of adiponectin as the dependent variable, with age, sex, changes of fasting plasma glucose, HbA<sub>1c</sub>, insulin resistance, and BMI, and treatment status as independent variables**

Independent variables	Change in adiponectin				
	Model 1	Model 2	Model 3	Model 4	Model 5
Intercept	-7.7 $\pm$ 4.7	-6.3 $\pm$ 4.7	-6.1 $\pm$ 4.7	-5.7 $\pm$ 4.1	-8.9 $\pm$ 3.8
Age	0.15 $\pm$ 0.08	0.13 $\pm$ 0.08	0.12 $\pm$ 0.08	0.12 $\pm$ 0.07	0.12 $\pm$ 0.06
Sex	1.8 $\pm$ 1.5	1.7 $\pm$ 1.5	1.7 $\pm$ 1.5	1.1 $\pm$ 1.3	1.5 $\pm$ 1.2
Change in fasting plasma glucose	-0.36 $\pm$ 0.18	-0.18 $\pm$ 0.18	0.05 $\pm$ 0.31	0.14 $\pm$ 0.27	0.18 $\pm$ 0.18
Change in HbA <sub>1c</sub>		-1.12 $\pm$ 0.60	-1.22 $\pm$ 0.60*	0.10 $\pm$ 0.62	0.24 $\pm$ 0.56
Change in insulin resistance			-2.17 $\pm$ 1.53	-2.84 $\pm$ 1.36*	-1.7 $\pm$ 1.26
Change in BMI				2.72 $\pm$ 0.67†	1.32 $\pm$ 0.72
Treatment					5.5 $\pm$ 1.5†

\* $P < 0.05$ ; † $P < 0.001$ .

oxidation in skeletal muscle increased in these animals, probably secondary to enhanced expression of genes involved in  $\beta$ -oxidation and energy dissipation, such as acyl-CoA oxidase and uncoupling protein-2 (14). As a result, adiponectin also decreased triglyceride content in muscle. Both the increased fatty acid levels in plasma and increased triglyceride content in muscle have been shown to cause insulin resistance (20). In addition, insulin-stimulated tyrosine phosphorylation of signaling molecules, including insulin receptor, insulin receptor substrate-1, and Akt in skeletal muscle, was also enhanced by adiponectin (14).

In view of its potential beneficial effects, any measures that could increase plasma adiponectin levels would likely have some clinical significance. We recently also showed that weight reduction with gastric partition surgery significantly increased plasma adiponectin levels and insulin sensitivity among severely obese individuals (21). Whether certain beneficial metabolic effects, such as improvement in insulin sensitivity by weight reduction or by treatment with PPAR- $\gamma$  agonists, were at least partially mediated by adiponectin remains to be investigated.

Although we did not investigate the expression of adiponectin in adipose tissue in these subjects, it should be plausible to speculate that the mechanism of increased plasma adiponectin by rosiglitazone treatment was secondary to increased transcription of adiponectin gene by activating the transcription factor, PPAR- $\gamma$ . We have also shown, in cultured mouse adipocytes, that rosiglitazone increased the steady-state mRNA of adi-

ponectin (16). The proximal promoter region (1 kb) of the human adiponectin gene contains only a half-site of PPAR- $\gamma$  response element (AGGTCA between -610 and -605 relative to ATG start codon) (22,23). Its proximal promoter also contains several regulatory elements commonly observed in the promoters of genes expressed in adipose tissue. These cis-elements include C/EBPs, SREBP, E-box, and GATA-1 (22,23). However, it cannot be excluded that rosiglitazone may enhance the stability of adiponectin mRNA, as well as the synthesis, stability, and secretion of adiponectin protein.

In this study, we observed an increase in plasma adiponectin along with weight gain after rosiglitazone treatment. This may seem contradictory to the previously reported negative correlation between plasma adiponectin level and body weight. It is plausible that activation of PPAR- $\gamma$  by rosiglitazone may, on one hand, promote body weight gain by increasing adipocyte differentiation and the number of small adipocytes as previously shown (24) and, on the other hand, enhance adiponectin gene transcription.

In conclusion, this study clearly demonstrates that treatment with rosiglitazone in type 2 diabetic patients increased plasma adiponectin levels. This effect may potentially protect diabetic patients from macrovascular complications and may improve their insulin sensitivity and glycemic control. Therefore, further studies on its clinical and biological relevance are warranted.

**Acknowledgments**—This study was supported, in part, by Smith-Kline Beecham Phar-

maceuticals and a grant from the Department of Education (89-B-FA01-1-4) of the R.O.C.

We thank Chia-Ling Chao for technical assistance.

## References

- Hu E, Liang P, Spiegelman BM: AdipoQ is a novel adipocyte-specific gene dysregulated in obesity. *J Biol Chem* 271:10697–10703, 1996
- Scherer PE, Williams S, Fogliano M, Baldini G, Lodish HF: A novel serum protein similar to C1q, produced exclusively in adipocytes. *J Biol Chem* 270:26746–26749, 1995
- Maeda K, Okubo K, Shimomura I, Funahashi T, Matsuzawa Y, Matsubara K: cDNA cloning and expression of a novel adipose specific collagen-like factor, apM1. *Biochem Biophys Res Commun* 221:286–289, 1996
- Nakano Y, Tobe T, Choi-Miura NH, Mazda T, Tomita M: Isolation and characterization of GBP28, a novel gelatin-binding protein purified from human plasma. *J Biochem* 120:803–812, 1996
- Ouchi N, Kihara S, Arita Y, Maeda K, Kuriyama H, Okamoto Y, Hotta K, Nishida M, Takahashi M, Nakamura T, Yamashita S, Funahashi T, Matsuzawa Y: Novel modulator for endothelial adhesion molecules: adipocyte-derived plasma protein adiponectin. *Circulation* 100:2473–2476, 1999
- Ouchi N, Kihara S, Arita Y, Okamoto Y, Maeda K, Kuriyama H, Hotta K, Nishida M, Takahashi M, Muraguchi M, Ohmoto Y, Nakamura T, Yamashita S, Funahashi T, Matsuzawa Y: Adiponectin, an adipocyte-derived plasma protein, inhibits endothelial NF- $\kappa$ B signaling through a cAMP-dependent pathway. *Circulation* 102:1296–1301, 2000
- Yokota T, Oritani K, Takahashi I, Ishikawa J, Matsuyama A, Ouchi N, Kihara S,

- Funahashi T, Tenner AJ, Tomiyama Y, Matsuzawa Y: Adiponectin, a new member of the family of soluble defense collagens, negatively regulates the growth of myelomonocytic progenitors and the functions of macrophages. *Blood* 96: 1723–1732, 2000
8. Ouchi N, Kihara S, Arita Y, Nishida M, Matsuyama A, Okamoto Y, Ishigami M, Kuriyama H, Kishida K, Nishizawa H, Hotta K, Muraguchi M, Ohmoto Y, Yamashita S, Funahashi T, Matsuzawa Y: Adipocyte-derived plasma protein, adiponectin, suppresses lipid accumulation and class A scavenger receptor expression in human monocyte-derived macrophages. *Circulation* 103:1057–1063, 2001
  9. Okamoto Y, Arita Y, Nishida M, Muraguchi M, Ouchi N, Takahashi M, Igura T, Inui Y, Kihara S, Nakamura T, Yamashita S, Miyagawa J, Funahashi T, Matsuzawa Y: An adipocyte-derived plasma protein, adiponectin, adheres to injured vascular walls. *Horm Metab Res* 32:47–50, 2000
  10. Arita Y, Kihara S, Ouchi N, Takahashi M, Maeda K, Miyagawa J, Hotta K, Shimomura I, Nakamura T, Miyaoka K, Kuriyama H, Nishida M, Yamashita S, Okubo K, Matsubara K, Muraguchi M, Ohmoto Y, Funahashi T, Matsuzawa Y: Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity. *Biochem Biophys Res Commun* 257:79–83, 1999
  11. Hotta K, Funahashi T, Arita Y, Takahashi M, Matsuda M, Okamoto Y, Iwahashi H, Kuriyama H, Ouchi N, Maeda K, Nishida M, Kihara S, Sakai N, Nakajima T, Hasegawa K, Muraguchi M, Ohmoto Y, Nakamura T, Yamashita S, Hanafusa T, Matsuzawa Y: Plasma concentrations of a novel, adipose-specific protein, adiponectin, in type 2 diabetic patients. *Arterioscler Thromb Vasc Biol* 20:1595–1599, 2000
  12. Weyer C, Funahashi T, Tanaka S, Hotta K, Matsuzawa Y, Pratley RE, Tatarani PA: Hypoadiponectinemia in obesity and type 2 diabetes: close association with insulin resistance and hyperinsulinemia. *J Clin Endocrinol Metab* 86:1930–1935, 2001
  13. Fruebis J, Tsao T-S, Javorschi S, Ebbets-Reed D, Erickson MRS, Yen FT, Bihain BE, Lodish HF: Proteolytic cleavage product of 30-kDa adipocyte complement-related protein increases fatty acid oxidation in muscle and causes weight loss in mice. *Proc Natl Acad Sci U S A* 98: 2005–2010, 2001
  14. Yamauchi T, Kamon J, Waki H, Terauchi Y, Kubota N, Hara K, Mori Y, Ide T, Murakami K, Tsuboyama-Kasaoka N, Ezaki O, Akanuma Y, Gavrilova O, Vinson C, Reitman ML, Kagechika H, Shudo K, Yoda M, Nakano Y, Tobe K, Nagai R, Kimura S, Tomita M, Froguel P, Kadowaki T: The fat-derived hormone adiponectin reverses insulin resistance associated with both lipoatrophy and obesity. *Nat Med* 7:941–946, 2001
  15. Berg AH, Combs TP, Du X, Brownlee M, Scherer P: The adipocyte-secreted protein Acrp30 enhances hepatic insulin action. *Nat Med* 7:947–953, 2001
  16. Lin WS, Chang HM, Tai TY, Chuang LM: Effect of thiazolidinedione on gene expression in NIH3T3–L1 adipocytes (Abstract). *Diabetes* 48 (Suppl. 1):A217, 1999
  17. Whitcomb RW, Saltiel AR: Thiazolidinediones. *Expert Opin Invest Drug* 4:1299–1309, 1995
  18. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC: Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 28:412–419, 1985
  19. Hotta K, Funahashi T, Bodkin NL, Ortmeier HK, Arita Y, Hansen BC, Matsuzawa Y: Circulating concentrations of the adipocyte protein adiponectin are decreased in parallel with reduced insulin sensitivity during the progression to type 2 diabetes in Rhesus monkeys. *Diabetes* 50:1126–1133, 2001
  20. Shulman GI: Cellular mechanism of insulin resistance. *J Clin Invest* 106:171–176, 2000
  21. Yang WS, Lee WJ, Funahashi T, Tanaka S, Matsuzawa Y, Chao CL, Chen CL, Tai TY, Chuang LM: Weight reduction increases plasma levels of an adipose-derived anti-inflammatory protein adiponectin. *J Clin Endocrinol Metab* 86:3815–3819, 2001
  22. Saito K, Tobe T, Minoshima S, Asakawa S, Sumiya J, Yoda M, Nakano Y, Shimizu N, Tomita M: Organization of the gene for gelatin-binding protein (GBP28). *Gene* 229:67–73, 1999
  23. Schäffler A, Langmann T, Palitzsch KD, Schölmerich J, Schmitz G: Identification and characterization of the human adipocyte apM-1 promoter. *Biochim Biophys Acta* 1399:187–197, 1998
  24. Okuno A, Tamemoto H, Tobe K, Ueki K, Mori Y, Iwamoto K, Umehara K, Akanuma Y, Fujiwara T, Horikoshi H, Yazaki Y, Kadowaki T: Troglitazone increases the number of small adipocytes without the change of white adipose tissue mass in obese Zucker rats. *J Clin Invest* 101:1354–1361, 1998