

The Association Between Diabetes Metabolic Control and Drug Adherence in an Indigent Population

JOEL M. SCHECHTMAN, MD, MPH

MOHAN M. NADKARNI, MD

JOHN D. VOSS, MD

OBJECTIVE— Studies of the association between diabetes metabolic control and adherence to drug therapy have yielded conflicting results. Because low socioeconomic and minority populations have poorer diabetes outcomes and greater barriers to adherence, we examined the relationship between adherence and diabetes metabolic control in a large indigent population.

RESEARCH DESIGN AND METHODS— The study population consisted of patients receiving medical care from a university-based internal medicine clinic serving a low-income population in rural central Virginia. The sample comprised 810 patients with type 2 diabetes who received oral diabetes medications from the clinic pharmacy and had at least one HbA_{1c} determination during the study period. Multiple linear regression was used to examine the association of HbA_{1c} level as well as change in HbA_{1c} level with medication adherence, demographic, and clinical characteristics.

RESULTS— Better metabolic control was independently associated with greater medication adherence, increasing age, white (versus African-American) race, and lower intensity of drug therapy. For each 10% increment in drug adherence, HbA_{1c} decreased by 0.16% ($P < 0.0001$). Controlling for other demographic and clinical variables, the mean HbA_{1c} of African-Americans was 0.29% higher than that of whites ($P = 0.04$). Additionally, the intensity of diabetes drug therapy for African-Americans was lower, as was their measured adherence to it. There was no association between metabolic control and gender, income, encounter frequency, frequency of HbA_{1c} testing, or continuity of care.

CONCLUSIONS— Adherence to medication regimens for type 2 diabetes is strongly associated with metabolic control in an indigent population; African-Americans have lower adherence and worse metabolic control. Greater efforts are clearly needed to facilitate diabetes self-management behaviors of low-income populations and foster culturally sensitive and appropriate care for minority groups.

Diabetes Care 25:1015–1021, 2002

Adherence to chronic drug regimens is often suboptimal; lower socioeconomic and minority populations have greater barriers to adherence, which may thwart efforts at improving care and outcomes (1,2). With the ever-expanding arsenal of pharmacotherapy for chronic disease and evidence linking such therapy to better outcomes, aware-

ness of the critical role of adherence to drug therapy has heightened. Because lower socioeconomic and minority groups generally have greater burdens of chronic disease, less vigorous treatment, and poorer disease outcomes (3–7), it is appropriate to focus adherence efforts on such populations.

Diabetes is a chronic condition in

which evidence clearly links improved metabolic control via drug therapy to better outcomes (8–10). Because of this linkage, one might expect that greater adherence to medical regimens would be associated with better metabolic control, both due to a direct effect and possibly as a marker of adherence to other diabetes self-management behaviors. Alternatively, it has been hypothesized that the absence of such an association could be due to prescription of inadequate medication regimens by physicians, failure to accurately measure adherence, or lack of relationship between medication taking behavior and other diabetes self-management behaviors (11,12). In fact, the concept of adherence with respect to diabetes has been considered dysfunctional, to the extent that blindly following a physician's orders may not promote the desired goal of patient empowerment and greater self-management (13).

Studies that have empirically evaluated the association between medication adherence and diabetes metabolic control have yielded conflicting results (14–22). However, most have relied on small patient samples and self-report as the measure of drug adherence and/or did not use HbA_{1c} as the metabolic outcome measure. The present study was therefore undertaken to evaluate the association between drug adherence and metabolic control in a large sample of patients with type 2 diabetes using objective adherence and outcome measures. We studied an indigent population because their greater barriers to adherence, higher diabetes prevalence, and higher rate of adverse diabetes outcomes are important public health concerns.

RESEARCH DESIGN AND METHODS

— The study was conducted at the University of Virginia (UVA) Health System's principal internal medicine primary care practice site. This site serves 13,138 patients, making 37,500 visits per year, and is staffed by 90 resident and faculty physicians. UVA is the largest provider of primary care to indi-

From the Department of Medicine, University of Virginia, Charlottesville, Virginia.

Address correspondence and reprint requests to Joel M. Schechtman, MD, MPH, Department of Medicine, University of Virginia, PO Box 800744, Charlottesville, VA 22908. E-mail: js3dv@virginia.edu.

Received for publication 22 August 2001 and accepted in revised form 5 March 2002.

Abbreviations: UVA, University of Virginia.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

Table 1—Study population description

	N (%)	Mean (SD)
Race		
African-American	346 (41.7)	
Caucasian	483 (58.3)	
Sex		
Male	327 (39.5)	
Female	502 (60.5)	
Age (years)		58.7 (11.5)
Income		
<Federal poverty	503 (60.7)	
100–110% poverty	61 (7.4)	
110–133% poverty	98 (11.8)	
133–167% poverty	71 (8.6)	
167–200% poverty	28 (3.4)	
>200% poverty	68 (8.2)	
HbA _{1c} level (most recent)		8.1% (2.0)
HbA _{1c} decrease		0.52% (2.16)
No. HbA _{1c} levels (20 months)		3.5 (1.6)
No. clinic visits per year		6.6 (5.1)
No. oral agents per patient		
1	403 (48.6)	
2	363 (43.8)	
3	63 (7.6)	
Concomitant insulin	252 (30.4)	
Oral agent adherence		79.7% (21)
Adherence to nondiabetes drugs		71.8% (17)
Visits to single provider (continuity)		73.5% (23)

gent patients in central Virginia and is the only institution within a 45-mile radius that is obliged to care for all patients regardless of ability to pay. Greater than 60% of the patients qualify for discounted care at UVA, using state family income guidelines. UVA uses a sliding fee scale based on income (as verified by federal income tax returns). With respect to prescription drug costs, the patient copayment ranges from 1 to 100% of the UVA drug acquisition cost. Although the patient population served is racially and socioeconomically diverse, 92% of the patients in the current study received a 50–99% subsidy of their prescription drug costs. Considering the high level of subsidy provided, lack of insurance drug coverage, and low income levels, it is unlikely that they had alternative sources of pharmaceuticals. Analysis did not reveal a difference in refill rates between patients with higher and lower income/copayment levels.

The study population was limited to patients taking at least one oral agent (sulfonylureas, metformin, α -glucosidase inhibitors, and thiazolidinediones) for type

2 diabetes. Although we controlled for insulin use, we were unable to determine drug adherence for insulin through use of an administrative data set because dosage instructions may change frequently and may not be well documented. There were 844 patients identified on one or more oral agents for diabetes during the study period of January 2000 through March 2001. The study population was further limited to the 829 patients that identified themselves as either African-American or Caucasian in their UVA registration data.

Adherence to prescribed drug therapy was calculated using prescription-refill data from the UVA pharmacy serving this patient population. The sum of the number of days of therapy dispensed on all but the last prescription in this period was used as the numerator of the adherence statistic. Using dates of the first and last prescriptions during this period as the boundaries, the number of days in the treatment interval served as the denominator. This method for calculation of proportional adherence is similar to that of previous adherence studies using prescription refill data (23–26). A

sample chart audit of 96 prescriptions indicated that the pharmacy database drug dosage and progress note dosage were in agreement 88.5% of the time and that only 3% of medications had been filled by outside pharmacies.

The association of diabetes metabolic control with patient demographic, clinical, and adherence characteristics was modeled using bivariable and multivariable statistical methods. Metabolic control, the dependent variable, was defined as the most recent HbA_{1c} level during the study period (the results were similar using the mean HbA_{1c} level). Improvement in metabolic control was defined as the difference between the first and last HbA_{1c} levels during a 20-month time frame concluding at the end of the study period. Adherence was modeled as the mean adherence for all diabetes drugs taken by each patient. The association of HbA_{1c} with patient sociodemographic variables (age, gender, race, income level) and prescription drug or disease characteristics (number of diabetes medications taken, mean adherence, number of clinic visits, proportion of visits to a single physician, number of HbA_{1c} levels obtained during the study period) was evaluated by simple and multivariable linear regression. Improvement in metabolic control (change in HbA_{1c}) was similarly modeled; the initial HbA_{1c} level added as an independent variable. Associations were considered statistically significant at a two-tailed α of 0.05. Of the 829 eligible study patients, 810 (98%) had at least one HbA_{1c} level and were included in the metabolic control model; 726 (88%) had at least two levels during the study period and were evaluated for improvement in metabolic control.

RESULTS— The study population demographics as well as means and variability of the analysis variables are summarized in Table 1. The mean age was 59 years, 42% were African-American, and 61% were women. A total of 61% of subjects had family income below the federal poverty line and 92% had family income less than twice the poverty level. Slightly more than half of subjects were taking two or more oral diabetic agents and 30% were also using insulin. The patients averaged more than six primary care visits (for all problems) per year and had a mean of 3.5 HbA_{1c} measurements recorded during the past 20 months (in-

Table 2—Bivariable associations with diabetes metabolic control

Independent variable	Dependent variable: most recent HbA _{1c} level			Dependent variable: change in HbA _{1c} level*		
	Parameter estimate	P value	R ²	Parameter estimate	P value	R ²
Race (white = 0)	0.29	0.04	0.5%	0.25	0.05	0.3%
Sex (male = 0)	-0.17	0.24	NA	-0.04	0.74	NA
Age	-0.026	<0.0001	2.3%	-0.01	0.04	0.4%
Income/copay	0.06	0.11	NA	-0.008	0.83	NA
Adherence (0–100%)	-0.019	<0.0001	4.0%	-0.013	<0.0001	1.7%
No. oral agents	0.79	<0.0001	6.1%	0.55	<0.0001	2.6%
Insulin use (0 or 1)	0.56	0.0003	1.6%	0.38	0.007	0.7%
No. HbA _{1c} levels	-0.004	0.93	NA	0.03	0.55	NA
No. encounters	-0.008	0.45	NA	0.006	0.50	NA
Continuity index	-0.09	0.78	NA	0.55	0.13	NA

*The initial HbA_{1c} level was controlled for in this model. A negative parameter estimate indicates HbA_{1c} improved with increment in independent variable.

cluding the 15-month period for which medication adherence was measured). The mean HbA_{1c} level was 8.1%; the average decrease was 0.5% over the study period.

The bivariable associations between HbA_{1c} level and the analysis variables are shown in Table 2. Metabolic control was associated with race, age, adherence, and intensity of drug therapy. For each 10% increment in adherence to oral diabetes agents, HbA_{1c} was 0.19% lower ($P < 0.0001$). The change in HbA_{1c} over time was also strongly associated with adherence: for each 10% adherence increment, there was a 0.13% greater decrease in HbA_{1c} ($P < 0.0001$). HbA_{1c} was 0.26% lower per decade increase in age ($P < 0.0001$) and tended to decrease more over time among older patients ($P = 0.04$). More intensive therapy (vis-à-vis number of oral agents and insulin use) was strongly associated with higher HbA_{1c} levels. The mean HbA_{1c} decreased over time by 0.25% more for whites than for African-Americans ($P = 0.05$), with the most recent HbA_{1c} being 0.29% higher among African-Americans ($P = 0.04$). No association was found between metabolic control or change in control over time and income/copayment level, gender, frequency of HbA_{1c} testing, frequency of visits, or proportion of visits to a single provider (continuity index).

In the multivariable analyses (Table 3), the independent association of diabetes medication adherence with metabolic control and its change over time was un-

diminished by controlling for the other demographic and clinical factors. The most recent HbA_{1c} was 0.16% lower for each 10% increment in adherence ($P < 0.0001$); the absolute change over time was 0.13% greater for each 10% increment in medication adherence ($P < 0.0001$). The intensity of drug therapy (both oral agents and insulin) also maintained very strong inverse associations with metabolic control and its change over time. The strength and magnitude of the association between race and HbA_{1c} was also unchanged in the multivariable models. The HbA_{1c} of African-Americans was still an average of 0.29% higher than that of whites ($P = 0.04$) and improved less over time. The strength of the association between age and HbA_{1c} level diminished after adjustment and there was no

longer an association between age and HbA_{1c} change over time. The model without interaction terms accounted for 16% of the variance in metabolic control, although race explained <1%.

Interactions between race, number of oral agents, insulin use, and adherence were assessed in both multivariable models. There was a significant interaction between race and number of oral agents ($P = 0.04$) in the metabolic control model, accounting for 1.7% of the variance in HbA_{1c} (model R² improved to 17%). As shown in Table 4, the HbA_{1c} level of African-Americans was progressively greater than that of whites with increasing intensity of oral therapy. The association of insulin use and adherence with metabolic control did not vary between African-American and white patients. No other interactions were statistically significant in either model.

Adherence to nondiabetes medications was highly correlated with adherence to diabetes medications in the study population ($r = 0.51, P < 0.0001$). However, when added to the multiple regression models, adherence to nondiabetes medications was not independently associated with HbA_{1c} or change in HbA_{1c} level.

Although adjusting for medication adherence did not alter the association between race and metabolic control, African-American patients in this study population had substantially lower calculated adherence levels to diabetes medications than white patients (mean adherence 76.5 vs. 82%, $P = 0.0002$). Despite having poorer metabolic control on

Table 3—Multivariable associations with diabetes metabolic control

Independent variable	Dependent variable: most recent HbA _{1c}			Dependent variable: change in HbA _{1c} *		
	Parameter estimate	P value	Partial R ²	Parameter estimate	P value	Partial R ²
Race (white = 0)	0.29	0.04	0.5%	0.28	0.04	0.4%
Sex (male = 0)	-0.21	0.13	NA	-0.12	0.36	NA
Age (years)	-0.014	0.02	0.7%	-0.005	0.38	NA
Income/copay	Multiple	0.12	NA	Multiple	0.57	NA
Adherence (0–100%)	-0.016	<0.0001	2.7%	0.013	<0.0001	1.5%
No. oral agents	0.89	<0.0001	7.5%	0.63	<0.0001	3.6%
Insulin use (0 or 1)	0.75	<0.0001	2.6%	0.50	0.001	1.1%
No. HbA _{1c} levels	-0.09	0.06	NA	-0.05	0.33	NA
No. encounters	-0.006	0.61	NA	0.004	0.68	NA
Continuity index	-0.10	0.74	NA	0.28	0.35	NA

*The initial HbA_{1c} level was controlled for in this model.

Table 4—HbA_{1c} level as a function of race and number of oral agents

Number of oral agents	African-American: mean HbA _{1c} (n)	White: mean HbA _{1c} (n)
One	7.79% (181)	7.76% (222)
Two	8.81% (150)	8.35% (213)
Three	10.78% (15)	9.37% (48)

average, African-Americans tended to take fewer oral diabetes drugs than white patients (mean number 1.52 vs. 1.64, $P = 0.006$); odds of taking two or more oral agents were 1.3 times higher for whites, and odds of taking three agents were 2.4 times higher ($P = 0.006$). The odds of taking a thiazolidinedione was 1.5 times higher for whites ($P = 0.04$). The proportion of African-Americans and whites taking insulin was similar (29.8 vs. 30.8%, $P = 0.74$). The number of physician encounters and the number of HbA_{1c} tests did not differ between African-American and white patients, nor did continuity of care, as defined by the proportion of visits to a single primary care physician.

CONCLUSIONS— In summary, type 2 diabetes metabolic control and its improvement over time were strongly associated with adherence to diabetes medication in a large relatively indigent clinic population in central Virginia. Other predictors of metabolic control included age, number of oral diabetes medications, insulin use, and race. Controlling for these and other factors by multivariable analysis did not diminish the adherence association.

The strong association between adherence to diabetes drug therapy and metabolic control stands in contrast to several previously published reports. A prospective study by Wooldridge (17) of 189 patients with diabetes, two-thirds with type 2 diabetes, demonstrated improved metabolic control after intensive individual education but no association of HbA_{1c} with self-reported adherence to therapy. Glasgow et al. (15) also found no association between self-reported insulin adherence and HbA_{1c} among 93 individuals with type 1 diabetes. Reports from the Medical Outcomes Study by Hays et al. (18) and Kravitz et al. (16) also demonstrated no association between self-reported adherence and HbA_{1c} levels,

except among insulin-requiring patients and a subset with myocardial infarction. However, the reliance of these and other studies (19) on self-reported adherence calls into question the validity of their negative findings (20,21).

Other small studies using more objective adherence measures have demonstrated associations with diabetes metabolic control. Diehl et al. (22) used pill counts to assess adherence in 77 patients with diabetes and found a trend toward higher fasting blood glucose levels in those taking <80% of prescribed dosages. Chousa et al. (23) similarly found adherence assessed by pill count to be associated with metabolic control measured by HbA_{1c} among 107 patients with type 2 diabetes. Using self-reported adherence plus the pharmacy records of 65 patients with type 2 diabetes, Peterson et al. (24) also demonstrated an association between adherence and a composite measure of metabolic control, which included HbA_{1c}.

Our study confirms the findings of these earlier reports in a much larger sample of patients. There are several possible explanations for the demonstrated association. Clearly, more intensive pharmacologic management of type 2 diabetes and adherence to such therapy should generally lead to lower HbA_{1c} levels. The widespread use in the study population of newer drug therapies and combinations not available in previously published analyses of adherence and metabolic control would be expected to enhance such an association. In this light, our finding that a 30% absolute increase in adherence was, on average, associated with a roughly 0.5% absolute improvement in HbA_{1c} level seems quite plausible. Alternative or contributory explanations for the association between drug adherence and metabolic control are also possible. Higher adherence to drug therapy may be a marker for better diabetes self-management practices with respect to diet, exercise, and monitoring. Because these other factors were not measured in our study, they may have confounded the demonstrated association of adherence and HbA_{1c}. It is also possible that many potential barriers to adherence (educational, social, cultural, financial, medical, psychiatric, and other), also unmeasured in this study, may contribute to poor metabolic control independent of their impact on adherence. The strong correlation

demonstrated between adherence to diabetes and nondiabetes medications is consistent with these alternative explanations. However, the lack of an independent association between nondiabetes medication adherence and metabolic control suggests that the demonstrated association of adherence to diabetes medication with HbA_{1c} level is not entirely due to nonspecific underlying patient adherence characteristics.

It has been recognized that an open and compassionate physician communication style can facilitate improved patient adherence and outcome (25–27), whereas prevailing judgmental styles may impede it (28). Therefore, unmeasured physician factors may have contributed to or confounded the association between medication adherence and metabolic control demonstrated in this study. Because the physician is a logical locus of intervention to improve patient adherence, this area may be worthy of further research.

An additional caveat that must be applied to the demonstrated association is that the use of prescription refill–based data to estimate drug adherence is clearly imperfect. However, it does have the advantage of being an unobtrusive and objective source of adherence estimates and has been validated as a reasonably accurate measure in the absence of alternative patient sources of medication (29–32). The patients with very low income in the present report were only able to receive heavily subsidized medications from a single source. Nevertheless, intercurrent hospitalizations or alternative sources may have falsely lowered apparent adherence, and medication sharing, hoarding, or wasting could have made adherence appear higher than actual. Furthermore, information on adherence to dosing schedules is not available through pharmacy data.

Our finding of an association between increasing age and better metabolic control is consistent with previous reports. Using National Health and Nutrition Examination Survey (NHANES) III data, Shorr et al. (33) demonstrated a significant inverse association between age and HbA_{1c} among those on oral hypoglycemic agents. Nichols et al. (34) found a similar association among 1,333 patients with type 2 diabetes from a managed care population. Previous studies have also shown significant associations between age and

adherence to diabetes regimens (33,35). Our study suggests that the association between age and HbA_{1c} may be largely mediated by adherence, because the strength of the association greatly diminished after controlling for drug adherence.

The association of number of oral medications and insulin use with worsened metabolic control was expected, based on the literature as well as customary clinical practice. Because oral hypoglycemics and insulin are generally added as a result of worsening metabolic control, it is logical that a strong association would exist between required intensity of therapy and HbA_{1c} level. Such an association has been demonstrated previously for oral agents and insulin in general (34,36,37). We found no mention of gradations based on number of oral agents as shown in our study, likely due to the relatively recent availability of multiple drug classes.

The study finding of an association between race, metabolic control, and adherence also deserves further comment. It is well established that African-Americans have a higher prevalence of diabetes (38,39), poorer metabolic control (34,36,40,41), and worse outcomes (42,43) than white Americans. This disparity has not previously been explained by between-group differences in insurance coverage (44), physician visits (42,45), or prescribed drug therapy (42,45,46). Although studies have shown that lower socioeconomic status and minority patients self-monitor blood glucose less often (42,47,48), little is known regarding the linkage between adherence to drug therapy and metabolic control in these groups. Because overall drug adherence may be lower among African-Americans (49,50), our study examined the hypothesis that variation in diabetes medication adherence might explain the racial association with metabolic control and disease outcomes.

Although our study confirmed a significant association between race, adherence, and metabolic control in this rural indigent population, we did not find that measured adherence disparities solely mediated the inferior metabolic control in African-American patients. The 0.3% racial difference in HbA_{1c} levels found by bivariable analysis persisted after controlling for multiple other variables, including adherence. Despite similar rates of physician visits and HbA_{1c} test-

ing, African-Americans were also found to have progressively higher HbA_{1c} levels than white patients as treatment intensified. Because proportionately fewer African-Americans were being treated with multiple drug regimens for diabetes, it seems that the HbA_{1c} threshold used to intensify their therapy may have been higher. Our findings are consistent with previous reports that have shown similar rates of insulin therapy overall but significantly fewer intensive multiple-dose regimens among African-Americans versus white patients with diabetes (42,45).

There are several potential explanations for these findings as well as cautions that must be applied to their interpretation. It is possible that the racial disparity was due to a confounding variable that was not accounted for in the model. Neither educational nor occupational status was available to the investigators, and although income was included in the model, there may have been racial differences within the income strata used. Unmeasured confounding factors such as comorbidity, blood glucose self-monitoring, or patient reluctance to intensify therapy might also explain the observed associations. These and other factors may have contributed to the differences noted in metabolic control, drug treatment, and adherence between African-American and white patients in the study population. In addition, the effect of concordance of physician-patient ethnicity on these disparities could not be evaluated due to the small number of African-American physicians at the study site. Others have shown that cultural and communication barriers, among other issues, associated with nonconcordant physician-patient ethnicity may lead to less participatory decision making, greater mistrust, and poorer outcomes (51–54).

It must also be noted that the racial difference in metabolic control may have been due to a type I error. However, this is made less likely by the consistency of the findings from the different study models and the fact that the results are similar to that of other reports. Nonetheless, it must be emphasized that race explained only a very small amount of the variability in metabolic control in our models (~1%). Additionally, due to the fact that this was a single low-income patient population receiving highly subsidized care at one academic institution, the

study results do not indicate that the same associations would be found in other settings. Although not necessarily generalizable, these results add to the pattern found elsewhere and suggest that racial disparities in diabetes metabolic control may be attributable, in part, to less intensive and delayed drug therapy among African-Americans, as well as lower adherence levels. Even if such findings are due to unmeasured intervening or confounding effects, they reinforce the call for greater attention to the healthcare needs of minority and otherwise underserved populations that are known to have poor diabetes outcomes.

In conclusion, medication adherence does seem to be associated with diabetes metabolic control in an indigent care setting. We believe that this further underlines the need in these populations to facilitate diabetes self-management behaviors in general. The association of race with diabetes metabolic control in the U.S. is also likely intertwined with issues of adherence behavior as well as differing intensity of diabetes therapy between African-Americans and whites, as demonstrated in our study. Clearly, efforts to foster culturally sensitive and appropriate care are especially critical in minority populations. As articulated by Anderson and Funnell (13), the goal should be to help educate, motivate, and empower patients to improve their self-care skills and take control of their disease rather than simply foster adherence to prescribed medications.

References

1. Haynes RB, McKibbon KA, Kanani R: Systematic review of randomized trials of interventions to assist patients to follow prescriptions for medications. *Lancet* 348: 383–386, 1996
2. Overcoming barriers to treatment adherence in minorities and persons living in poverty [article online], 2001. Available from <http://grants.nih.gov/grants/guide/rfa-files/RFA-HL-01-005.html>. Accessed 1 September 2001.
3. Polednak AP: *Racial and Ethnic Differences in Disease*. New York, Oxford University Press, 1989
4. Curry CL: Coronary artery disease in African-Americans. *Circulation* 83:1474–1475, 1991
5. Johnson PA, Lee TH, Cook EF, Rouan GW, Goldman L: Effect of race on the presentation and management of patients with acute chest pain. *Ann Intern Med* 118:

- 593–601, 1993
6. Ayanian JZ, Udvarhelyi IS, Gatsonis CA, Pashos CL, Epstein AM: Racial differences in the use of revascularization procedures after coronary angiography. *JAMA* 269: 2642–2646, 1993
 7. Eggers PW, Greenberg LG: Racial and ethnic differences in hospitalization rates among aged Medicare beneficiaries, 1998. *Health Care Fin Rev* 21:91–105, 2000
 8. The Diabetes Control and Complications Trial Research Group: The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med* 329:977–986, 1993
 9. UK Prospective Diabetes Study (UKPDS) Group: Effect of intensive blood-glucose control with metformin on complications in overweight patients with type 2 diabetes (UKPDS 34). *Lancet* 352:854–865, 1998
 10. UK Prospective Diabetes Study (UKPDS) Group: Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet* 352:837–853, 1998
 11. McNabb WL: Adherence in diabetes: can we define it and can we measure it? *Diabetes Care* 20:215–218, 1997
 12. Johnson SB: Methodological issues in diabetes research: measuring adherence. *Diabetes Care* 15:1658–1667, 1992
 13. Anderson RM, Funnell MM: Compliance and adherence are dysfunctional concepts in diabetes care. *Diabetes Educator* 26: 597–604, 2000
 14. Watkins JD, Williams TF, Martin DA, Hogan MD, Anderson E: A study of diabetic patients at home. *Am J Public Health* 57:452–457, 1967
 15. Glasgow RE, McCaul KD, Schafer LC: Self-care behaviors and glycemic control in type I diabetes. *J Chron Dis* 40:399–412, 1987
 16. Kravitz RL, Hays RD, Sherbourne CD, et al: Recall of recommendations and adherence to advice among patients with chronic medical conditions. *Arch Intern Med* 153:1869–1878, 1993
 17. Wooldridge KL, Wallston KA, Graber AL, et al: The relationship between health beliefs, adherence, and metabolic control of diabetes. *Diabetes Educator* 18:495–500, 1992
 18. Hays RD, Kravitz RL, Mazel RM, et al: The impact of patient adherence on health outcomes for patients with chronic disease in the Medical Outcomes Study. *J Behav Med* 17:347–360, 1994
 19. Hepburn DA, Langan SJ, Deary IJ, Macleod KM, Frier BM: Psychological and demographic correlates of glycaemic control in adult patients with type 1 diabetes. *Diabet Med* 11:578–582, 1994
 20. Caron HS: Adherence: the case for objective measurement. *J Hypertens Suppl* 3: S11–S17, 1985
 21. Bond WS, Hussar DA: Detection methods and strategies for improving medication adherence. *Am J Hosp Pharm* 48:1978–1988, 1991
 22. Diehl AK, Bauer RL, Sugarek NJ: Correlates of medication compliance in non-insulin-dependent diabetes mellitus. *South Med J* 80:332–335, 1987
 23. Chousa FP, Guillen VFG, Otero MD, Beltran DO, Lopez RP, Sanchez JM: Usefulness of six indirect methods to evaluate drug therapy compliance in non insulin dependent diabetes mellitus. *Revista Clinica Espanola* 197:555–559, 1997
 24. Peterson GM, McLean S, Senator GB: Determinants of patient compliance, control, presence of complications, and handicap in non-insulin-dependent diabetes. *Aust N Z J Med* 14:135–141, 1984
 25. O'Connor PJ: From blame to understanding: moving diabetes care forward. *J Fam Pract* 46:205–206, 1998
 26. Bultman DC, Svarstad BL: Effects of physician communication style on client medication beliefs and adherence with antidepressant treatment. *Pat Educ Counsel* 40:173–185, 2000
 27. Coleman VR: Physician behavior and compliance. *Hypertension* 3:S69–S71, 1985
 28. Aaker E, Knudsen A, Wynn R, Lund A: General practitioners' reactions to non-compliant patients. *Scand J Prim Health Care* 19:103–106, 2001
 29. Choo PW, Rand CS, Inui TS, Lee ML, Cain E, Cordeiro-Breault M, Canning C, Platt R: Validation of patient reports, automated pharmacy records, and pill counts with electronic monitoring of adherence to antihypertensive therapy. *Med Care* 37:846–857, 1999
 30. Farmer KC: Methods for measuring and monitoring medication regimen adherence in clinical trials and clinical practice. *Clin Ther* 21:1074–1090, 1999
 31. Grymonpre RE, Didur CD, Montgomery PR, Sitar DS: Pill count, self-report, and pharmacy claims: data to measure medication adherence in the elderly. *Ann Pharmacother* 32:749–754, 1998
 32. Christensen DB, Williams B, Goldberg HI, et al: Assessing adherence to antihypertensive medication using computer-based pharmacy records. *Med Care* 25: 1164–1170, 1997
 33. Shorr RI, Franse LV, Resnick HE, Di Bari M, Johnson KC, Pahor M: Glycemic control of older adults with type 2 diabetes: findings from the Third National Health and Nutrition Examination Survey, 1988–1994. *J Am Geriatr Soc* 48:264–267, 2000
 34. Nichols GA, Hillier TA, Javor K, Brown JB: Predictors of glycemic control in insulin-using adults with type 2 diabetes. *Diabetes Care* 23:273–277, 2000
 35. Alogna M: Perception of severity of disease and health locus of control in compliant and noncompliant diabetic patients. *Diabetes Care* 3:533–534, 1980
 36. Harris MI, Eastman RC, Cowie CC, Flegal KM, Eberhardt MS: Racial and ethnic differences in glycemic control of adults with type 2 diabetes. *Diabetes Care* 22: 403–408, 1999
 37. Cook CB, Ziemer DC, El-Kebbi IM, Gallina DL, Dunbar VG, Ernst KL, Phillips LS: Diabetes in urban African-Americans: XVI. Overcoming clinical inertia improves glycemic control in patients with type 2 diabetes. *Diabetes Care* 22: 1494–1500, 1999
 38. Harris MI: Noninsulin-dependent diabetes mellitus in black and white Americans. *Diabetes Metab Rev* 6:71–90, 1990
 39. Harris MI, Flegal KM, Cowie CC, Eberhardt MS, Goldstein DE, Little RR, Wiedmeyer H-M, Byrd-Holt DD: Prevalence of diabetes, impaired fasting glucose, and impaired glucose tolerance in U.S. adults: the Third National Health and Nutrition Examination Survey 1988–1994. *Diabetes Care* 21:518–524, 1998
 40. Delamater AM, Shaw KH, Applegate EB, Pratt IA, Eidson M, Lancelotta GX, Gonzalez-Mendoza L, Richton S: Risk for metabolic control problems in minority youth with diabetes. *Diabetes Care* 22: 700–705, 1999
 41. Summerson JH, Konen JC, Dignan MB: Race-related differences in metabolic control among adults with diabetes. *South Med J* 85:953–956, 1992
 42. Harris MI: Racial and ethnic differences in health care access and health outcomes for adults with type 2 diabetes. *Diabetes Care* 24:454–459, 2001
 43. Harris EL, Sherman S, Georgopoulos A: Black-white differences in risk of developing retinopathy among individuals with type 2 diabetes. *Diabetes Care* 22:779–783, 1999
 44. Harris MI: Racial and ethnic differences in health insurance coverage for adults with diabetes. *Diabetes Care* 22:1679–1682, 1999
 45. Cowie CC, Harris MI: Ambulatory medical care for non-Hispanic whites, African-Americans, and Mexican-Americans with NIDDM in the U.S. *Diabetes Care* 20:142–147, 1997
 46. Harris MI: Epidemiological correlates of NIDDM in Hispanics, whites, and blacks in the U.S. population. *Diabetes Care* 14: 639–648, 1991
 47. Harris MI, Cowie CC, Howie LJ: Self-monitoring of blood glucose by adults with diabetes in the United States population. *Diabetes Care* 16:1116–1123, 1993

48. Karter AJ, Ferrara A, Darbinian JA, Ackerson LM, Selby JV: Self-monitoring of blood glucose: language and financial barriers in a managed care population with diabetes. *Diabetes Care* 23:477–483, 2000
49. Laine C, Newschaffer CJ, Zhang DZ, Cosler L, Hauck WW, Turner BJ: Adherence to antiretroviral therapy by pregnant women infected with human immunodeficiency virus: a pharmacy claims-based analysis. *Obstet Gynecol* 95: 167–173, 2000
50. Monane M, Bohn RL, Gurwitz JH, Glynn RJ, Levin R, Avorn J: Compliance with antihypertensive therapy among elderly Medicaid enrollees: the roles of age, gender, and race. *Am J Public Health* 86:1805–1808, 1996
51. Cooper-Patrick L, Gallo JJ, Gonzales JJ, Vu HT, Powe NR, Nelson C, Ford DE: Race, gender, and partnership in the patient-physician relationship. *JAMA* 282: 583–589, 1999
52. LaVeist TA, Nickerson KJ, Bowie JV: Attitudes about racism, medical mistrust, and satisfaction with care among African American and white cardiac patients. *Med Care Res Rev* 57 (Suppl. 1): 146–161, 2000
53. Doescher MP, Saver BG, Franks P, Fiscella K: Racial and ethnic disparities in perceptions of physician style and trust. *Arch Fam Med* 9:1156–1163, 2000
54. Perez-Stable EJ, Napoles-Springer A, Miramontes JM: The effects of ethnicity and language on medical outcomes of patients with hypertension or diabetes. *Med Care* 35:1212–1219, 1997