

# Prospective Study of C-Reactive Protein in Relation to the Development of Diabetes and Metabolic Syndrome in the Mexico City Diabetes Study

THANG S. HAN, MD, PHD<sup>1</sup>  
NAVEED SATTAR, MD, PHD<sup>2</sup>  
KEN WILLIAMS, MS<sup>3</sup>

CLICERIO GONZALEZ-VILLALPANDO, MD<sup>4</sup>  
MICHAEL E.J. LEAN, MD, FRCP<sup>5</sup>  
STEVEN M. HAFFNER, MD, MPH<sup>3</sup>

**OBJECTIVE** — Recent evidence suggests that C-reactive protein (CRP) may predict development of diabetes in Caucasian populations. We evaluated CRP as a possible risk factor of the development of diabetes and metabolic syndrome in a 6-year study of 515 men and 729 women from the Mexico City Diabetes Study.

**RESEARCH DESIGN AND METHODS** — Baseline CRP, indexes of adiposity, and insulin resistance (homeostasis model assessment [HOMA-IR]) were used to predict development of the metabolic syndrome, defined as including two or more of the following: 1) dyslipidemia (triglyceride  $\geq 2.26$  mmol/l or HDL cholesterol  $\leq 0.91$  mmol/l in men and  $\leq 1.17$  mmol/l in women;  $< 35$  and  $40$  mg/dl for men and women); 2) hypertension (blood pressure  $> 140/90$  mmHg or on hypertensive medication); or 3) diabetes (1999 World Health Organization criteria).

**RESULTS** — At baseline, CRP correlated significantly ( $P < 0.001$ ) with all metabolic indexes in women, but less so in men. After 6 years, 14.2% of men and 16.0% of women developed the metabolic syndrome. Compared with tertile 1, women with CRP in the highest tertile had an increased relative risk of developing the metabolic syndrome by 4.0 (95% CI 2.0–7.9) and diabetes by 5.5 (2.2–13.5); these risks changed minimally after adjusting for BMI or HOMA-IR. The area under receiver-operating characteristic (ROC) curve for the prediction of the development of the syndrome was 0.684 for CRP, increasing to 0.706 when combined with BMI and to 0.710 for a complex model of CRP, BMI, and HOMA-IR.

**CONCLUSIONS** — CRP was not a significant predictor of the development of the metabolic syndrome in men. Our data strongly support the notion that inflammation is important in the pathogenesis of diabetes and metabolic disorders in women.

*Diabetes Care* 25:2016–2021, 2002

**C**-reactive protein (CRP) is an acute-phase reactant that is a marker of inflammation in the body. Mild chronic elevations of CRP concentrations,

even when within the clinically “normal” range, are independently predictive of future cardiovascular events (1,2). Recent cross-sectional studies show that elevated

From <sup>1</sup>Addenbrooke’s Hospital, Cambridge University Medical School, Cambridge, U.K.; the <sup>2</sup>University Department of Pathological Biochemistry, Glasgow Royal Infirmary, Glasgow, U.K.; the <sup>3</sup>Department of Medicine #7873, University of Texas Health Science Center at San Antonio, San Antonio, Texas; the <sup>4</sup>Center de Estudios in Diabetes, Mexico City, Mexico; and the <sup>5</sup>Department of Human Nutrition, Glasgow Royal Infirmary, Glasgow, U.K.

Address correspondence and reprint requests to Steven M. Haffner, Department of Medicine #7873, University of Texas Health Science Center at San Antonio, 7703 Floyd Curl Dr., San Antonio, TX 78229-3900. E-mail: haffner@uthscsa.edu.

Received for publication 17 December 2001 and accepted in revised form 4 August 2002.

**Abbreviations:** CRP, C-reactive protein; HOMA-IR, homeostasis model assessment of insulin resistance; ROC, receiver-operating characteristic.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

CRP levels correlate significantly with features of the metabolic syndrome (insulin resistance syndrome), including adiposity, hyperinsulinemia, insulin resistance, hypertriglyceridemia, and low HDL cholesterol (3,4). A recent study has also demonstrated that high levels of CRP are related to increased accumulation of visceral and subcutaneous fat depots measured by computerized tomography scan (5). Two studies have reported that CRP predicts development of diabetes in postmenopausal women (6) and the elderly (7). These studies require confirmation in other populations and extension to other features of the metabolic syndrome.

Therefore, we assessed the 6-year development of diabetes and the metabolic syndrome in relation to baseline CRP levels in 515 men and 729 women from the Mexico City Diabetes Study. The metabolic syndrome was defined as subjects developing two or more of the metabolic disorders (dyslipidemia, hypertension, or diabetes) (8). We also compared the clinical value of CRP with indexes of adiposity and insulin resistance as a simple risk factor for the development of metabolic disorders.

## RESEARCH DESIGN AND METHODS

The Mexico City Diabetes Study is a population-based study of diabetes and cardiovascular risk factors. Between 1990 and 1992, 3,326 men and nonpregnant women aged 35–64 years from several low income “colonias” in Mexico City were eligible for the survey. A home interview was completed by 93% of the eligible subjects, and 68% of these completed a medical examination in a clinic (8). The protocol was approved by the Institutional Review Boards of the University of Texas Health Science Center at San Antonio and the American British Cowdray Hospital in Mexico City, and all subjects gave written informed consent. As described in the present article, we studied 515 men and 729 women who, at

baseline, had serum CRP concentrations measured and were free of diabetes.

Subjects' weight, height, and waist and hip circumferences were measured using standard methods for the calculations of BMI ( $\text{kg}/\text{m}^2$ ) and waist-to-hip ratio (8). Systolic (first Korotkov phase) and diastolic (fifth Korotkov phase) blood pressures were measured in a seated position after 5 min at rest three times to the nearest even digit with a random-zero sphygmomanometer (Hawksley-Gelman, Lancing, Sussex, U.K.). The average of the second and third readings was defined as the subject's blood pressure.

Each subject's blood was sampled after a 12- to 14-h fast for the measurements of serum lipids, lipoproteins, CRP, insulin, and plasma glucose concentrations. Glucose and insulin concentrations were also measured 2 h after a 75-g oral glucose load. Plasma insulin was measured by a solid-phase radioimmunoassay (8). CRP concentrations were measured by an ultra-sensitive competitive immunoassay (antigens and antibodies from Calbiochem, Darmstadt, Germany) with an interassay coefficient of variation of 8.9% (9). Insulin resistance was calculated from the homeostasis model assessment (HOMA-IR):  $\text{Insulin resistance} = [\text{fasting insulin (units/ml)} \times \text{fasting glucose (mmol/l)}] / 22.5$  (8).

Information of subjects' lifestyle was obtained from their responses to questions about the number of cigarettes smoked, the number of alcoholic drinks consumed per week, and total (leisure and work) metabolic equivalents of activity performed.

Hypertension was defined as systolic blood pressure  $\geq 140$  or diastolic blood pressure  $\geq 90$  mmHg or hypertension diagnosed by a physician, and was confirmed by the use of antihypertensive medication that subjects were asked to bring to the examination (10). Hypertriglyceridemia was defined as  $\geq 2.26$  mmol/l ( $\geq 200$  mg/dl). HDL cholesterol concentration was considered low if it was  $\leq 0.91$  mmol/l ( $\leq 35$  mg/dl) in men and  $\leq 1.17$  mmol/l ( $\leq 40$  mg/dl) in women (10). Dyslipidemia was defined as subjects with either hypertriglyceridemia or low HDL cholesterol. Diabetes was diagnosed as fasting plasma glucose  $\geq 7$  mmol/l (126 mg/dl) or a 2-h glucose level  $\geq 11.1$  mmol/l (200 mg/dl). Subjects who did not meet these criteria but who were receiving oral antidiabetic agents or insu-

**Table 1—Baseline characteristics of 1,244 men and women in the Mexico City Study**

	Men (n = 515)	Women (n = 729)
Age (years)	46.0 $\pm$ 0.4	46.0 $\pm$ 0.3
Weight (kg)	72.6 $\pm$ 0.5	65.6 $\pm$ 0.4
Height (cm)	164.5 $\pm$ 0.2	151.4 $\pm$ 0.2
BMI ( $\text{kg}/\text{m}^2$ )	26.8 $\pm$ 0.2	28.6 $\pm$ 0.2
Waist circumference (cm)	93.1 $\pm$ 0.4	97.4 $\pm$ 0.5
Waist-to-hip ratio	0.964 $\pm$ 0.002	0.964 $\pm$ 0.003
CRP (mg/l)	2.29 $\pm$ 0.32	2.74 $\pm$ 0.20
Fasting insulin (IU/ml)	14.7 $\pm$ 0.6	16.7 $\pm$ 0.6
Fasting glucose (mmol/l)	4.72 $\pm$ 0.03	4.73 $\pm$ 0.02
Insulin resistance (HOMA)*	3.22 $\pm$ 0.13	3.68 $\pm$ 0.14
Triglycerides (mmol/l)	2.79 $\pm$ 0.09	1.98 $\pm$ 0.05
HDL cholesterol (mmol/l)	0.78 $\pm$ 0.01	0.90 $\pm$ 0.01
Systolic blood pressure (mmHg)	117.5 $\pm$ 0.6	114.4 $\pm$ 0.6
Diastolic blood pressure (mmHg)	74.9 $\pm$ 0.5	71.2 $\pm$ 0.4
Current smoking (%)	48.2	20.3

Data are means  $\pm$  SE. \*Insulin resistance calculated using the HOMA as  $\text{fasting insulin (units/ml)} \times \text{fasting glucose (mmol/l)} / 22.5$ .

lin were also considered to have diabetes regardless of their plasma glucose value. The metabolic syndrome was defined as two or more metabolic disorders (dyslipidemia, hypertension, or diabetes) (10). All independent variables were stratified into sex-specific tertiles for analyses of the risks of the development of metabolic disorders and the metabolic syndrome. CRP was stratified into tertiles at 0.91 and 1.66 mg/l in men and at 1.24 and 2.17 mg/l in women. After 6 years, subjects were recalled to ascertain the incidence of dyslipidemia, diabetes, and hypertension. All anthropometric, biochemical, and physiological measurements were performed in an identical manner to those used at the baseline examination (10).

All statistical analyses were performed using SAS statistical package version 8.01 (SAS Institute, Cary, NC) for men and women separately and were adjusted for age and lifestyle factors. Spearman's correlations were used to assess the relationship between CRP and biochemical factors. Logistic regression analyses were carried out to calculate the odds ratios for the development of the metabolic syndrome and its components (dependent variables) in subjects with elevated baseline CRP (second and third tertiles) compared with those with low baseline CRP (lowest tertile, i.e., the reference group), with additional adjustments for baseline indexes of adiposity and insulin resistance (independent variables).

**RESULTS**— Table 1 shows the base-

line characteristics of 515 men and 729 women who had similar age. Compared with men, women were lighter and shorter, but they had higher BMI and larger waist circumference. Women had higher concentrations of CRP, HOMA-IR, and HDL cholesterol, and they had lower levels of triglyceride and systolic and diastolic blood pressures. Of the women, 4% received hormonal replacement therapy.

CRP correlated significantly ( $P < 0.01$ ) with BMI ( $r = 0.34$ ), waist circumference ( $r = 0.32$ ), HOMA-IR (0.22), fasting glucose ( $r = 0.11$ ), triglyceride ( $r = 0.23$ ), HDL cholesterol ( $r = -0.17$ ), and systolic blood pressure ( $r = 0.16$ ) in women but only significantly ( $P < 0.05$ ) in men with BMI ( $r = 0.14$ ), waist circumference ( $r = 0.12$ ), HOMA-IR ( $r = 0.12$ ), and triglyceride ( $r = 0.10$ ).

Six years after the baseline examination of subjects who were free of diabetes but had one or none of other metabolic disorders (hypertension or dyslipidemia), 14.2% of men and 16.0% of women developed the metabolic syndrome, 45.4% of whom (44.1% of men and 46.2% of women) developed diabetes. The incidence of the metabolic syndrome adjusted for age, cigarette smoking, alcohol consumption, and physical activity was significantly higher in women with higher levels of CRP: 7.5, 18.1, and 23.8% in the first, second, and third CRP tertiles, respectively ( $P < 0.001$  for trend). This relationship was not significant in men (14.2, 11.9, and 14.1%, respectively, trend  $P = 0.54$ ).

**Table 2—Multivariate logistic regression analysis of baseline CRP in the prediction of features of the development of the metabolic syndrome with adjustment for age, cigarette smoking, alcohol consumption, and physical activity**

Dependent variable*	CRP tertile 2			CRP tertile 3		
	OR	95% CI	P	OR	95% CI	P
<b>Men</b>						
Hypertriglyceridemia	2.2	0.92–5.4	0.078	1.2	0.5–3.3	0.69
Low HDL cholesterol	1.2	0.4–3.6	0.71	1.07	0.3–3.5	0.92
Hypertension	0.8	0.3–2.1	0.71	1.0	0.4–2.4	0.99
Diabetes	0.7	0.3–1.6	0.37	0.8	0.4–2.0	0.70
Metabolic syndrome	0.8	0.4–1.7	0.54	0.9	0.5–2.0	0.88
<b>Women</b>						
Hypertriglyceridemia	1.2	0.6–2.3	0.67	1.9	0.97–3.6	0.064
Low HDL cholesterol	2.3	1.01–5.2	0.047	2.2	0.92–5.3	0.077
Hypertension	2.1	0.98–4.7	0.058	2.1	0.96–4.7	0.064
Diabetes	2.8†	1.1–7.3	0.036	5.4‡	2.2–13.4	<0.001
Metabolic syndrome	2.8†	1.4–5.5	0.003	4.1‡	2.1–8.0	<0.001

\*In each model, subjects were free of the corresponding disorder under analysis, but they might have had none or one of the other disorders at baseline (e.g., for the prediction of the development of hypertriglyceridemia, subjects were free of baseline hypertriglyceridemia, but they might have had none or just one of the other disorders). †P < 0.05, ‡P < 0.01 for heterogeneity by sex.

Table 2 shows that compared with women with low baseline CRP levels (lowest tertile, i.e., reference group), those with high baseline CRP levels (highest tertile) had a significant (P < 0.01) increase in the risk for the development of diabetes and for the metabolic syndrome. There were no significant relationships shown in men.

Table 3 shows that compared with


women in the lowest tertile of baseline CRP, the risk for developing the metabolic syndrome was increased by 2.8 (1.4–5.5) in women with baseline CRP in the second tertile and by 4.0 (2.0–7.9) in the third tertile. These odds ratios did not change substantially after adjusting for indexes of adiposity and insulin resistance. There were no significant relationships between CRP and the development of the

metabolic syndrome in men. There were little changes in the relationship after adjusting for BMI and insulin resistance. We repeated this analysis in “lean” and “overweight” subjects (stratified by median of baseline BMI) separately, and found that the incidence was higher in overweight subjects in each CRP tertile and highest in overweight subjects with the highest levels of CRP (Fig. 1). In both lean and over-

**Table 3—Odds ratio for the development of metabolic syndrome**

	CRP					
	Tertile 2*			Tertile 3*		
	OR	95% CI	P	OR	95% CI	P
<b>Men</b>						
CRP alone	0.8	0.4–1.7	0.590	1.0	0.5–2.1	0.696
CRP + BMI	0.7	0.3–1.6	0.408	0.9	0.4–2.0	0.869
CRP + waist circumference	0.8	0.4–1.7	0.610	1.0	0.5–2.1	0.992
CRP + waist-to-hip ratio	0.8	0.4–1.7	0.607	1.0	0.5–2.1	0.980
CRP + insulin resistance	0.8	0.4–1.7	0.563	1.0	0.5–2.1	0.972
CRP + BMI + insulin resistance	0.8	0.3–1.7	0.480	1.0	0.5–2.1	0.946
CRP + waist circumference + insulin resistance	0.8	0.4–1.8	0.634	1.0	0.5–2.2	0.996
<b>Women</b>						
CRP alone	2.8	1.4–5.5	0.004	4.0	2.0–7.9	<0.001
CRP + BMI	2.6	1.3–5.1	0.008	3.4	1.7–6.9	<0.001
CRP + waist circumference	2.6	1.3–5.1	0.007	3.7	1.9–7.3	<0.001
CRP + waist-to-hip ratio	2.6	1.3–5.2	0.007	3.7	1.9–7.3	<0.001
CRP + insulin resistance	3.1	1.5–6.4	0.002	3.8	1.9–7.7	<0.001
CRP + BMI + insulin resistance	3.0	1.5–6.1	0.003	3.5	1.7–7.2	<0.001
CRP + waist circumference + insulin resistance	3.1	1.5–6.3	0.002	3.7	1.8–7.4	<0.001

Data was pooled with sex interaction with baseline CRP levels in the second and third (highest) tertiles compared with those in the first (lowest) tertile, after adjustment for indexes of adiposity and insulin resistance as covariates in various models. Data were adjusted for baseline age, cigarette smoking, alcohol consumption, and physical activity. \*Reference group = CRP tertile 1.



**Figure 1**—Six-year incidence of the metabolic syndrome in “lean” and “overweight” women (stratified by median of baseline BMI) by tertiles of baseline CRP levels. Data were adjusted for age, cigarette smoking, alcohol consumption, and physical activity.


weight groups, high CRP levels remained significantly predictive of the metabolic syndrome (results not shown).

Figure 2 shows that the area under the receiver-operating characteristic (ROC) curve was 0.684 for CRP in the prediction of the development of metabolic syndrome, and increased to 0.706 when CRP was combined with BMI, and a further 0.004 increase was obtained when insulin resistance was added to this model.

**CONCLUSIONS**— The findings in the present study showed that Mexican women with elevated CRP levels had a significantly increased risk of the development of both diabetes and the metabolic syndrome independent of their adiposity and insulin resistance. In addition, the results showed that CRP alone was as good as other single risk factors in

the prediction of the syndrome. When CRP was combined with indexes of adiposity, the area under the ROC curve improved by  $\sim 0.02$ , but further inclusion of biochemical measures such as insulin resistance added only another 0.004.

Thus, our data in Mexican women complement the recent study by Pradhan et al. (6) who demonstrated a similar observation in predominantly Caucasian women where diabetes was initially identified by self-report and subsequently verified by a spectrum of criteria. Another study (7) demonstrates that CRP levels are associated with the development of diabetes in the elderly. This latter study did not examine men and women separately, was based on fewer individuals ( $n = 45$ ) who developed diabetes, and only assessed fasting glucose on one occasion after baseline at 3–4 years of follow-up. Our analysis, which identified a larger number of



**Figure 2**—ROC analyses of CRP, BMI, and insulin as covariates in the prediction of 6-year development of the metabolic syndrome in women. Data were adjusted for age, cigarette smoking, alcohol consumption, and physical activity. The higher the area under the curve (i.e., the greater the curvature away from the 50% line [zero prediction]), the greater the predictive power.

middle-aged subjects of both sexes with a more complete diagnosis of diabetes on the basis of both fasting glucose and 2-h post-glucose tolerance test data, is a significant addition to the above observations.

The other novel and important aspect of our study is that, in addition to the development of diabetes, we examined for the development of other features of the metabolic syndrome. In line with our expectations, CRP also predicted the development of the syndrome in women independently of obesity and insulin resistance, thereby strengthening its potential role in predicting metabolic disorders. The clinical relevance of our results in women is that CRP is the most commonly used and best standardized inflammatory marker and thus has the potential to be used in the prediction of diabetes and the metabolic syndrome. This issue has become more important since recent reports have suggested prevention/delay of diabetes with lifestyle intervention (11,12). A recent report (13) suggested that a fasting prediction equation may identify individuals at high risk of developing diabetes without requiring an oral glucose tolerance test. The ability of CRP to improve such diabetes risk assessments should be studied.

CRP is an inflammatory marker produced and released by the liver under the stimulation of cytokines including interleukin-6, interleukin-1, and tumor necrosis factor- $\alpha$ . The relationships between CRP levels and adiposity could be explained by recent findings of adipose tissue as a source for the production and release of cytokines (14). These inflammatory factors have also been linked to dyslipidemia, hypertension, and insulin action in previous cross-sectional studies (3,4) and such observations have been extended in the present prospective study. There exist plausible biochemical mechanisms for linking some of these components. For example, cytokines promote de novo hepatic fatty acid synthesis and interfere with lipoprotein lipase activity (the endothelial enzyme responsible for catabolism of triglyceride-rich lipoproteins) (15). Cytokines may also directly impede insulin-stimulated glucose uptake (16).

Most studies of CRP have analyzed men and women together and have been adjusted for sex, although a recent article by Hak et al. (17) did show a stronger association of inflammatory markers with

insulin concentrations in elderly women than in men. The present study explicitly tested for sex-CRP interactions. Elevated CRP was a significant predictor of the development of diabetes and the metabolic syndrome in women only, independent of obesity and insulin resistance. This relationship was not significant in men. There are several possible explanations for these sex differences. Firstly, it should be noted that many more women were included in this study and that more women developed both diabetes and the metabolic syndrome. Thus, the study had greater statistical power to detect meaningful results in women. Secondly, since CRP correlated with indexes of adiposity and other metabolic parameters more strongly in women than in men, low-grade inflammation may have a greater role in perturbing insulin action in women. Inflammatory factors may interact with female sex hormones such that chronic inflammation may reduce the protective effects of estrogen on body fat distribution and insulin action. Data from the present study cannot clarify this suggestion. Nevertheless, there is some evidence for an interaction between these two pathways. Previous studies have suggested that cytokines may interfere with estradiol secretion (18), whereas oral, but not transdermal (which avoids hepatic first-pass metabolism) estrogens, can increase CRP levels (19,20). Only 4% of women in the present study received hormonal replacement therapy, and adjustment for this factor (or any previous hormone use) did not change the results (data not presented). Clearly, further prospective data are needed to examine the potential of CRP to predict diabetes and metabolic disorders in men. In the present study, <50% of subjects who developed the metabolic syndrome had developed diabetes as a manifestation of the syndrome; thus, the risks attributable to the metabolic syndrome cannot simply be accounted for by diabetes alone.

It is now known that both lipid-lowering therapies (statins) and ACE inhibitors have anti-inflammatory properties (21,22), and preliminary findings have suggested that they may lower the risk of diabetes (23,24). In addition, insulin-sensitizing agents, such as thiazolidinediones (25), and physical exercise (26) also appear to have anti-inflammatory properties. More recent animal data suggest that salicylates prevent obe-

sity and diet-induced insulin resistance (27). Thus, reducing inflammation may have beneficial effects on the development of diabetes and other metabolic disorders. Future investigations are clearly needed to examine this possibility.

In conclusion, we show that the predictive value of CRP for the development of diabetes extends to Mexican women 35–64 years of age. We also show for the first time that CRP predicts the development of the metabolic syndrome independently of the levels of adiposity and insulin resistance in Mexican women, but this risk was not evident in men. We suggest that measurement of CRP alone or combined with BMI or waist circumference can be used instead of complicated measures (e.g., fasting insulin, which requires a fasting state) as a risk factor for developing diabetes and the metabolic syndrome. Thus, CRP may be of clinical value for monitoring steps for the prevention of diabetes and metabolic disorders.

**Acknowledgments**— This study was supported by grants from the National Heart, Lung, and Blood Institute (RO1HL24799 and R37HL36820). N.S. was supported by a SHERT Medical Research Foreign Travel Grant.

The authors thank Dr. Chris Allen (Clinical Dean) and Dr. Owen Edwards (Clinical Tutor) of Addenbrooke's Hospital (University of Cambridge, U.K.) for their support of this study by T.S.H. as part of his Clinical Elective Study.

#### References

- Ridker PM, Hennekens CH, Buring JE, Rifai N: C-reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. *N Engl J Med* 342:836–843, 2000
- Koenig W, Sund M, Frohlich M, Fischer HG, Lowel H, Doring A, Hutchinson WL, Pepys MB: C-reactive protein, a sensitive marker of inflammation, predicts future risk of coronary heart disease in initially healthy middle-aged men: results from the MONICA (Monitoring Trends and Determinants in Cardiovascular Disease) Augsburg Cohort Study, 1984 to 1992. *Circulation* 99:237–242, 1999
- Festa A, D'Agostino R Jr, Howard G, Mykkanen L, Tracy RP, Haffner SM: Chronic subclinical inflammation as part of the insulin resistance syndrome: the Insulin Resistance Atherosclerosis Study (IRAS). *Circulation* 102:42–47, 2000
- Yudkin JS, Stehouwer CD, Emeis JJ, Coppack SW: C-reactive protein in healthy subjects: associations with obesity, insulin resistance, and endothelial dysfunction: a potential role for cytokines originating from adipose tissue? *Arterioscler Thromb Vasc Biol* 19:972–978, 1999
- Forouhi NG, Sattar N, McKeigue PM: Relation of C-reactive protein to body fat distribution and features of the metabolic syndrome in Europeans and South Asians. *Int J Obes Relat Metab Disord* 25: 1327–1331, 2001
- Pradhan AD, Manson JE, Rifai N, Buring JE, Ridker PM: C-reactive protein, interleukin 6, and risk of developing diabetes mellitus. *JAMA* 286:327–334, 2001
- Barzilay JI, Abraham L, Heckbert SR, Cushman M, Kuller LH, Resnick HE, Tracy RP: The relation of markers of inflammation to the development of glucose disorders in the elderly: the Cardiovascular Health Study. *Diabetes* 50:2384–2389, 2001
- Haffner SM, Kennedy E, Gonzalez C, Stern MP, Miettinen H: A prospective analysis of the HOMA-IR model: the Mexico City Diabetes Study. *Diabetes Care* 19: 1138–1141, 1996
- Macy EM, Hayes TE, Tracy RP: Variability in the measurement of C-reactive protein in healthy subjects: implications for reference intervals and epidemiological applications. *Clin Chem* 43:52–58, 1997
- Haffner SM, Valdez RA, Hazuda HP, Mitchell BD, Morales PA, Stern MP: Prospective analysis of the insulin-resistance syndrome (syndrome X). *Diabetes* 41: 715–722, 1992
- Tuomilehto J, Lindström J, Erickson JG, Valle TT, Hämäläinen H, Hanne-Parikka P, Keinänen-Kiukaanniemi S, Lassko M, Louheranta A, Rastas M, Salminen V, Uusitupa M, for the Finnish Diabetes Prevention Study Group: Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med* 344:1343–1350, 2001
- Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, Nathan DM: Reduction of the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 346:393–403, 2002
- Stern MP, Williams K, Haffner SM: Identification of persons at high risk for type 2 diabetes mellitus: do we need the oral glucose tolerance test? *Ann Intern Med* 136: 575–581, 2002
- Mohamed-Ali V, Pinkney JH, Coppack SW: Adipose tissue as an endocrine and paracrine organ. *Int J Obes* 22:1145–1158, 1998
- Grunfeld C, Feingold KR: Regulation of lipid metabolism by cytokines during host defense. *Nutrition* 12 (Suppl. 1):

- S24–S26, 1996
16. Youd JM, Rattigan S, Clark MG: Acute impairment of insulin-mediated capillary recruitment and glucose uptake in rat skeletal muscle in vivo by TNF- $\alpha$ . *Diabetes* 49:1904–1909, 2000
  17. Hak AE, Pols HA, Stehouwer CD, Meijer J, Kiliaan AJ, Hofman A, Breteler MM, Witteman JC: Markers of inflammation and cellular adhesion molecules in relation to insulin resistance in nondiabetic elderly: the Rotterdam study. *J Clin Endocrinol Metab* 86:4398–4405, 2001
  18. Alpizar E, Spicer LJ: Effects of interleukin-6 on proliferation and follicle-stimulating hormone-induced estradiol production by bovine granulosa cells in vitro: dependence on size of follicle. *Biol Reprod* 50:38–43, 1994
  19. Cushman M, Legault C, Barrett-Connor E, Stefanick ML, Kessler C, Judd HL, Sakkinen PA, Tracy RP: Effect of postmenopausal hormones on inflammation-sensitive proteins: the Postmenopausal Estrogen/Progestin Interventions (PEPI) Study. *Circulation* 100:717–722, 1999
  20. Sattar N, Forouhi NG, Wild RA: C-reactive protein and hormone replacement therapy. *Circulation* 102: E96–E97, 2000
  21. Palinski W: New evidence for beneficial effects of statins unrelated to lipid lowering. *Arterioscler Thromb Vasc Biol* 21:3–5, 2001
  22. Gullestad L, Aukrust P, Ueland T, Espevik T, Yee G, Vagelos R, Froland SS, Fowler M: Effect of high- versus low-dose angiotensin converting enzyme inhibition on cytokine levels in chronic heart failure. *J Am Coll Cardiol* 34:2061–2067, 1999
  23. Freeman DJ, Norrie J, Sattar N, Neely RD, Cobbe SM, Ford I, Isles C, Lorimer AR, Macfarlane PW, McKillop JH, Packard CJ, Shepherd J, Gaw A: Pravastatin and the development of diabetes mellitus: evidence for a protective treatment effect in the West of Scotland Coronary Prevention Study. *Circulation* 103:357–362, 2001
  24. Yusuf S, Sleight P, Pogue J, Bosch J, Davies R, Dagenais G: Effects of an angiotensin-converting-enzyme inhibitor, ramipril, on cardiovascular events in high-risk patients: the Heart Outcomes Prevention Evaluation Study Investigators. *N Engl J Med* 342:145–153, 2000
  25. Jiang C, Ting AT, Seed B: PPAR-gamma agonists inhibit production of monocyte inflammatory cytokines. *Nature* 391:82–86, 1998
  26. Smith JK, Dykes R, Douglas JE, Krishnaswamy G, Berk S: Long-term exercise and atherogenic activity of blood mononuclear cells in persons at risk of developing ischemic heart disease. *JAMA* 281:1722–1727, 1999
  27. Yuan M, Konstantopoulos N, Lee J, Hansen L, Li ZW, Karin M, Shoelson SE: Reversal of obesity- and diet-induced insulin resistance with salicylates or targeted disruption of Ikkbeta. *Science* 293:1673–1677, 2001