



# Trends and Characteristics of Self-reported Case Presentation of Diabetes Diagnosis Among Youth From 2002 to 2010: Findings From the SEARCH for Diabetes in Youth Study

*Diabetes Care* 2015;38:e84–e85 | DOI: 10.2337/dc15-0157

Sharon H. Saydah,<sup>1</sup>  
Giuseppina Imperatore,<sup>1</sup>  
Leora Henkin,<sup>2</sup> Ralph D'Agostino Jr.,<sup>2</sup>  
Jasmin Divers,<sup>2</sup>  
Elizabeth J. Mayer-Davis,<sup>3</sup>  
Dana Dabelea,<sup>4</sup>  
Georgeanna Klingensmith,<sup>5</sup>  
Catherine Pihoker,<sup>6</sup> and  
Jean M. Lawrence<sup>7</sup>

Diagnosis of diabetes in youth is increasing in the U.S. (1,2). It is not known how much of this change is due to an increase in diabetes and how much is due to improved case detection, especially for type 2 diabetes. Some researchers have hypothesized that part of the explanation for the increase in diabetes diagnosis in youth is increased screening, resulting in a higher percentage of cases being identified. The objective of this study was to assess whether the change in diabetes could be explained by changes in case identification by examining trends from 2002 to 2010 in self-reported case presentation of diabetes.

Briefly, there were 9,054 youth aged <20 years with newly diagnosed diabetes between 2002 and 2010 in the SEARCH for Diabetes in Youth study (3). Participants were asked, "How did you find out you had diabetes?" Responses were grouped into symptoms, checkup, community screening, or other. Self-reported case presentation patterns were examined in 3-year blocks to assess change over time, reported by diabetes type. We explored trends in

self-reported modes of diabetes diagnosis (i.e., symptoms, checkup, screening, and other method) and reported results unadjusted and then adjusted for age-group, sex, and race/ethnicity.

Results are presented in Table 1. Among youth with type 1 diabetes, >95% of them reported diabetes diagnosis due to symptoms and many fewer reported diagnosis due to checkup, health screening, or other. Self-report of case presentation remained stable from 2002 to 2010 for youth with type 1 diabetes. Among youth with type 2 diabetes, 65% reported diagnosis due to symptoms and 30% reported diagnosis during a regular checkup. Unlike type 1 diabetes, there were significant changes in reported case presentation for type 2 diabetes with presentation due to symptoms decreasing from 72.1% in 2002–2004 to 59.1% in 2008–2010.

Observed differences in patterns of the self-reported modes of case presentation by age and sex among youth with type 1 diabetes and by sex and race/ethnicity among youth with type 2 diabetes may reflect differences in how

diabetes presents, medical-seeking practices, or community awareness.

Of the few previous studies reporting on modes of diabetes diagnosis, none have included youth (4,5). Previous studies have found that adults with diabetes are most likely to report diagnosis due to symptoms (4,5).

While we found no evidence that increased incidence of type 1 diabetes was due to improvements in case finding, there was evidence of increased case finding among youth with type 2 diabetes. The changes in reported modes of case presentation over time for type 2 diabetes suggest that some of the trends in prevalence over this time period may be attributable to changes in health care or community screening patterns.

**Acknowledgments.** The authors thank Tony Pearson-Clarke for his invaluable editorial assistance. The SEARCH for Diabetes in Youth Study is indebted to the many youth and their families and health care providers, whose participation made this study possible.

**Funding.** SEARCH for Diabetes in Youth is funded by the Centers for Disease Control and Prevention (PA numbers 00097, DP-05-069,

<sup>1</sup>Division of Diabetes Translation, National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention, Atlanta, GA

<sup>2</sup>Wake Forest School of Medicine, Winston-Salem, NC

<sup>3</sup>University of North Carolina at Chapel Hill, Chapel Hill, NC

<sup>4</sup>Colorado School of Public Health, University of Colorado Denver, Aurora, CO

<sup>5</sup>Barbara Davis Center for Diabetes and Department of Pediatrics, University of Colorado School of Medicine, Aurora, CO

<sup>6</sup>Department of Pediatrics, University of Washington, Seattle, WA

<sup>7</sup>Department of Research & Evaluation, Kaiser Permanente Southern California, Pasadena, CA

Corresponding author: Sharon H. Saydah, [ssaydah@cdc.gov](mailto:ssaydah@cdc.gov).

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention and the National Institute of Diabetes and Digestive and Kidney Diseases.

© 2015 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered.

**Table 1—Self-reported case presentation of diabetes diagnosis by incident year and diabetes type among youth 20 years of age and younger, SEARCH for Diabetes in Youth Study 2002–2010**

	Type 1 diabetes (n = 7,554)					Type 2 diabetes (n = 1,500)				
	Symptoms	Checkup	Community screening	Other	P value	Symptoms	Checkup	Community screening	Other	P value
Incident years										
2002–2004	95.4	3.7	0.2	0.7	0.0512	72.1	23.7	2.8	1.4	<0.0001
2005–2007	94.8	3.6	0.4	0.4		66.4	29.5	2.5	1.6	
2008–2010	94.2	4.1	0.3	0.3		59.1	37.1	3.3	0.5	
Adjusted estimates*										
2002–2004	95.9	3.2	0.2	0.2	0.0505	72.9	23.7	2.4	1.0	<0.0001
2005–2007	95.5	3.3	0.2	0.2		66.8	29.9	2.7	0.6	
2008–2010	94.9	3.4	0.3	0.3		59.8	36.9	3.0	0.4	

Data are %. \*Adjusted for age-groups (0 to < 5 years, 5 to < 10 years, 10 years to < 15 years, 15 years and older), sex, and race/ethnicity (Asian/Pacific).

and DP-10-001) and supported by the National Institute of Diabetes and Digestive and Kidney Diseases. Site contract numbers include Kaiser Permanente Southern California (U48/CCU919219, U01 DP000246, and U18DP002714), University of Colorado Denver (U48/CCU819241-3, U01 DP000247, and U18DP000247-06A1), Kuakini Medical Center (U58CCU919256 and U01 DP000245), Children's Hospital Medical Center (Cincinnati) (U48/CCU519239, U01 DP000248, and U18DP002709), University of North Carolina at Chapel Hill (U48/CCU419249, U01 DP000254, and U18DP002708), University of Washington School of Medicine (U58/CCU019235-4, U01 DP000244, and U18DP002710-01), and Wake Forest University School of Medicine (U48/CCU919219, U01 DP000250, and 200-2010-35171). The authors wish to acknowledge the involvement of General Clinical Research Centers at the South Carolina Clinical & Translational Research Institute at the Medical University of South Carolina (National Institutes of Health [NIH]/National Center for Research Resources grant number UL1RR029882), Seattle Children's Hospital (NIH Clinical and Translational Science Award grant UL1 TR00423 of the University of Washington), University of Colorado Pediatric Clinical Translational

Research Center (grant number UL1 TR000154), Barbara Davis Center for Diabetes at the University of Colorado Denver (Diabetes and Endocrinology Research Center NIH P30 DK57516), National Center for Research Resources and National Center for Advancing Translational Sciences, NIH (grant 8 UL1 TR000077), and the Children with Medical Handicaps Program managed by the Ohio Department of Health.

**Duality of Interest.** No potential conflicts of interest relevant to this article were reported.

**Author Contributions.** S.H.S., G.I., L.H., R.D., J.D., E.J.M.-D., D.D., G.K., C.P., and J.M.L. contributed to the conception and design of the work, the acquisition of the data, the interpretation of the data, and the revision of the manuscript. All authors provided final approval for publication and ensure the accuracy and integrity of the work. S.H.S. drafted the manuscript. S.H.S. and J.D. contributed to the analysis of the data. S.H.S. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

**Prior Presentation.** Parts of this study were presented in abstract form at the 73rd Scientific

Sessions of the American Diabetes Association, Chicago, IL, 21–25 June 2013.

## References

- Lawrence JM, Imperatore G, Dabelea D, et al.; SEARCH for Diabetes in Youth Study Group. Trends in incidence of type 1 diabetes among non-Hispanic white youth in the U.S., 2002-2009. *Diabetes* 2014;63:3938–3945
- Dabelea D, Mayer-Davis EJ, Saydah S, et al.; SEARCH for Diabetes in Youth Study. Prevalence of type 1 and type 2 diabetes among children and adolescents from 2001 to 2009. *JAMA* 2014; 311:1778–1786
- SEARCH Study Group. SEARCH for Diabetes in Youth: a multicenter study of the prevalence, incidence and classification of diabetes mellitus in youth. *Control Clin Trials* 2004;25:458–471
- Rodbard HW, Green AJ, Fox KM, Grandy S. Trends in method of diagnosis of type 2 diabetes mellitus: results from SHIELD. *Int J Endocrinol* 2009;2009:796206
- Clark NG, Fox KM, Grandy S; SHIELD Study Group. Symptoms of diabetes and their association with the risk and presence of diabetes: findings from the Study to Help Improve Early evaluation and management of risk factors Leading to Diabetes (SHIELD). *Diabetes Care* 2007;30:2868–2873