



RESPONSE TO COMMENT ON HUGHES ET AL.

## An Early Pregnancy HbA<sub>1c</sub> $\geq$ 5.9% (41 mmol/mol) Is Optimal for Detecting Diabetes and Identifies Women at Increased Risk of Adverse Pregnancy Outcomes. *Diabetes Care* 2014;37:2953–2959

*Diabetes Care* 2015;38:e93 | DOI: 10.2337/dc15-0401

We appreciate the opportunity to respond to the comments made by Anderson et al. (1) on our recent article (2). We agree, as highlighted under our study limitations, that the low participation rate in the oral glucose tolerance test (OGTT) could affect the validity of the HbA<sub>1c</sub> sensitivity and specificity calculations. As women with lower HbA<sub>1c</sub> (particularly <5.6% [38 mmol/mol]) were least likely to consent to an early OGTT, it is possible that the optimal HbA<sub>1c</sub> threshold for identifying women with undiagnosed preexisting diabetes (not gestational diabetes mellitus [GDM]) is in fact lower than 5.9% (41 mmol/mol) and we invite others to replicate our study.

The pregnancy outcome data, however, are available for all women who delivered at our center. The finding that a booking HbA<sub>1c</sub> within the prediabetes range (5.9–6.4%) was associated with adverse pregnancy outcomes in the absence of a subsequent diagnosis of GDM by OGTT criteria is significant, as current guidelines (2,3) do not recommend early intervention unless the HbA<sub>1c</sub> is  $\geq$ 6.5% (48 mmol/mol). In addition, as mentioned in our discussion, in a separate cohort of New Zealand women, Rowan et al. (4) reported that those with a nondiagnostic OGTT but an HbA<sub>1c</sub>  $\geq$ 5.9% have significant hyperglycemia with >90% referred before 24 weeks' gestation requiring pharmacotherapy. It is therefore possible that an

HbA<sub>1c</sub> measure is clinically more relevant than an OGTT in early pregnancy.

Women recruited for our study were from a primary care setting and invitations for an OGTT were by both telephone and mail. Women were approached in the same manner irrespective of their social or ethnic backgrounds. The ethnic split of women attending the early OGTT was similar to that of the women attending routine GDM screening in later pregnancy, with Maori women underrepresented despite being a high-risk group for diabetes. Interestingly, a greater proportion of Maori women attended for an early pregnancy HbA<sub>1c</sub>. HbA<sub>1c</sub> measurement may be a more acceptable screening test for preexisting diabetes, with a greater uptake than an OGTT in early pregnancy for certain ethnic populations.

We excluded the random plasma glucose (RPG) data from this article, as our goal was to present the HbA<sub>1c</sub> data as clearly as possible. RPG and HbA<sub>1c</sub> measurements did not correlate well; these data and the pregnancy outcome data related to RPG measurements will be published separately.

We acknowledge that iron deficiency, which is particularly prevalent in later pregnancy, can cause a slight upwards shift in HbA<sub>1c</sub> measurements equating to about 0.1% (1–2 mmol/mol) (5). In our study, we measured HbA<sub>1c</sub> in early pregnancy and ferritin was not measured at that time.

Ruth C.E. Hughes,<sup>1</sup> M. Peter Moore,<sup>2</sup> Joanna E. Gullam,<sup>1</sup> and Janet Rowan<sup>3</sup>

Importantly, because a diagnosis of diabetes or prediabetes has significant implications, we do not advocate labeling women based on an early pregnancy HbA<sub>1c</sub> measurement and believe that glycemic status should be reassessed postpartum. However, in our population, HbA<sub>1c</sub> was a useful marker to identify women with a higher risk of adverse pregnancy outcomes who may benefit from early intervention.

**Duality of Interest.** No potential conflicts of interest relevant to this article were reported.

### References

- Anderson HB, Rezaei S, Setlur A, Henderson CE. Comment on Hughes et al. An early pregnancy HbA<sub>1c</sub>  $\geq$ 5.9% (41 mmol/mol) is optimal for detecting diabetes and identifies women at increased risk of adverse pregnancy outcomes. *Diabetes Care* 2014;37:2953–2959 (Letter). *Diabetes Care* 2015; 38:e92. DOI: 10.2337/dc14-2962
- Hughes RCE, Moore MP, Gullam JE, Mohamed K, Rowan J. An early pregnancy HbA<sub>1c</sub>  $\geq$ 5.9% (41 mmol/mol) is optimal for detecting diabetes and identifies women at increased risk of adverse pregnancy outcomes. *Diabetes Care* 2014;37:2953–2959
- American Diabetes Association. Standards of medical care in diabetes—2015. *Diabetes Care* 2015;38(Suppl. 1):S1–S94
- Rowan JA, Budden A, Sadler LC. Women with a nondiagnostic 75 g glucose tolerance test but elevated HbA<sub>1c</sub> in pregnancy: an additional group of women with gestational diabetes. *Aust N Z J Obstet Gynaecol* 2014;54:177–180
- Ahmad J, Rafat D. HbA<sub>1c</sub> and iron deficiency: a review. *Diabetes Metab Syndr* 2013;7:118–122

<sup>1</sup>Department of Obstetrics and Gynecology, University of Otago, Christchurch Women's Hospital, Christchurch, New Zealand

<sup>2</sup>Diabetes Center, Christchurch Hospital, Christchurch, New Zealand

<sup>3</sup>National Women's Health, Auckland City Hospital, Auckland, New Zealand

Corresponding author: Ruth C.E. Hughes, ruth.hughes@cdhb.health.nz.

© 2015 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered.