



COMMENT ON LEESE ET AL.

Progression of Diabetes Retinal Status Within Community Screening Programs and Potential Implications for Screening Intervals. *Diabetes Care* 2015;38:488–494

Diabetes Care 2015;38:e207–e208 | DOI: 10.2337/dc15-1356

Focke Ziemssen, Lydia Marahrens, and Daniel Roeck

There is a discussion about the optimum screening interval for the timely detection of diabetic retinopathy (DR) (1). A low rate of progression in patients without any preexisting retinopathy leads to the hypothesis that extending screening intervals can generally reduce total costs. Unfortunately, the British Four Nations Diabetic Retinopathy Screening Study Group's recent article on the evaluation of DR progression has severe limitations (1).

1. When adhering to just the broad categorization of “referable” and “treatable” disease, the exact stage of DR remains nontransparent (1). This is a major drawback, as the most appropriate treatment might change in the future when new studies' data emerge or when the prediction of the individualized risk might improve. The authors best illustrate the restriction, describing that even within the U.K. there is no consensus between England and Scotland on the need for the specialist physicians' examinations. The team of authors mixed up the different stages of DR within the vague “composite terms.” Necessarily, the data have to be stratified as per the type of diabetes.
2. The statement by Leese et al. that “. . . usually only patients with proliferative retinopathy require immediate treatment at the first visit to an

ophthalmology clinic” is just incorrect. The authors ignore the recent evidence that a lag in the treatment of diabetic macular edema (DME) cannot be fully compensated for later on. At the very least, irreversible impairment has been proven by trials using a crossover design. Treating DME 1 year later cannot achieve the same level of improvement as immediate treatment. The most recent research characterizing anti-vascular endothelial growth factor (anti-VEGF) studies shows a strong ceiling effect with baseline acuity determining the final function (2). Thus, early therapy is very likely more cost-efficient—similar to neovascular macular degeneration therapy (3).

3. When systematically screening the literature, the bottom line of both of the most comprehensive reviews includes warnings against the high risks of prolonging the intervals between eye examinations for all patients (4,5). In addition, there is a large degree of heterogeneity regarding the screening methods (photographs, recall system, telemedicine, funduscopy), as access to the health system differs greatly. It is very likely that those staying away from free and promoted screenings have a lower level of adherence and health awareness (6). Therefore, the study might underestimate the rate

of progression by excluding a considerable number of people that have inadequate metabolic control. We kindly ask that the issue of nonappearance be precisely addressed in future publications.

4. The exact screening protocol and the imaging modality used play an important role in the sensitivity of the screening. The Diabetes Control and Complications Trial (DCCT) has already discovered a considerable rate of DR (22%) using angiography in individuals who only have had diabetes for a short time, but there was no evidence of DR on color photographs (7). How can the use of modern imaging devices such as ultrawidefield imaging or ocular coherence tomography further increase the ability to detect clinically significant stages of DR (8)?

We are very grateful that responsible British scientists, such as Simon Harding from Liverpool, are planning prospective trials with randomization for different screening intervals before drawing far-reaching conclusions.

Duality of Interest. F.Z. has received consulting fees from Alimera, Allergan, Bayer HealthCare, and Novartis and speakers' fees from Alcon, Alimera, Allergan, Bayer HealthCare, Heidelberg Engineering, and Novartis. No other potential conflicts of interest relevant to this article were reported.

Centre for Ophthalmology, University Eye Hospital, Tübingen, Germany

Corresponding author: Focke Ziemssen, focke.ziemssen@med.uni-tuebingen.de.

© 2015 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered.

References

1. Leese GP, Stratton IM, Land M, et al.; Four Nations Diabetic Retinopathy Screening Study Group. Progression of diabetes retinal status within community screening programs and potential implications for screening intervals. *Diabetes Care* 2015;38:488–494
2. Sophie R, Lu N, Campochiaro PA. Predictors of functional and anatomic outcomes in patients with diabetic macular edema treated with ranibizumab. *Ophthalmology* 2015;122:1395–1401
3. Butt T, Lee A, Lee C, Tufail A; UK AMD EMR Study Group. The cost-effectiveness of initiating ranibizumab therapy in eyes with neovascular AMD with good vision: an economic model using real-world outcomes. *BMJ Open* 2015;5:e006535
4. Echouffo-Tcheugui JB, Ali MK, Roglic G, Hayward RA, Narayan KM. Screening intervals for diabetic retinopathy and incidence of visual loss: a systematic review. *Diabet Med* 2013;30:1272–1292
5. Taylor-Phillips S, Mistry H, Leslie R, et al. Extending the diabetic retinopathy screening interval beyond 1 year: systematic review. *Br J Ophthalmol*. 13 January 2015 [Epub ahead of print]. DOI: 10.1136/bjophthalmol-2014-305938
6. Millett C, Dodhia H. Diabetes retinopathy screening: audit of equity in participation and selected outcomes in South East London. *J Med Screen* 2006;13:152–155
7. Malone JI, Morrison AD, Pavan PR, Cuthbertson DD; Diabetic Control and Complications Trial. Prevalence and significance of retinopathy in subjects with type 1 diabetes of less than 5 years' duration screened for the Diabetes Control and Complications Trial. *Diabetes Care* 2001;24:522–526
8. Silva PS, Cavallerano JD, Haddad NM, et al. Peripheral lesions identified on ultrawide field imaging predict increased risk of diabetic retinopathy progression over 4 years. *Ophthalmology* 2015;122:949–956