



Effects of Frequency of Sensor-Augmented Pump Use on HbA_{1c} and C-Peptide Levels in the First Year of Type 1 Diabetes

Diabetes Care 2016;39:e61–e62 | DOI: 10.2337/dc15-2201

Taylor M. Triolo,¹ David M. Maahs,¹ Laura Pyle,¹ Robert Slover,¹ Bruce Buckingham,² Peiyao Cheng,³ Linda A. DiMeglio,⁴ Andrew A. Bremer,⁵ Stuart A. Weinzimer,⁶ H. Peter Chase,¹ and Diabetes Research in Children Network (DirecNet) and Type 1 Diabetes TrialNet Study Groups

Intensive glucose control after the onset of type 1 diabetes has been suggested to preserve C-peptide production (1–3). In people with type 1 diabetes, sensor-augmented pump (SAP) use improved glycemic control, particularly when used >6 days/week (4). As previously reported, subjects with type 1 diabetes were randomized to either 3 days of in-hospital hybrid closed-loop therapy followed by SAP therapy or usual care (5). In the primary intent-to-treat analysis, no significant differences existed in HbA_{1c} or C-peptide between the two groups at 1 year. In our post hoc analysis, we hypothesized that more frequent sensor use in the SAP group would be associated with lower HbA_{1c} levels and preservation of C-peptide production at 1 year.

Subjects in the SAP group ($n = 46$) were stratified by median SAP use of 12.4 h/day. HbA_{1c} and C-peptide levels were compared at baseline and 3, 6, 9, 12, and 24 months. At 12 months of follow-up, the median HbA_{1c} values for those with SAP use above versus below the median were lower (7.0% [Q1, Q3 6.0, 7.5] vs. 7.7% [7.1, 8.5], $P = 0.007$).

All three C-peptide measures were 50–79% higher at 12 months in the above versus below median SAP use group. However, these nonsignificant trends were no longer present at 24 months. No statistically significant differences were seen in fasting, peak, or area under the curve C-peptide levels between the two groups at any time period over 2 years of follow-up. All three C-peptide measures declined >50% from 12 to 24 months in the high SAP use group, although HbA_{1c} levels remained similar (Fig. 1).

As HbA_{1c} levels remained similar at 12 and 24 months in the above median SAP use group, it can be concluded that factors other than glycemic control were likely related to the >50% reduction in C-peptide production during the second year after diagnosis. In subjects having type 1 diabetes for >1 year, only consistent (≥ 6 days/week) use of a SAP system was shown to improve glycemic control (2). In our analysis, we stratified subjects by the frequency of use of SAP over the year of follow-up (the primary end point), and median use (12.4 h) was about half the day. This frequency of SAP use may not have been sufficient to

reduce HbA_{1c} levels adequately to preserve β -cell function. The drop-off in frequency of continuous glucose monitor use in this and previous studies may be due to less sophisticated earlier-generation continuous glucose monitor technologies, such as the Medtronic Sof-sensor used in the initial study (5). Further studies with larger numbers of subjects who are followed for a longer time period and are using improved technology will be important to further evaluate this hypothesis in the future.

Funding. This research was supported by National Institutes of Health/Eunice Kennedy Shriver National Institute of Child Health and Human Development grants for the Diabetes Research in Children Network (DirecNet) Study Group (HD-41890-10, HD-41906-10, and HD-41908-10) and by grants for the Type 1 Diabetes TrialNet Study Group (U01-DK-085509, U01-DK-06104211, 5U01-DK-085466-05, U01-DK-085505-02, and 5U01-DK-085465-04). Other more complete information is available elsewhere (2,3).

Duality of Interest. D.M.M. is on the advisory board for Insulet, and his institution has received research funding from Medtronic and Dexcom. R.S. has received grant support from Medtronic MiniMed. B.B. received payment from Sanofi, GlySens, and Roche for serving on medical advisory boards; consulted for BD Biosciences;

¹University of Colorado Anschutz Medical Campus, Aurora, CO

²Stanford University, Stanford, CA

³Jaeb Center for Health Research, Tampa, FL

⁴Section of Pediatric Endocrinology and Diabetology, Department of Pediatrics, Indiana University School of Medicine, Indianapolis, IN

⁵Division of Pediatric Endocrinology, Vanderbilt University, Nashville, TN

⁶Yale University School of Medicine, New Haven, CT

Corresponding author: H. Peter Chase, peter.chase@ucdenver.edu.

Received 7 October 2015 and accepted 14 January 2016.

Clinical trial reg. no. NCT00760526, clinicaltrials.org.

© 2016 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered.




Figure 1—Comparison over 2 years of those who used the SAP above the median compared with those who used the SAP below the median use in the first 12 months. **A:** HbA_{1c} levels (%). **B:** Fasting C-peptide (nmol/L). **C:** Peak C-peptide (nmol/L). **D:** Area under the curve (AUC) C-peptide (nmol min/L). * $P < 0.05$ at 9 months and $P < 0.01$ at 12 and 24 months.

received payment for a lecture (symposium at the European Association for the Study of Diabetes) from Dexcom; received payment from Medtronic for serving on the Data and Safety Monitoring Board for Sensor-Augmented Pump Therapy for A1C Reduction (STAR 3) study; and received institutional payment from Medtronic for other principal investigator–initiated studies

and sponsored research. S.A.W. is on the advisory board for Insulet and has received honoraria and grant support from Medtronic. H.P.C. has received grant support from Dexcom. No other potential conflicts of interest relevant to this article were reported.

Author Contributions. T.M.T., D.M.M., L.P., R.S., B.B., A.A.B., P.C., and H.P.C. researched the

data and reviewed and edited the manuscript. T.M.T., D.M.M., L.P., and H.P.C. wrote the manuscript. L.A.D. and S.A.W. reviewed and edited the manuscript. H.P.C. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Prior Presentation. Data from this study were presented at the 75th Scientific Sessions of the American Diabetes Association, Boston, MA, 5–9 June 2015.

References

- Shah SC, Malone JJ, Simpson NE. A randomized trial of intensive insulin therapy in newly diagnosed insulin-dependent diabetes mellitus. *N Engl J Med* 1989;320:550–554
- Kordonouri O, Pankowska E, Rami B, et al. Sensor-augmented pump therapy for the diagnosis of childhood type 1 diabetes: results of the Paediatric Onset Study (ONSET) after 12 months of treatment. *Diabetologia* 2010;53:2487–2495
- Kordonouri O, Hartmann R, Pankowska E, et al. Sensor augmented pump therapy from onset of type 1 diabetes: late follow-up results of the Paediatric Onset Study. *Pediatr Diabetes* 2012;13:515–518
- Tamborlane WV, Beck RW, Bode BW, et al.; Juvenile Diabetes Research Foundation Continuous Glucose Monitoring Study Group. Continuous glucose monitoring and intensive treatment of type 1 diabetes. *N Engl J Med* 2008;359:1464–1476
- Buckingham B, Beck RW, Ruedy KJ, et al.; Diabetes Research in Children Network (DirecNet) Study Group; Type 1 Diabetes TrialNet Study Group. Effectiveness of early intensive therapy on β -cell preservation in type 1 diabetes. *Diabetes Care* 2013;36:4030–4035