



# Folic Acid Supplement Intake in Early Pregnancy Increases Risk of Gestational Diabetes Mellitus: Evidence From a Prospective Cohort Study

*Diabetes Care* 2016;39:e36–e37 | DOI: 10.2337/dc15-2389

Beibei Zhu,<sup>1,2</sup> Xing Ge,<sup>1</sup> Kun Huang,<sup>1,2</sup>  
Leijing Mao,<sup>1</sup> Shuangqin Yan,<sup>3</sup>  
Yeqing Xu,<sup>3</sup> Sanhuan Huang,<sup>1</sup>  
Jiahu Hao,<sup>1,2</sup> Peng Zhu,<sup>1,2</sup> Ying Niu,<sup>1</sup>  
Shilu Tong,<sup>2,4</sup> and Fangbiao Tao<sup>1,2</sup>

Folic acid (FA) is well known for its preventive effect on neural tube defects, and recommendations for FA supplement before and during pregnancy have been well established all over the world. Since increased consumption of FA has been prevalent, concerns about its negative consequences have been raised. Diverse adverse effects of FA have been identified, ranging from an increased risk of oral cleft malformations, asthma/respiratory tract infections, and small-for-gestational-age infants to the more serious effect of a potential increased risk of cancers.

Previous studies have suggested that a high maternal folate concentration is associated with an increased risk of insulin resistance in children and gestational diabetes mellitus (GDM) (1). In addition, animal experiments have reported that maternal high-FA supplement promotes glucose intolerance and insulin resistance in male mouse offspring fed a high-fat diet (2). However, whether FA supplement consumption preconception and during pregnancy increases the risk of GDM has not yet been investigated. We used data from a prospective cohort study to explore the association between FA supplement consumption and risk of GDM.

The China-Anhui Birth Cohort Study (C-ABCS) is a population-based study that recruited pregnant women ( $n = 3,474$ )

from Ma'anshan City in the Anhui province of China between May 2013 and September 2014. Women underwent a 75-g oral glucose tolerance test at approximately 28 weeks of gestation, and diagnosis of GDM was made when any of the following criteria were met: fasting plasma glucose  $\geq 5.1$  mmol/L, at 1 h  $\geq 10$  mmol/L, and at 2 h  $\geq 8.5$  mmol/L, according to current guidelines from the American Diabetes Association. We used multivariable logistic regression to evaluate the effect of FA supplement consumption on risk of GDM and further assessed the interactions between prepregnancy BMI, a family history of diabetes, and FA supplement consumption. Only women who had either used FA supplements or never used any vitamin supplements ( $n = 1,938$ ) were included in the analysis. GDM was diagnosed in 249 of the 1,938 (12.8%) women. We found that daily FA supplement consumption in the first trimester was associated with an increased risk of GDM (adjusted odds ratio [OR] 2.25 [95% CI 1.35–3.76]). Women with a prepregnancy BMI  $\geq 25$  kg/m<sup>2</sup> and taking FA supplements daily in the first trimester had a much higher risk of GDM (OR 5.63 [95% CI 2.77–11.46]) compared with women with a prepregnancy BMI  $< 25$  kg/m<sup>2</sup> and not taking any FA supplements. An increased risk of GDM was not

apparent for women using FA before pregnancy alone or in the second trimester alone.

The underlying mechanism of this adverse effect is unclear. There are at least two possible explanations. The first may be an imbalance between vitamin B<sub>12</sub> and folate. High folate status could exaggerate the metabolic effects of vitamin B<sub>12</sub> deficiency (3) and might participate in the pathogenesis of GDM through worsening insulin resistance. The second possible explanation may be the harmful effects of unmetabolized plasma FA. Unmetabolized plasma FA has been reported to be related to decreased natural killer cell cytotoxicity (4), which has also been suggested to be involved in the pathogenesis of GDM (5).

In conclusion, our study, for the first time, suggests that daily FA supplement consumption in early pregnancy increases the risk of GDM, and further larger cohort studies are warranted to examine this adverse effect.

**Funding.** This work is supported by National Natural Science Foundation of China (NSCF-81330068 and NSCF-81573168).

**Duality of Interest.** No potential conflicts of interest relevant to this article were reported.

**Author Contributions.** B.Z. contributed to the acquisition of data, statistical analysis and interpretation of data, and manuscript writing. X.G., L.M., S.H., and Y.N. recruited participants and collected questionnaires. K.H. contributed to

<sup>1</sup>Department of Maternal, Child and Adolescent Health, Anhui Medical University, Hefei, China

<sup>2</sup>Anhui Provincial Key Laboratory of Population Health & Aristogenics, Anhui Medical University, Hefei, China

<sup>3</sup>Ma'anshan Maternity and Child Health Care Hospital, Ma'anshan, China

<sup>4</sup>School of Public Health and Social Work and Institute of Health and Biomedical Innovation, Queensland University of Technology, Kelvin Grove, Australia

Corresponding author: Fangbiao Tao, [fbaotao@ahmu.edu.cn](mailto:fbaotao@ahmu.edu.cn).

Received 4 November 2015 and accepted 19 November 2015.

© 2016 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered.

sample and data management. S.Y. and Y.X. collected samples. J.H. and P.Z. interpreted the results. S.T. revised the manuscript. F.T. contributed to the conception and design of the study and study supervision. F.T. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

## References

1. Krishnaveni GV, Hill JC, Veena SR, et al. Low plasma vitamin B12 in pregnancy is associated with gestational 'diabesity' and later diabetes. *Diabetologia* 2009;52:2350–2358
2. Huang Y, He Y, Sun X, He Y, Li Y, Sun C. Maternal high folic acid supplement promotes glucose intolerance and insulin resistance in male mouse offspring fed a high-fat diet. *Int J Mol Sci* 2014;15:6298–6313
3. Selhub J, Morris MS, Jacques PF. In vitamin B12 deficiency, higher serum folate is associated with increased total homocysteine and methylmalonic acid concentrations. *Proc Natl Acad Sci U S A* 2007;104:19995–20000
4. Troen AM, Mitchell B, Sorensen B, et al. Un-metabolized folic acid in plasma is associated with reduced natural killer cell cytotoxicity among postmenopausal women. *J Nutr* 2006; 136:189–194
5. Zhao YH, Wang DP, Zhang LL, Zhang F, Wang DM, Zhang WY. Genomic expression profiles of blood and placenta reveal significant immune-related pathways and categories in Chinese women with gestational diabetes mellitus. *Diabet Med* 2011;28:237–246