



Measuring the Stigma Surrounding Type 2 Diabetes: Development and Validation of the Type 2 Diabetes Stigma Assessment Scale (DSAS-2)

Diabetes Care 2016;39:2141–2148 | DOI: 10.2337/dc16-0117

Jessica L. Browne,^{1,2} Adriana D. Ventura,^{1,2}
Kylie Mosely,^{2,3} and Jane Speight^{1,2,4}

OBJECTIVE

To develop and validate a self-report measure of perceived and experienced stigma for use with adults with type 2 diabetes: the Type 2 Diabetes Stigma Assessment Scale (DSAS-2).

RESEARCH DESIGN AND METHODS

An item pool was drafted based on qualitative data from 25 adults with type 2 diabetes and content from other health-related stigma questionnaires. Thirteen adults with type 2 diabetes completed 57 draft diabetes stigma items and participated in cognitive debriefing interviews. Based on participant feedback, the pool was reduced to 48 items with a 5-point Likert scale (strongly disagree to strongly agree). A total of 1,064 adults with type 2 diabetes completed a survey including these 48 items and other validated measures. Data were subject to principal components analysis and Spearman ρ correlations.

RESULTS

The scale was reduced to 19 items, with an unforced three-factor solution indicative of three subscales: Treated Differently (6 items, $\alpha = 0.88$), Blame and Judgment (7 items, $\alpha = 0.90$), and Self-stigma (6 items, $\alpha = 0.90$). A forced one-factor solution supported the calculation of a total score ($\alpha = 0.95$). Satisfactory concurrent, convergent, and discriminant validity were demonstrated.

CONCLUSIONS

The 19-item DSAS-2 is a reliable and valid measure of type 2 diabetes stigma. A rigorous design and validation process has resulted in a relatively brief measure of perceived and experienced stigma in type 2 diabetes. The novel scale has satisfactory psychometric properties and is now available to facilitate much-needed research in this field.

For the person with type 2 diabetes, managing the condition is a relentless activity. This not only places an emotional burden on the individual (1) but may also invite negative attention, comment, and judgment from others, suggesting a socioemotional burden. Whereas our previous review demonstrates that people without diabetes do not perceive the condition as stigmatized (2), there is now a reasonable body of evidence to indicate that people with type 2 diabetes perceive and experience stigma related to their condition. Qualitative research has provided a rich description of diabetes stigma from the perspective of adults with type 2 diabetes

¹School of Psychology, Deakin University, Geelong, Australia

²The Australian Centre for Behavioural Research in Diabetes, Diabetes Victoria, Melbourne, Australia

³School of Psychology, Australian Catholic University, Sydney, Australia

⁴AHP Research, Hornchurch, U.K.

Corresponding author: Jessica L. Browne, jlbrowne@acbrd.org.au.

Received 19 January 2016 and accepted 10 July 2016.

© 2016 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <http://www.diabetesjournals.org/content/license>.

See accompanying articles, pp. 2122, 2126, 2149, 2158, 2165, 2174, 2182, 2190, and 2197.

(3–6). They feel blamed and judged by others (e.g., for “bringing it on themselves”) and report being subject to negative stereotyping (e.g., fat, lazy), leading to exclusion, rejection, discrimination, and restricted/lost life opportunities (6).

The multinational Diabetes Attitudes, Wishes, and Needs second study (DAWN-2) corroborates these qualitative findings, with one in five (19.2%) respondents reporting being discriminated against (7). However, in this study, discrimination (just one specific manifestation of stigmatization) was assessed using a single item, and so we remain unable to quantify the full extent and experience of diabetes stigma. The International Diabetes Federation has identified diabetes stigma as a problem that needs urgent attention, with a key priority being to “champion a world free from discrimination and stigma for people with diabetes” (8). However, before we can fully address diabetes stigma at the individual, environmental, and societal levels, we need research to investigate the following: the extent and severity of the problem, its determinants and consequences (e.g., for emotional well-being, diabetes self-management, and clinical outcomes), and changes over time or in response to intervention (9).

Rigorous research cannot take place without a validated, self-report measure of diabetes stigma. Previous studies have used generic measures of health-related stigma (10,11) or unvalidated diabetes-specific questions (7,12). Although valid and appropriate for use in some contexts, generic measures do not capture the nuances of negative social judgment and stereotyping that are associated specifically with type 2 diabetes, and some include content that is irrelevant or inappropriate for people with type 2 diabetes (e.g., about fear of touching due to contagion). Study-specific or ad hoc approaches to quantifying diabetes stigma are typically unsystematic and unvalidated and make comparisons across studies impossible due to variations in operationalization and measurement. Thus, our aim was to develop and validate a self-report measure of perceived and experienced stigma for use with people with type 2 diabetes: the Type 2 Diabetes Stigma Assessment Scale (DSAS-2).

RESEARCH DESIGN AND METHODS

All study activities were approved by the Deakin University Human Research Ethics Committee (2011-046).

Study 1: Design and Debriefing of Questionnaire

Item Development

The content of the questionnaire was informed by qualitative data about diabetes stigma from 25 adults with type 2 diabetes, published elsewhere (6). One author drafted an initial 125-item pool based on participant quotes from interview transcripts, representing all key themes in the data. The items were constructed as statements with a 5-point Likert response scale (strongly disagree to strongly agree). An additional 30 items were drafted based on items used in health-related stigma scales (e.g., Weight Self-Stigma Scale [13] and Chronic Illness Anticipated Stigma Scale [11]).

The 155-item pool was reduced to 91 items through review by all authors, with consensus about item redundancy and revisions reached through discussion. Four external experts (in health psychology, health stigma, and/or diabetes) were invited to review the 91 items. Feedback was collated and reviewed by all authors, who made the final decisions about item revisions and deletions. A total of 57 items were taken forward into the cognitive debriefing phase.

Cognitive Debriefing Interviews

Recruitment and Procedure. Cognitive debriefing interviews were conducted to assess the clarity, relevance, and comprehensiveness of the questionnaire instructions, items, and response options from the perspective of adults with type 2 diabetes. Participants were recruited through advertisements in diabetes media (e.g., e-newsletters) and social media (e.g., Facebook pages). Upon making contact with researchers, potential participants were screened for eligibility (self-reported type 2 diabetes diagnosis, aged 18–75 years, fluent in English, able to access/use Internet, and living in Australia) and asked to provide written informed consent. Then a telephone interview was scheduled. Participants were then directed to a website and requested to complete an online version of the stigma questionnaire while making notes for discussion on a paper copy (sent to them by post), in the 24 h

prior to their scheduled interview. Interviews lasted an average of 28 min (range 15–45). Participants were entered into a prize draw to win a year’s supply of (12) movie tickets.

Study 2: Scale Reduction and Psychometric Validation of DSAS-2

Procedure

An online survey was conducted to collect response data on the type 2 diabetes stigma item pool, and a selection of previously validated questionnaires (see MATERIALS). The website featured a plain language description of the study, a consent form, eligibility screening, and the online survey (securely hosted by Qualtrics, Provo, UT). Participants who did not have access to the internet could contact researchers to request a hard-copy survey.

Participants and Recruitment

Eligible participants were English-speaking adults with type 2 diabetes, aged 18–75 years, living in Australia. An invitation letter was mailed to a random sample of 12,000 National Diabetes Services Scheme (NDSS) registrants with type 2 diabetes, stratified by insulin use (50% insulin treated, intentionally oversampled) and by state of residence. The letter introduced the study and directed people to the study website. The study was also advertised nationwide in diabetes-related media (e.g., e-newsletters and social media). Twenty-five people requested and returned hard copy versions of the consent form and survey. Participants were entered into a prize draw to win one of three iPad Minis.

Materials

The survey included the diabetes stigma item pool as well as a selection of psychosocial measures for validation purposes and questions about demographic and clinical characteristics. Table 1 summarizes the types of validity assessed and measures used.

Concurrent Validity. The Stigma Scale for Chronic Illness 8-item scale (SSCI-8) (14), a generic scale that assesses internalized and enacted stigma for chronic conditions, was used to assess concurrent validity. It comprises eight statements (e.g., “Because of my condition, people were unkind to me”) and a 5-point Likert response scale (1 = never to 5 = always). Item scores are summed to form a total score (range 8–40), where

Table 1—Summary of validity assessments used in the current study

Validity	Definition	Measure
Concurrent validity	Concurrent validity is confirmed if the correlation between the new measure and an established measure for the same/similar construct is at least moderate ($r > 0.4$)	SSCI-8 to assess perceived and enacted stigma for chronic conditions
Convergent validity	Convergent validity is confirmed if the correlation between two measures thought to be related to each other is at least moderate ($r > 0.4$)	PHQ-8 to assess depressive symptoms GAD-7 to assess anxiety symptoms PAID to assess diabetes distress RSE to assess self-esteem
Discriminant validity	Discriminant validity is confirmed when a weak (or nonexistent) correlation ($r < 0.3$) is demonstrated with other constructs thought to bear little or no relationship with the construct at hand	Diabetes duration PRMQ to assess memory

higher scores indicate more internalized/enacted stigma. Positive correlations were expected with the DSAS-2.

Convergent Validity. Convergent validity was assessed against the following.

Patient Health Questionnaire 8-Item Scale and Generalized Anxiety Disorder 7-Item Scale. The Patient Health Questionnaire 8-item scale (PHQ-8) and Generalized Anxiety Disorder 7-item scale (GAD-7) (15,16) assess depressive and anxiety symptomatology, respectively, by asking respondents to indicate how frequently they have experienced eight symptoms of depression (e.g., “little interest or pleasure in doing things”) and seven symptoms of anxiety (e.g., “feeling afraid as if something awful might happen”) over the last 2 weeks. For both scales, items are scored on a 4-point Likert scale (0 = not at all to 3 = nearly every day) and summed to produce a total score (PHQ-8 range 0–24; GAD-7 range 0–21). Higher scores indicate more depressive/anxiety symptoms. Positive correlations were expected with the DSAS-2.

Problem Areas in Diabetes Scale. Problem Areas in Diabetes (PAID) scale (17) is a 20-item measure of diabetes-specific distress. Respondents indicate how much of a problem each statement (e.g., “feeling ‘burned out’ by the constant effort needed to manage diabetes”) is for them on a 5-point Likert scale (0 = not a problem to 4 = serious problem). Items are summed and converted to a percentage score, with higher scores indicating greater diabetes distress. A positive correlation was expected with the DSAS-2.

Rosenberg Self-esteem Scale. Rosenberg Self-esteem scale (RSE) (18) is a 10-item measure of self-esteem. Respondents rate statements (e.g., “I take a positive

attitude toward myself”) on a 4-point Guttman scale (0 = strongly disagree to 3 = strongly agree). A total score (range 0–30) is calculated by summing responses to individual items, after reverse scoring negatively worded items. Higher scores indicate higher self-esteem. A negative correlation was expected with the DSAS-2.

Discriminant Validity. Discriminant validity was assessed against diabetes duration (years) and scores on the Prospective and Retrospective Memory Questionnaire (PRMQ) (19). The PRMQ assesses general prospective and retrospective memory. Respondents indicate how frequently they experience 16 different memory “slips” (e.g., “Do you fail to mention or give something to a visitor that you were asked to pass on?”) by responding on a 5-point Likert scale (1 = never to 5 = very often). Responses are summed to produce a total score (range 16–80); higher scores indicate more memory problems.

All the validated scales demonstrated high internal consistency reliability in the current sample (all $\alpha \geq 0.90$, range 0.90–0.96). The measures were ordered in the survey in such a way as to minimize priming 1) clinical characteristics, 2) PHQ-8, 3) GAD-7, 4) PAID, 5) diabetes stigma items, 6) SSCI-8, 7) RSE, and 8) demographics.

Data Preparation and Analyses

Kolmogorov-Smirnov tests demonstrated that all scale data had nonnormal distributions, necessitating the use of nonparametric statistics.

Data on all scales were not missing completely at random (Little MCAR test, all $P < 0.05$); however, missing data patterns did not differ by any key demographic, clinical, or psychosocial factors,

suggesting that imputation was appropriate. Where respondents had $\leq 10\%$ of data points missing on a measure, their missing data were imputed using expectation maximization techniques (20) to facilitate calculation of a total score. Total scores were not calculated when $>10\%$ of data points were missing. No missing data were tolerated for the calculation of subscale scores. High overall completion rates ($\geq 90\%$) were taken as evidence of acceptability.

Prior to proceeding with dimension reduction analyses, items with a large number (five or more) of very high ($r_s > 0.7$) or very low ($r_s < 0.3$) inter-item correlations were screened out using two-tailed inter-item Spearman ρ correlations. Participants with missing data were deleted pairwise, minimizing loss of data by using all available cases for each analysis.

A Kaiser-Meyer-Olkin statistic of 0.98 indicates that our sample size was more than adequate to proceed with dimension reduction analyses (21). The structural validity of the questionnaire was assessed using principal components analysis with direct oblimin rotation. Decisions about how many factors to retain were informed by Kaiser eigenvalue (≥ 1), percentage variance explained by each factor, and factor loadings. Modern guidelines suggest that multiple factor loadings of ≥ 0.5 indicate likely strong and stable factors, and thus this cutoff was used to identify meaningful loadings (22). Cronbach α was used to assess internal consistency reliability of factors (subscales) and a forced one-factor solution (total scale).

Item scores were summed to produce subscale and scale total scores. Participant response patterns on the final scale were examined using descriptive statistics.

Concurrent, convergent, and discriminant validity were assessed using two-tailed Spearman ρ correlations. Participants with missing total scores were deleted pairwise, minimizing loss of data by using all available cases for each analysis. In accordance with Cohen's guidelines, medium-to-large correlations ($r_s > \pm 0.4$) were taken as evidence for concurrent and convergent validity, and small correlations ($r_s < \pm 0.3$) were taken as evidence of discriminant validity (23).

All statistical analyses were conducted using IBM SPSS for Mac, version 22 (Chicago, IL). An α level of $P < 0.05$ was taken to indicate significance.

RESULTS

Study 1: Design and Debriefing of Questionnaire

Thirteen adults with type 2 diabetes (median age 60 years [range 39–75]) took part in cognitive debriefing interviews. All had at least some high school education; a minority ($n = 4$, 31%) had a university degree; two (15%) were born outside Australia; and five (38%) participants used insulin.

Participants indicated that the questionnaire items were easy to understand, relevant, and comprehensive. Participants demonstrated a clear and accurate understanding of the word "stigma." Iterative changes to the questionnaire were made throughout the cognitive debriefing phase if consistent feedback was received from three or more participants. Examples include the following: deletion of redundant items, changing the middle response option from "neither disagree nor agree" to "unsure," and adding additional questionnaire instructions.

Where feedback was inconsistent between participants, no immediate action was taken, and instead all authors reviewed these outstanding issues and achieved consensus regarding revisions through discussion. This process resulted in a reduced and revised type 2 diabetes stigma item pool of 48 items (down from 57), each with a 5-point Likert response scale (1 = strongly disagree, 2 = disagree, 3 = unsure, 4 = agree, and 5 = strongly agree).

Study 2: Scale Reduction and Psychometric Validation of DSAS-2 Sample Characteristics

A total of 1,264 eligible adults with type 2 diabetes participated. Of those,

1,064 completed the survey section that included the diabetes stigma items ($n = 200$ discontinued prior to this). Thus, analyses were performed on this sample of $n = 1,064$ and their characteristics are displayed in Table 2.

Acceptability

A total of 989 (93.0%) respondents completed all 48 diabetes stigma items (i.e., no missing data), indicating a high level of acceptability, especially given the large number of items and relatively sensitive content. Forty-three (4.0%)

respondents had one missing data point, 11 (1.0%) had two missing data points, and 21 (2.0%) had more than two missing data points (range 3–48).

Scale Reduction

Very high multicollinearity was observed among the 48 items (determinant = 7.00^{e-19}). Screening of inter-item correlations resulted in deletion of eight items (three due to high correlations and five due to low correlations).

An unforced principal components analysis performed on the remaining

Table 2—Sample characteristics (n = 1,064)

	Mean \pm SD (range) or median (quartile 1, quartile 3) or n (%)*
Demographic and clinical characteristics	
Women	458 (43.0)
Age (years)	61.20 \pm 9.40 (22–75)
Diabetes duration (years)	11.12 \pm 7.59 (0–44)
Primary treatment	
Insulin pump	2 (0.2)
Insulin injections	462 (43.2)
Noninsulin injectable	33 (3.1)
Oral hypoglycemic agents only	427 (40.1)
Lifestyle only	142 (13.3)
Geographic area	
Metropolitan	619 (58.3)
Regional	254 (23.9)
Rural	189 (17.8)
Relationship status: single	288 (27.1)
Living situation: alone	192 (18.0)
Employment	
Paid employment	385 (36.2)
Retired	502 (47.2)
Other	117 (16.6)
Household income (\$AUD)	
$\leq 20,000$	188 (18.0)
20,001–40,000	240 (22.9)
40,001–60,000	164 (15.7)
60,001–100,000	151 (14.4)
100,001–150,000	99 (9.5)
>150,000	59 (5.6)
Don't know/prefer not to say	146 (13.9)
Education	
Some high school	287 (27.0)
Completed high school	122 (11.5)
Vocational education/diploma	322 (30.3)
University, undergraduate	185 (17.4)
University, postgraduate	146 (13.7)
Born in Australia	745 (70.0)
Main language spoken at home: English	1,025 (96.4)
Aboriginal or Torres Strait Islander	20 (1.9)
Psychosocial characteristics	
Depressive symptoms: PHQ-8	6.61 \pm 5.70 (0–24)
	5.00 (2.00, 10.00)
Anxiety symptoms: GAD-7	4.71 \pm 5.04 (0–21)
	3.00 (1.00, 7.00)
Diabetes-specific distress: PAID	19.07 \pm 19.57 (0–100)
	12.50 (3.75, 27.50)
Self-esteem: RSE	20.59 \pm 5.70 (3–30)
	20.00 (17.00, 25.00)

*Frequencies do not always add to total $n = 1,064$ due to missing data on some items.

40 items revealed four factors (eigenvalue ≥ 1). However, the fourth factor had a low eigenvalue (1.14), explained little variance (2.9%), and comprised just one item. A forced three-factor solution revealed three potential subscales best described as Treated Differently (11 items), Blame and Judgment (12 items), and Self-stigma (7 items). Ten items did not load on any factor using the ≥ 0.5 criteria and were therefore excluded from further analyses.

Cronbach α coefficients for the three subscales were high (all ≥ 0.92), indicating potential redundancy, so further scale reduction was undertaken by reviewing item content, inter-item and item-total correlations, and α coefficients. Scale reduction decisions for each subscale were as follows:

Treated Differently: Five items were removed (two due to redundant content and numerous high inter-item correlations and three due to content not being well aligned with other items in the subscale).

Blame and Judgment: Six items were removed (three due to redundancy and multiple, very high inter-item and item-total correlations and three that asked participants to compare stigma associated with type 1 and type 2 diabetes on the basis of concerns that they may serve to perpetuate negative feelings/attitudes toward other groups). One item (previously dropped from the scale on the basis that it did not load ≥ 0.5) was reinstated, as it uniquely represented the experience of feeling blamed for “bringing diabetes on myself,” a key theme identified from the qualitative interviews (6).

Self-stigma: One item was removed because it was almost identical to another item but used more complicated language.

An unforced three-factor solution comprising the final 19 items is presented in Table 3, along with a forced one-factor solution, supporting calculation of a total scale score. In the three-factor solution, one item (“I feel embarrassed in social situations because of my type 2 diabetes”) double loaded on the Treated Differently and Self-stigma factors. However, it loaded on the latter marginally more

strongly, and the wording was considered more relevant to the self-stigma construct; thus, it was included in the scoring of the latter subscale only.

The three-factor and one-factor solutions were stable, with findings being replicated on two randomly selected subsamples ($n = 518$ and $n = 546$).

The final scale had a Flesch-Kincaid reading ease score of 69.7, suggesting that the scale is constructed in plain English that is readable for people with some secondary school education (8th to 9th grade reading level).

Response Patterns and Descriptive Statistics

The full range of response options was used for every item, although for most, the distribution of responses was skewed positively (Table 4). Missing data ranged from 1.2 to 1.9% per item.

The mean \pm SD Total Diabetes Stigma score was 41.0 ± 15.9 (range 19–90); median was 40.0 (27, 52). Subscale scores were as follows: Treated Differently: 11.0 ± 4.8 (range 6–30), 10 (6, 14); Blame and Judgment: 18.1 ± 7.2 (range 7–35), 18 (12, 24); and Self-stigma: 11.9 ± 5.8 (range 6–30), 11 (7, 15). A total of 205 (19.3%) respondents scored more than one SD above the mean total diabetes stigma score, suggestive of potentially problematic perceived and experienced diabetes stigma.

Validity of the DSAS-2

Satisfactory concurrent validity was demonstrated by a large positive correlation between the DSAS-2 and the SSCI-8 ($r_s = 0.78$, $P < 0.001$). Convergent validity was confirmed with a strong positive correlation with the PAID ($r_s = 0.67$, $P < 0.001$), medium positive associations with the PHQ-8 and GAD-7 ($r_s = 0.44$ and 0.46 , respectively, both $P < 0.001$), and a medium negative correlation with the RSE ($r_s = -0.51$, $P < 0.001$). Discriminant validity was confirmed as the DSAS-2 had weak(er) associations with the PRMQ ($r_s = 0.28$, $P < 0.001$) and diabetes duration ($r_s = -0.04$, NS).

CONCLUSIONS

The DSAS-2 meets the need for a self-report measure of type 2 diabetes stigma, which can now be used in research to quantify the extent and impact of stigma, as perceived and experienced by adults with type 2 diabetes. This

novel measure has a comprehensive yet simple three-factor structure, which encapsulates both enacted and self-stigma, consistent with established theoretical perspectives (2,9,24,25). Importantly, the measure can also be reduced to a single dimension, facilitating the calculation of a total scale score. Our scale development and validation process has resulted in an acceptable, relatively brief, and usable measure of diabetes stigma, with strong psychometric properties.

Small proportions of missing data and full use of response options point to the acceptability of the scale. The 19-item scale is comparable in length to other widely used diabetes-specific measures, such as the PAID (17) and the Insulin Treatment Appraisal Scale (26). The DSAS-2 scale and subscales all have high internal consistency reliability, and the measure has good concurrent, convergent, and discriminant validity. Although the discriminant validity analyses resulted in a statistically significant correlation between the DSAS-2 and PRMQ scores (likely due to the large sample size), the effect size was small, indicating a negligible relationship between the variables. As diabetes stigma is likely to have pervasive cognitive, emotional, and social impacts, we anticipated that many constructs would have some relationship, however small, with diabetes stigma. Diabetes duration and the PRMQ were considered good candidates for assessing discriminant validity of the DSAS-2 because we made a priori predictions that these relationships would be small, negligible, or indirect; thus discriminant validity of the DSAS-2 was confirmed (27).

The Cronbach α for the full 19-item scale is high (0.95), suggesting further scale reduction might be possible. However, we believe that further item deletion will reduce the comprehensiveness and face validity of the scale, and reduce the internal consistency reliability of the subscales.

The item “I feel embarrassed in social situations because of my type 2 diabetes” double loaded on the Treated Differently and Self-stigma factors (but was only used for calculation of the latter subscale). For future use, we recommend removing the phrase “in social situations” from this item, as this will further differentiate it from the concept

Table 3—Structure and internal consistency reliability of the 19-item DSAS-2

Item wording	Unforced three-factor solution*			Forced one-factor solution*
	Treated Differently	Blame and Judgment	Self-stigma	
Some people think I cannot fulfill my responsibilities (e.g., work, family) because I have type 2 diabetes	0.73	—	—	0.69
Some people treat me like I'm "sick" or "ill" because I have type 2 diabetes	0.65	—	—	0.70
Some people see me as a lesser person because I have type 2 diabetes	0.51	—	—	0.84
Some people exclude me from social occasions that involve food/drink they think I shouldn't have	0.73	—	—	0.70
I have been discriminated against in the workplace because of my type 2 diabetes	0.73	—	—	0.65
I have been rejected by others (e.g., friends, colleagues, romantic partners) because of my type 2 diabetes	0.74	—	—	0.68
I have been told that I brought my type 2 diabetes on myself	—	0.64	—	0.62
There is blame and shame surrounding type 2 diabetes	—	0.74	—	0.77
Because I have type 2 diabetes, some people judge me for my food choices	—	0.63	—	0.75
Health professionals think that people with type 2 diabetes don't know how to take care of themselves	—	0.78	—	0.65
Because of my type 2 diabetes, health professionals have made negative judgments about me	—	0.70	—	0.74
There is a negative stigma about type 2 diabetes being a "lifestyle disease"	—	0.82	—	0.73
Because I have type 2 diabetes, some people assume I must be overweight, or have been in the past	—	0.80	—	0.73
I feel embarrassed in social situations because of my type 2 diabetes†	0.53†	—	0.56	0.73
I'm ashamed of having type 2 diabetes	—	—	0.82	0.72
I blame myself for having type 2 diabetes	—	—	0.77	0.57
Because I have type 2 diabetes, I feel like I am not good enough	—	—	0.62	0.79
Having type 2 diabetes makes me feel like a failure	—	—	0.70	0.80
I feel guilty for having type 2 diabetes	—	—	0.74	0.80
Eigenvalue	9.85	1.40	1.33	N/A
% variance explained	51.8	7.4	7.0	
Cumulative % variance explained		66.1		
Cronbach α	0.88	0.90	0.90	0.95
Determinant		0.000003		

*Factor loadings (principal components analysis); factor loadings <0.5 suppressed. For the three-factor solution, the rotation converged in 14 iterations. †Not included in the scoring of the treated differently subscale. For future use, we recommend removing the phrase "in social situations" from this item.

of being treated differently by others and will keep the focus on self-perception.

A key strength of this study is the rigorous process used to design, debrief, and validate the DSAS-2. Content for the questionnaire was developed directly from the themes that emerged from a qualitative study of stigma with 25 adults with type 2 diabetes (6). The stigma item pool was debriefed in full with a further 13 people with type 2 diabetes, and as such, adults with type 2 diabetes were fully engaged as key informants during questionnaire development. Survey participants constituted

a national sample of adults with type 2 diabetes, and a wide range of ages and socioeconomic backgrounds were represented.

It is likely that the sample constitutes a representative spread of views about diabetes stigma among Australians with type 1 or type 2 diabetes, as substantial efforts were made to reduce recruitment bias in this regard. The true purpose of the study (i.e., "diabetes stigma") was not disclosed but rather it was advertised as a survey about what it is like to live with diabetes. Furthermore, the full survey included many other questions about

psychosocial aspects of diabetes, so the advertised purpose was not misleading.

This study also has a number of limitations. Although the primary recruitment method for study 2 involved sending invitations to a national random sample of NDSS registrants, participation was voluntary and thus the sample (while large) is self-selected and potentially biased, although not in relation to their experience of social stigma as the true objective was concealed during recruitment. The sample may be more likely to have high health literacy or be more engaged in issues of diabetes

Table 4—Response patterns by item for the 19-item DSAS-2

Subscale and item wording	Response options					Missing data
	Strongly disagree	Disagree	Unsure	Agree	Strongly agree	
Treated differently						
Some people think I cannot fulfill my responsibilities (e.g., work, family) because I have type 2 diabetes	512 (48.1)	323 (30.4)	105 (9.9)	90 (8.5)	20 (1.9)	14 (1.3)
Some people treat me like I'm "sick" or "ill" because I have type 2 diabetes	436 (41.0)	336 (31.6)	109 (10.2)	149 (14.0)	17 (1.6)	17 (1.6)
Some people see me as a lesser person because I have type 2 diabetes	446 (41.9)	306 (28.8)	156 (14.7)	114 (10.7)	24 (2.3)	18 (1.7)
Some people exclude me from social occasions that involve food/drink they think I shouldn't have	531 (49.9)	352 (33.1)	94 (8.8)	54 (5.1)	13 (1.2)	20 (1.9)
I have been discriminated against in the workplace because of my type 2 diabetes	523 (49.2)	351 (33.0)	111 (10.4)	41 (3.9)	19 (1.8)	19 (1.8)
I have been rejected by others (e.g., friends, colleagues, romantic partners) because of my type 2 diabetes	555 (52.2)	363 (34.1)	75 (7.0)	42 (3.9)	12 (1.1)	17 (1.6)
Blame and judgment						
I have been told that I brought my type 2 diabetes on myself	423 (39.8)	255 (24.0)	101 (9.5)	217 (20.4)	55 (5.2)	13 (1.2)
There is blame and shame surrounding type 2 diabetes	316 (29.7)	250 (23.5)	137 (12.9)	271 (25.5)	77 (7.2)	13 (1.2)
Because I have type 2 diabetes, some people judge me for my food choices	293 (27.5)	252 (23.7)	113 (10.6)	305 (28.7)	85 (8.0)	16 (1.5)
Health professionals think that people with type 2 diabetes don't know how to take care of themselves	302 (28.4)	297 (27.9)	195 (18.3)	202 (19.0)	51 (4.8)	17 (1.6)
Because of my type 2 diabetes, health professionals have made negative judgments about me	379 (35.6)	299 (28.1)	178 (16.7)	149 (14.0)	43 (4.0)	16 (1.5)
There is a negative stigma about type 2 diabetes being a "lifestyle disease"	245 (23.0)	196 (18.4)	134 (12.6)	353 (33.2)	121 (11.4)	15 (1.4)
Because I have type 2 diabetes, some people assume I must be overweight, or have been in the past	218 (20.5)	155 (14.6)	148 (13.9)	404 (38.0)	124 (11.7)	15 (1.4)
Self-stigma						
I feel embarrassed in social situations because of my type 2 diabetes	566 (53.2)	302 (28.4)	65 (6.1)	95 (8.9)	22 (2.1)	14 (1.3)
I'm ashamed of having type 2 diabetes	612 (57.5)	255 (24.0)	54 (5.1)	92 (8.6)	36 (3.4)	15 (1.4)
I blame myself for having type 2 diabetes	413 (38.8)	209 (19.6)	97 (9.1)	250 (23.5)	75 (7.0)	20 (1.9)
Because I have type 2 diabetes, I feel like I am not good enough	530 (49.8)	303 (28.5)	87 (8.2)	101 (9.5)	26 (2.4)	17 (1.6)
Having type 2 diabetes makes me feel like a failure	478 (44.9)	331 (31.1)	85 (8.0)	124 (11.7)	31 (2.9)	15 (1.4)
I feel guilty for having type 2 diabetes	405 (38.1)	318 (29.9)	81 (7.6)	188 (17.7)	58 (5.5)	14 (1.3)

Data presented are *n* (%). Within rows, percentages do not always total 100 due to rounding.

management than the general population of Australian adults with diabetes. The sample appears mostly representative of Australian adults with type 2 diabetes, although the proportion who use insulin is higher here (42 vs. 24% [28]) due to intentional oversampling. The questionnaire was developed and made available in English only, due to funding constraints, which may explain the relative lack of ethnic diversity in the sample. Finally, we were unable to assess responsiveness or predictive validity in the current study, and these aspects of psychometric validation will need to be established in future work.

The next steps in our research program will be to 1) quantitatively assess the relationships between diabetes stigma and other outcomes of interest, e.g., psychological well-being and

self-management, and 2) develop a comparable self-report measure of perceived and experienced stigma for adults with type 1 diabetes (work ongoing). Future research will need to confirm the psychometric properties of the DSAS-2 in other samples. As the DSAS-2 has been developed and validated with an Australian sample, it will be important to debrief the measure with English-speaking participants from other countries and cultures and for appropriate cultural adaptation and translation techniques to be used to enable use in non-English-speaking populations.

Our findings indicate that the DSAS-2 is a reliable and valid measure of perceived and experienced stigma for use with adults with type 2 diabetes. The DSAS-2 fulfills the need for a validated, diabetes-specific measure of stigma and will facilitate new

research in the field to further our understanding of the determinants, extent, and impact of diabetes stigma.

Acknowledgments. The authors thank Prof. Jorg Huber (University of Brighton), Dr. Giesje Nefs (Tilburg University), Prof. Frans Pouwer (Tilburg University), and Dr. Rebecca Puhl (University of Connecticut Rudd Center for Food Policy and Obesity) for their expert review of the draft diabetes stigma item pool. The authors also thank Shaira Baptista and Elizabeth Holmes-Truscott (The Australian Centre for Behavioural Research in Diabetes) for their assistance in setting up the online survey and data cleaning. Finally, the authors thank all of the study participants (debriefing and online survey) for volunteering their time, insights, and experiences.

The DSAS-2 is the copyright of The Australian Centre for Behavioural Research in Diabetes, a partnership for better health between Diabetes Victoria and Deakin University. It is available

free of charge to academic researchers, clinicians, and students for use in non-commercially funded research. For commercially funded studies (initiated or sponsored by industry), a license fee will apply. Potential users are advised to email info@acbrd.org.au to enquire about or access the latest version of the questionnaire and scoring guidance.

Funding. The design and development of the questionnaire was funded by a Diabetes Australia Research Trust grant.

Duality of Interest. Enhancement of the sample size for the validation study was enabled by an unrestricted educational grant from Sanofi. No other potential conflicts of interest relevant to this article were reported.

Author Contributions. J.L.B. conceptualized the study, developed the initial diabetes stigma item pool, conducted the data collection and analyses, and wrote the first draft of the manuscript. A.D.V. provided substantive intellectual input into the study protocol, reviewed and revised the DSAS-2, conducted the data collection and analyses, and provided feedback and revisions on the first and subsequent drafts of the manuscript. K.M. provided substantive intellectual input into the study protocol, reviewed and revised the DSAS-2, reviewed the data collection and analyses for the purposes of making decisions about scale reduction, and provided feedback and revisions on the first and subsequent drafts of the manuscript. J.S. conceptualized the study, reviewed and revised the DSAS-2, reviewed the data collection and analyses for the purposes of making decisions about scale reduction, and provided feedback and revisions on the first and subsequent drafts of the manuscript. All authors approved the final submitted manuscript. J.L.B. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Prior Presentation. Parts of this study were presented in poster form at the 76th Scientific Sessions of the American Diabetes Association, New Orleans, LA, 10–14 June 2016.

References

- Ducat L, Philipson LH, Anderson BJ. The mental health comorbidities of diabetes. *JAMA* 2014; 312:691–692
- Schabert J, Browne JL, Mosely K, Speight J. Social stigma in diabetes: a framework to understand a growing problem for an increasing epidemic. *Patient* 2013;6:1–10
- Lawton J, Ahmad N, Hanna L, Douglas M, Hallowell N. 'I can't do any serious exercise': barriers to physical activity amongst people of Pakistani and Indian origin with type 2 diabetes. *Health Educ Res* 2006;21:43–54
- Aikins AdG. Reframing applied disease stigma research: a multilevel analysis of diabetes stigma in Ghana. *J Community Appl Soc Psychol* 2006;16:426–441
- Tak-Ying Shiu A, Kwan JY-M, Wong RY-M. Social stigma as a barrier to diabetes self-management: implications for multi-level interventions. *J Clin Nurs* 2003;12:149–150
- Browne JL, Ventura A, Mosely K, Speight J. 'I call it the blame and shame disease': a qualitative study about perceptions of social stigma surrounding type 2 diabetes. *BMJ Open* 2013;3:e003384
- Nicolucci A, Kovacs Burns K, Holt RIG, et al.; DAWN2 Study Group. Diabetes Attitudes, Wishes and Needs second study (DAWN2™): cross-national benchmarking of diabetes-related psychosocial outcomes for people with diabetes. *Diabet Med* 2013;30:767–777
- International Diabetes Federation. *IDF Strategic Plan 2013–2015*. Brussels, IDF, 2013
- Van Brakel WH. Measuring health-related stigma—a literature review. *Psychol Health Med* 2006;11:307–334
- Kato A, Takada M, Hashimoto H. Reliability and validity of the Japanese version of the self-stigma scale in patients with type 2 diabetes. *Health Qual Life Outcomes* 2014;12:179
- Earnshaw VA, Quinn DM, Kalichman SC, Park CL. Development and psychometric evaluation of the chronic illness anticipated stigma scale. *J Behav Med* 2013;36:270–282
- Della LJ. Exploring diabetes beliefs in at-risk Appalachia. *J Rural Health* 2011;27:3–12
- Lillis J, Luoma JB, Levin ME, Hayes SC. Measuring weight self-stigma: the weight self-stigma questionnaire. *Obesity (Silver Spring)* 2010;18:971–976
- Molina Y, Choi SW, Cella D, Rao D. The stigma scale for chronic illnesses 8-item version (SSCI-8): development, validation and use across neurological conditions. *Int J Behav Med* 2013;20:450–460
- Kroenke K, Strine TW, Spitzer RL, Williams JBW, Berry JT, Mokdad AH. The PHQ-8 as a measure of current depression in the general population. *J Affect Disord* 2009;114:163–173
- Spitzer RL, Kroenke K, Williams JBW, Löwe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med* 2006;166:1092–1097
- Polonsky WH, Anderson BJ, Lohrer PA, et al. Assessment of diabetes-related distress. *Diabetes Care* 1995;18:754–760
- Rosenberg M. *Society and the Adolescent Child*. Princeton, NJ, Princeton University Press, 1965
- Crawford JR, Smith G, Maylor EA, Della Sala S, Logie RH. The Prospective and Retrospective Memory Questionnaire (PRMQ): normative data and latent structure in a large non-clinical sample. *Memory* 2003;11:261–275
- Shrive FM, Stuart H, Quan H, Ghali WA. Dealing with missing data in a multi-question depression scale: a comparison of imputation methods. *BMC Med Res Methodol* 2006;6:57
- Hutcheson GD, Sofroniou N. *The Multivariate Social Scientist: Introductory Statistics Using Generalized Linear Models*. London, SAGE, 1999
- Costello A, Osborne J. Best practices in exploratory factor analysis: four recommendations for getting the most from your analysis. *Pract Assess, Res Eval* 2005;10
- Cohen J. A power primer. *Psychol Bull* 1992; 112:155–159
- Link BG, Phelan JC. Conceptualizing stigma. *Annu Rev Sociol* 2001;27:363–385
- Jacoby A. Felt versus enacted stigma: a concept revisited. Evidence from a study of people with epilepsy in remission. *Soc Sci Med* 1994;38: 269–274
- Snoek FJ, Skovlund SE, Pouwer F. Development and validation of the insulin treatment appraisal scale (ITAS) in patients with type 2 diabetes. *Health Qual Life Outcomes* 2007;5:69
- Furr RM, Bacharach VR. *Psychometrics: An Introduction*. 2nd ed. Thousand Oaks, CA, SAGE, 2014
- National Diabetes Services Scheme. *Type 2 Diabetes Statistics at 30 September 2015*. Canberra, Australia, Diabetes Australia, 2015