



Cost-effectiveness of the 2014 U.S. Preventive Services Task Force (USPSTF) Recommendations for Intensive Behavioral Counseling Interventions for Adults With Cardiovascular Risk Factors

Diabetes Care 2017;40:640–646 | DOI: 10.2337/dc16-1186

Ji Lin,¹ Xiaohui Zhuo,²
Barbara Bardenheier,^{1,3}
Deborah B. Rolka,¹ W. Edward Gregg,¹
Yuling Hong,⁴ Guijing Wang,⁴
Ann Albright,¹ and Ping Zhang¹

OBJECTIVE

In 2014, the U.S. Preventive Services Task Force (USPSTF) recommended behavioral counseling interventions for overweight or obese adults with the following known cardiovascular disease risk factors: impaired fasting glucose (IFG), hypertension, dyslipidemia, or metabolic syndrome. We assessed the long-term cost-effectiveness (CE) of implementing the recommended interventions in the U.S.

RESEARCH DESIGN AND METHODS

We used a disease progression model to simulate the 25-year CE of the USPSTF recommendation for eligible U.S. adults and subgroups defined by a combination of the risk factors. The baseline population was estimated using 2005–2012 National Health and Nutrition Examination Survey (NHANES). The cost and effectiveness of the intervention were obtained from systematic reviews. Incremental CE ratios (ICERs), measured in cost/quality-adjusted life-year (QALY), were used to assess the CE of the intervention compared with no intervention. Future QALYs and costs (reported in 2014 U.S. dollars) were discounted at 3%.

RESULTS

We estimated that ~98 million U.S. adults (44%) would be eligible for the recommended intervention. Compared with no intervention, the ICER of the intervention would be \$13,900/QALY. CE varied widely among subgroups, ranging from a cost saving of \$302 per capita for those who were obese with IFG, hypertension, and dyslipidemia to a cost of \$103,200/QALY in overweight people without these conditions.

CONCLUSIONS

The recommended intervention is cost effective based on the conventional CE threshold. Considerable variation in CE across the recommended subpopulations suggests that prioritization based on risk level would yield larger total health gains per dollar spent.

¹Division of Diabetes Translation, Centers for Disease Control and Prevention, Atlanta, GA

²Merck Research Laboratory, North Wales, PA

³Immunization Safety Office, Centers for Disease Control and Prevention, Atlanta, GA

⁴Division for Heart Disease and Stroke Prevention, Centers for Disease Control and Prevention, Atlanta, GA

Corresponding author: Ji Lin, xhi6@cdc.gov.

Received 1 June 2016 and accepted 28 January 2017.

This article contains Supplementary Data online at <http://care.diabetesjournals.org/lookup/suppl/doi:10.2337/dc16-1186/-/DC1>.

The opinions expressed are those of the authors and do not necessarily reflect the views of the Centers for Disease Control and Prevention.

This article is featured in a podcast available at <http://www.diabetesjournals.org/content/diabetes-core-update-podcasts>.

© 2017 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <http://www.diabetesjournals.org/content/license>.

Cardiovascular disease (CVD) imposes an enormous health and economic burden in the U.S. leading to ~800,000 deaths (nearly one in every three deaths) and \$320 billion in costs in 2011 (1,2). CVD is preventable through the successful management of risk factors, including preventing type 2 diabetes and lowering blood pressure and cholesterol levels (3,4).

In August 2014, the U.S. Preventive Services Task Force (USPSTF) recommended an intensive behavioral counseling intervention to reduce CVD risks in overweight or obese adults with one or more of the following risk factors: hypertension, dyslipidemia, impaired fasting glucose (IFG), or metabolic syndrome (5). The intervention promotes a healthy diet and physical activity through health education, individual feedback, problem-solving skills, and an individualized plan. Interventions are delivered by trained professionals, such as diabetes educators, dietitians, and behavioral interventionists; or by trained lay persons, such as community health workers. The intervention involves multiple contacts, either individually or in a group setting, over an extended period of time (e.g., 1 year). The USPSTF based its recommendation largely on the consistent evidence that behavioral counseling for lifestyle change improves CVD risk factors and reduces type 2 diabetes incidence (6).

The new recommendation is rated “grade B” and thus must be covered by private insurance under the Affordable Care Act without a copay. Because a significant proportion of U.S. adults will be eligible for the intervention (7), scaling the intervention nationwide will require substantial resources from public and private sectors. Similar interventions have been estimated to cost \$400 to \$1,000 per person per year (8). However, such interventions may be an efficient use of health care resources in the long run if substantial medical costs are saved by preventing or delaying CVD, type 2 diabetes, and diabetes-related complications. No previous study has estimated the long-term health and economic implications of implementing the USPSTF-recommended intensive behavioral counseling intervention in the U.S. population.

The primary objectives of our study are 1) to assess the long-term cost-effectiveness (CE) of USPSTF-recommended intensive behavioral counseling interventions in the overall target population and 2) to examine the variation in CE across

subgroups, as defined by combinations of risk factors specified in the USPSTF recommendation.

RESEARCH DESIGN AND METHODS

We measured the CE of the USPSTF-recommended intervention using incremental CE ratios (ICERs), expressed in cost per quality-adjusted life-year (QALY) gained by implementing the recommended intervention, compared with no intervention. First, we identified the population of U.S. adults who were eligible for the intervention based on the USPSTF-defined criteria. Second, we specified the intervention and its assumed effectiveness and costs based on previous studies. Third, we applied a discrete Markov disease progression model to simulate the long-term health and cost impacts with and without implementing the intervention. Finally, we calculated the ICER by dividing the difference in costs by the difference in QALYs

with and without the intervention. We evaluated the ICER over a time horizon of 25 years because health benefits may not be realized in a short period (9). The conventionally used \$50,000/QALY threshold for adopting a new intervention was applied to assess CE (10).

Study Population

We used the National Health and Nutrition Examination Survey (NHANES) 2005–2012 to identify demographic characteristics and health profiles of the intervention population in the U.S. (Table 1). Estimates were weighted according to the NHANES sampling design. We used a BMI of ≥ 25 and < 30 kg/m² to define overweight and a BMI ≥ 30 kg/m² to define obesity. People with hypertension included those with systolic blood pressure ≥ 140 mmHg or diastolic blood pressure ≥ 90 mmHg (11), and those who had received a diagnosis of hypertension by a physician

Table 1—Characteristics of the eligible population for USPSTF-recommended intensive behavioral counseling interventions

Risk factors	Criteria	Proportion of the eligible population
Total number of adults recommended for the intervention		98 million*
Overweight	BMI ≥ 25 and < 30 kg/m ²	50.8%
Obese	BMI ≥ 30 kg/m ²	49.2%
Hypertension and overweight/obese	$\geq 140/90$ mmHg	35.2%
Dyslipidemia and overweight/obese	ATP III borderline high definitions under the following conditions (12)	82.4%
High cholesterol	≥ 200 mg/dL	58.0%
High LDL	≥ 130 mg/dL	46.2%
High triglyceride	≥ 150 mg/dL	39.6%
HDL	< 40 mg/dL	23.7%
IFG and overweight/obese	FPG ≥ 100 and < 126 mg/dL	45.7%
Metabolic syndrome and overweight/obese	ATP III definition† (14)	41.8%
Risk factors	Average value among eligible population	
BMI	31.5	
Weight	90.9 kg	
A1C	5.5% (37 mmol/mol)	
Total cholesterol	209.3 mg/dL	
HDL	49.2 mg/dL	
LDL	125.7 mg/dL	
Systolic blood pressure	123.5 mmHg	
Diastolic blood pressure	72.9 mmHg	
Triglycerides	152.8 mg/dL	

†Metabolic syndrome was defined using ATP III definition by having three or more of the five following conditions: abdominal (central) obesity (> 102 cm [40 inches] or > 88 cm [35 inches] in women); elevated blood pressure ($\geq 130/85$ mmHg); IFG; high serum triglyceride levels (≥ 150 mg/dL); and low HDL levels (HDL < 40 or < 50 mg/dL for women). *Based on data from the 2005–2012 NHANES. The total adult population aged ≥ 18 years was estimated at 225 million, which is comparable with the U.S. Census estimate of 227 million for 2009 (36).




Figure 1—25-year ICERs (\$USD/QALY) of the intensive behavioral counseling intervention compare with no intervention by risk factor. Those with none of the three major risk factors (dyslipidemia, hypertension, and IFG) have only metabolic syndrome (women who have abdominal obesity and specific blood pressure level [SBP 130–140 mmHg or DBP 85–90 mmHg] with HDL from 40 to 50 mg/dL). Willingness-to-pay (WTP) = \$50,000/QALY is used as the CE threshold.

one of the other three risk factors and overweight adults with IFG (cost-saving and CE groups) would reduce the people eligible for the targeted program from 98 million to 66.7 million nationally and would lower the intervention cost by ~\$20 billion.

Sensitivity Analyses

Detailed results of the sensitivity analysis are described in Supplementary Table 2. For the one-way sensitivity analysis, shortening the analytic horizon would increase the ICER while increasing the analytic horizon to a lifetime would lower ICER. Assuming zero and 5% discount rates yielded \$5,200/QALY and \$20,200/QALY, respectively. Varying intervention effectiveness had a large effect on CE. Assuming the lower effectiveness bounds yielded a higher ICER, while assuming the upper bounds yielded a lower ICER. Adding screening costs increased the ICERs. Delivering the intervention in a group setting would lower the ICER.

For the two-way sensitivity analysis, under the most favorable scenario (i.e., achieving the upper bound effectiveness with a group-based intervention cost), the intervention led to a cost savings of \$64/person receiving the intervention. In contrast, with the worst scenario (i.e., achieving the lower bound effectiveness with additional screening costs), the intervention yielded an ICER of \$26,300/QALY.

For extended effectiveness and cost scenarios, scenario 1 yielded a cost

saving at \$90/person in 25 years, while under scenario 2 the intervention would have an ICER of \$11,600/QALY.

CONCLUSIONS

We estimate that under the new USPSTF recommendation on behavioral counseling for CVD prevention, ~98 million Americans are eligible for the intervention, which would cost \$64 billion if all were to participate. Applying the conventional “willingness-to-pay” cutoff of \$50,000/QALY (10), the intervention is cost effective for the overall targeted population as well as for each age-group. However, CE varies substantially depending on the risk factor profile of the participants; the intervention is cost effective for overweight adults with IFG and for obese adults with at least one of three risk factors (dyslipidemia, hypertension, or IFG); these two groups account for ~68% of the eligible population. The intervention is cost saving if it was implemented in persons who are obese with IFG and hypertension, dyslipidemia, or both, ~19.8 million or 20.2% of all eligible population. CE could be improved substantially by targeting these higher-risk subgroups and/or delivering the intervention in group settings.

Our results are consistent with those of previous studies that found intensive lifestyle interventions aimed at reducing the incidence of type 2 diabetes among people with prediabetes to be cost effective, with a median cost of approximately

\$14,000/QALY gained (15). Our risk group analysis was also consistent with a previous study (30) of diabetes prevention interventions, in which those with higher levels of fasting plasma glucose or A1C had more favorable CE ratios than those at the lower end of the prediabetes spectrum. Applying the USPSTF recommendation to those with a relatively low risk (overweight rather than obese, and one additional risk factor rather than multiple) diminishes CE because, while the costs for implementing the intervention are the same, the number of cases of diabetes and CVD averted is smaller.

Reductions in BMI and blood glucose levels have more impact on CE than reductions in blood pressure and lipid levels, because of their greater risk reduction of diabetes. Previous studies (3,4) have shown that behavioral counseling intervention can reduce the risk of type 2 diabetes by 38–60%. A recent study (30) suggested that delaying or preventing type 2 diabetes for 10 years for a person at age 40 years might save more than \$30,000 in lifetime medical spending. However, the effects of this intervention on other risk factors (i.e., lipid levels and blood pressure) are modest. The reductions in systolic/diastolic blood pressure reported by the USPSTF were 1–3 mmHg, and for LDL were 1.4–6 mg/dL (5,6). It is likely that the use of hypertension and dyslipidemia medications among those who already had these conditions may also have diluted the impact of the intervention.

As expected, we found the recommended intervention to be more cost effective in the longer simulation horizon. Chronic disease prevention typically provides more benefit over the long term than the short term. CVD events may not occur in the short term, and diabetes-related complications typically do not occur until years after diabetes onset. While policy makers and program planners are often interested in short-term results, it may be more appropriate to take a longer perspective when evaluating CVD and diabetes prevention.

There are four notable limitations of the study. First, in the model, we included only the health benefits reported in the USPSTF recommendation. Excluding other potential health benefits, such as reducing cancer incidence (31), would underestimate the CE of the recommended intervention. Second, while existing evidence has provided compelling evidence of the efficacy of the recommended interventions in clinical settings, its effectiveness in real-world settings and sustainable effectiveness in the long term remain to be demonstrated. Lack of long-term effectiveness data in real-world settings adds some uncertainty to our modeling results. Thus, we conducted extensive sensitivity analyses, and the results support the robustness of our conclusions. Third, similar to all disease-modeling studies, our results are subject to the limitations of the model structure, the risk equations used, and the model assumptions. The CVD risk equations that were developed for the original CDC-RTI model were based on data from the UKPDS (19,32), a study of people with type 2 diabetes in the U.K. Because our study population included both persons with and without diabetes in the U.S., we used the recently developed ASCVD equations (20). The ASCVD equations were developed using data from the CARDIA study, the ARIC study, the CHS, and the Framingham Study and have been validated using contemporary clinical data. However, the accuracy and validity of the equations in some populations remain subject to debate (33,34). The CDC-RTI simulation model uses the UKPDS microvascular complication risk equations (19,32) and their ability to predict the future risk of microvascular complications among contemporary patients may be subject to treatment changes.

However, there are no other risk equations that are superior to the UKPDS equations (35). Finally, our model uses a cohort-based approach, and thus the results represent average effects without considering individual-level stochasticity. Because our study is intended to assist policy decisions at a population level, variations in CE results at an individual level are less relevant.

In summary, we found that the USPSTF-recommended intervention is likely to be cost effective for overweight and obese adults with CVD risk factors. However, CE varies by risk subgroups, suggesting that those who are obese with at least one CVD risk factor (dyslipidemia, hypertension, or IFG), or who are overweight with IFG may be priority populations. Further refinement of recommendations for risk stratification, and focusing on a group-only delivery method with comparable effectiveness, may improve the CE of the intervention.

Acknowledgments. The authors thank Thomas J. Hoerger, RTI UNC, Center of Excellence in Health Promotion Economics, for suggestions on setting up the simulation; Rui Li, Centers for Disease Control and Prevention, for sharing the cost estimate of the intervention before its publication; and Elizabeth Luman and Clarice Conley, Centers for Disease Control and Prevention, for their editorial comments.

Duality of Interest. The study was conducted by the U.S. Government employees as part of their duties. No other potential conflicts of interest relevant to this article were reported.

Author Contributions. J.L. designed the study, reviewed the literature, analyzed the baseline data and the results, modified the simulation model, and wrote the first draft of the manuscript. X.Z. designed the study, reviewed the literature, and analyzed the baseline data. B.B. analyzed the baseline data. D.B.R., W.E.G., Y.H., G.W., and A.A. designed the study. P.Z. designed the study and reviewed the literature. All authors contributed to the discussion and reviewed and edited the manuscript. J.L., D.B.R., W.E.G., and P.Z. are the guarantors of this work and, as such, had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Prior Presentation. Parts of this study were presented in abstract form at the 75th Scientific Sessions of the American Diabetes Association, Boston, MA, 5–9 June 2015.

References

1. Mozaffarian D, Benjamin EJ, Go AS, et al. Heart disease and stroke statistics-2016 update: a report from the American Heart Association. *Circulation* 2016;133:e38–e360
2. Mozaffarian D, Benjamin EJ, Go AS, et al.; American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart

disease and stroke statistics-2015 update: a report from the American Heart Association. *Circulation* 2015;131:e29–e322

3. Li G, Zhang P, Wang J, et al. The long-term effect of lifestyle interventions to prevent diabetes in the China Da Qing Diabetes Prevention Study: a 20-year follow-up study. *Lancet* 2008;371:1783–1789
4. Knowler WC, Fowler SE, Hamman RF, et al.; Diabetes Prevention Program Research Group. 10-year follow-up of diabetes incidence and weight loss in the Diabetes Prevention Program Outcomes Study. *Lancet* 2009;374:1677–1686
5. LeFevre ML; U.S. Preventive Services Task Force. Behavioral counseling to promote a healthful diet and physical activity for cardiovascular disease prevention in adults with cardiovascular risk factors: U.S. Preventive Services Task Force Recommendation Statement. *Ann Intern Med* 2014;161:587–593
6. Lin JS, O'Connor E, Evans CV, Senger CA, Rowland MG, Groom HC. Behavioral counseling to promote a healthy lifestyle in persons with cardiovascular risk factors: a systematic review for the U.S. Preventive Services Task Force. *Ann Intern Med* 2014;161:568–578
7. Omura JD, Carlson SA, Paul P, et al. Adults eligible for cardiovascular disease prevention counseling and participation in aerobic physical activity—United States, 2013. *MMWR Morb Mortal Wkly Rep* 2015;64:1047–1051
8. Hernan WH, Brandle M, Zhang P, et al.; Diabetes Prevention Program Research Group. Costs associated with the primary prevention of type 2 diabetes mellitus in the diabetes prevention program. *Diabetes Care* 2003;26:36–47
9. Zhuo X, Zhang P, Gregg EW, et al. A nationwide community-based lifestyle program could delay or prevent type 2 diabetes cases and save \$5.7 billion in 25 years. *Health Aff (Millwood)* 2012;31:50–60
10. Grosse SD. Assessing cost-effectiveness in healthcare: history of the \$50,000 per QALY threshold. *Expert Rev Pharmacoecon Outcomes Res* 2008;8:165–178
11. Chobanian AV, Bakris GL, Black HR, et al.; Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. National Heart, Lung, and Blood Institute; National High Blood Pressure Education Program Coordinating Committee. Seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Hypertension* 2003;42:1206–1252
12. Antonopoulos S; National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). Third report of the National Cholesterol Education Program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adult Treatment Panel III) final report. *Circulation* 2002;106:3143–3421
13. American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2014;37(Suppl. 1):S81–S90
14. Grundy SM, Brewer HB Jr, Cleeman JI, Smith SC Jr, Lenfant C; American Heart Association; National Heart, Lung, and Blood Institute. Definition of metabolic syndrome: report of the National Heart, Lung, and Blood Institute/American Heart

Association conference on scientific issues related to definition. *Circulation* 2004;109:433–438

15. Li R, Qu S, Zhang P, et al. Economic evaluation of combined diet and physical activity promotion programs to prevent type 2 diabetes among persons at increased risk: a systematic review for the Community Preventive Services Task Force. *Ann Intern Med* 2015;163:452–460
16. CDC Diabetes Cost-effectiveness Group. Cost-effectiveness of intensive glycemic control, intensified hypertension control, and serum cholesterol level reduction for type 2 diabetes. *JAMA* 2002;287:2542–2551
17. Zhuo X, Zhang P, Selvin E, et al. Alternative HbA1c cutoffs to identify high-risk adults for diabetes prevention: a cost-effectiveness perspective. *Am J Prev Med* 2012;42:374–381
18. Hoerger TJ, Segel JE, Zhang P, Sorensen SW. *Validation of the CDC-RTI Diabetes Cost-Effectiveness Model*. Research Triangle Park, NC, RTI Press, 2009
19. Clarke PM, Gray AM, Briggs A, et al.; UK Prospective Diabetes Study (UKPDS) Group. A model to estimate the lifetime health outcomes of patients with type 2 diabetes: the United Kingdom Prospective Diabetes Study (UKPDS) Outcomes Model (UKPDS no. 68). *Diabetologia* 2004;47:1747–1759
20. Goff DC, Lloyd-Jones DM, Bennett G, O'Donnell C, Coady S, Robinson J. 2013 ACC/AHA guideline on the assessment of cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2014;63(Suppl. 25):2935–2959
21. Friedman GD, Cutter GR, Donahue RP, et al. CARDIA: study design, recruitment, and some

characteristics of the examined subjects. *J Clin Epidemiol* 1988;41:1105–1116

22. ARIC Investigators. The Atherosclerosis Risk in Communities (ARIC) Study: design and objectives. The ARIC investigators. *Am J Epidemiol* 1989;129:687–702
23. Fried LP, Borhani NO, Enright P, et al. The cardiovascular health study: design and rationale. *Ann Epidemiol* 1991;1:263–276
24. Kannel WB, Feinleib M, McNamara PM, Garrison RJ, Castelli WP. An investigation of coronary heart disease in families. The Framingham offspring study. *Am J Epidemiol* 1979;110:281–290
25. Muntner P, Colantonio LD, Cushman M, et al. Validation of the atherosclerotic cardiovascular disease Pooled Cohort risk equations. *JAMA* 2014;311:1406–1415
26. Henderson KH, Kaufman BG, Stearns S, et al. Validation of the atherosclerotic cardiovascular disease (ASCVD) pooled cohort risk equations by education level: the Atherosclerosis Risk in Communities (ARIC) Study. *J Am Coll Cardiol* 2016;67:1842
27. Herman WH, Hoerger TJ, Brandle M, et al.; Diabetes Prevention Program Research Group. The cost-effectiveness of lifestyle modification or metformin in preventing type 2 diabetes in adults with impaired glucose tolerance. *Ann Intern Med* 2005;142:323–332
28. U.S. Centers for Medicare & Medicaid Services. Physician Fee Schedule Look-Up Tool [Internet], 2016. Available from <https://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/PFSlookup/index.html>. Accessed 27 January 2016
29. Balk EM, Earley A, Raman G, Avendano EA, Pittas AG, Remington PL. Combined diet and

physical activity promotion programs to prevent type 2 diabetes among persons at increased risk: a systematic review for the Community Preventive Services Task Force. *Ann Intern Med* 2015;163:437–451

30. Zhuo X, Zhang P, Barker L, Albright A, Thompson TJ, Gregg E. The lifetime cost of diabetes and its implications for diabetes prevention. *Diabetes Care* 2014;37:2557–2564
31. Anand P, Kunnumakkara AB, Sundaram C, et al. Cancer is a preventable disease that requires major lifestyle changes. *Pharm Res* 2008;25:2097–2116
32. Hayes AJ, Leal J, Gray AM, Holman RR, Clarke PM. UKPDS outcomes model 2: a new version of a model to simulate lifetime health outcomes of patients with type 2 diabetes mellitus using data from the 30 year United Kingdom Prospective Diabetes Study: UKPDS 82. *Diabetologia* 2013;56:1925–1933
33. Cook NR, Ridker PM. Response to Comment on the reports of over-estimation of ASCVD risk using the 2013 AHA/ACC risk equation. *Circulation* 2014;129:268–269
34. Ridker PM, Cook NR. Statins: new American guidelines for prevention of cardiovascular disease. *Lancet* 2013;382:1762–1765
35. Palmer AJ, Clarke P, Gray A, et al.; Mount Hood 5 Modeling Group. Computer modeling of diabetes and its complications: a report on the Fifth Mount Hood challenge meeting. *Value Health* 2013;16:670–685
36. *Age and Sex Composition in the United States: 2009 [Internet], 2009*. Washington, DC, United States Census Bureau. Available from <https://www.census.gov/population/age/data/2009comp.html>. Accessed 26 April 2016