



Prevalence of and Risk Factors for Diabetic Peripheral Neuropathy in Youth With Type 1 and Type 2 Diabetes: SEARCH for Diabetes in Youth Study

Diabetes Care 2017;40:1226–1232 | <https://doi.org/10.2337/dc17-0179>

Mamta Jaiswal,¹ Jasmin Divers,²
Dana Dabelea,³ Scott Isom,²
Ronny A. Bell,⁴ Catherine L. Martin,⁵
David J. Pettitt,⁶ Sharon Saydah,⁷
Catherine Pihoker,⁸ Debra A. Standiford,⁹
Lawrence M. Dolan,⁹ Santica Marcovina,¹⁰
Barbara Linder,¹¹ Angela D. Liese,¹²
Rodica Pop-Busui,⁵ and Eva L. Feldman¹

OBJECTIVE

We assessed the prevalence of and risk factors for diabetic peripheral neuropathy (DPN) in youth with type 1 diabetes (T1D) and type 2 diabetes (T2D) enrolled in the SEARCH for Diabetes in Youth (SEARCH) study.

RESEARCH DESIGN AND METHODS

The Michigan Neuropathy Screening Instrument (MNSI) was used to assess DPN in 1,734 youth with T1D (mean \pm SD age 18 ± 4 years, T1D duration 7.2 ± 1.2 years, and HbA_{1c} $9.1 \pm 1.9\%$) and 258 youth with T2D (age 22 ± 3.5 years, T2D duration 7.9 ± 2 years, and HbA_{1c} $9.4 \pm 2.3\%$) who were enrolled in the SEARCH study and had ≥ 5 years of diabetes duration. DPN was defined as an MNSI exam score of >2 . Glycemic control over time was estimated as area under the curve for HbA_{1c}.

RESULTS

The prevalence of DPN was 7% in youth with T1D and 22% in youth with T2D. Risk factors for DPN in youth with T1D were older age, longer diabetes duration, smoking, increased diastolic blood pressure, obesity, increased LDL cholesterol and triglycerides, and lower HDL cholesterol (HDL-c). In youth with T2D, risk factors were older age, male sex, longer diabetes duration, smoking, and lower HDL-c. Glycemic control over time was worse among those with DPN compared with those without for youth with T1D (odds ratio 1.53 [95% CI 1.24; 1.88]) but not for youth with T2D (1.05 [0.7; 1.56]).

CONCLUSIONS

The high rates of DPN among youth with diabetes are a cause of concern and suggest a need for early screening and better risk factor management. Interventions in youth that address poor glycemic control and dyslipidemia may prevent or delay debilitating neuropathic complications.

The increasing prevalence of type 1 (T1D) and type 2 diabetes (T2D) in children and adolescents in the U.S. (1) is likely to result in increased numbers of individuals with diabetes-related complications in their early adulthood. Diabetic peripheral neuropathy (DPN) is a debilitating complication that has been well characterized in adults, with prevalence rates ranging from 10–26% in newly diagnosed adults with diabetes (2). The prevalence and predictors of DPN in youth with diabetes in the U.S., however, have not

¹Department of Neurology, University of Michigan, Ann Arbor, MI

²Department of Biostatistical Sciences, Wake Forest School of Medicine, Winston-Salem, NC

³Department of Epidemiology, Colorado School of Public Health, Aurora, CO

⁴Department of Epidemiology and Prevention, Wake Forest School of Medicine, Winston-Salem, NC

⁵Division of Metabolism, Endocrinology and Diabetes, University of Michigan, Ann Arbor, MI

⁶Santa Barbara, CA

⁷Division of Diabetes Translation, Centers for Disease Control and Prevention, Atlanta, GA

⁸Department of Pediatrics, University of Washington, Seattle, WA

⁹Division of Endocrinology, Cincinnati Children's Hospital Medical Center, Cincinnati, OH

¹⁰Northwest Lipid Metabolism and Diabetes Research Laboratories, Department of Medicine, University of Washington, Seattle, WA

¹¹National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD

¹²Department of Epidemiology and Biostatistics and Center for Research in Nutrition and Health Disparities, Arnold School of Public Health, University of South Carolina, Columbia, SC

Corresponding author: Eva L. Feldman, efeldman@umich.edu.

Received 30 January 2017 and accepted 2 June 2017.

This article contains Supplementary Data online at <http://care.diabetesjournals.org/lookup/suppl/doi:10.2337/dc17-0179/-/DC1>.

© 2017 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <http://www.diabetesjournals.org/content/license>.

been systematically examined. Several small cross-sectional clinic-based studies using various definitions of DPN have reported a wide range of prevalence rates from 5 to 62% (3–6). Population-based longitudinal studies in Australia and Denmark have extensively analyzed the prevalence and predictors of microvascular complications, including DPN, in children and adolescents with diabetes (7,8). In the Australian cohort of 1,433 adolescents with T1D and 68 with T2D aged <18 years, the prevalence of DPN was 21% and 27%, respectively (8), while in the Danish cohort of 339 adolescents with T1D, the prevalence of DPN was 62% (7).

We previously reported results from a small pilot study comparing the prevalence of DPN in a subset of youth enrolled in the SEARCH for Diabetes in Youth (SEARCH) study and found that 8.5% of 329 youth with T1D (mean \pm SD age 15.7 \pm 4.3 years and diabetes duration 6.2 \pm 0.9 years) and 25.7% of 70 youth with T2D (age 21.6 \pm 4.1 years and diabetes duration 7.6 \pm 1.8 years) had evidence of DPN (9). Recently, we also reported the prevalence of microvascular and macrovascular complications in youth with T1D and T2D in the entire SEARCH cohort (10).

In the current study, we examined the cross-sectional and longitudinal risk factors for DPN. The aims were 1) to estimate prevalence of DPN in youth with T1D and T2D, overall and by age and diabetes duration, and 2) to identify risk factors (cross-sectional and longitudinal) associated with the presence of DPN in a multiethnic cohort of youth with diabetes enrolled in the SEARCH study.

RESEARCH DESIGN AND METHODS

SEARCH Study

SEARCH is a population-derived prospective cohort study following children and adolescents of diverse racial and ethnic backgrounds diagnosed with diabetes at <20 years of age in the U.S. The SEARCH participants are subjects with incident diabetes diagnoses identified at four geographically defined populations in Ohio, Washington, South Carolina, and Colorado; from health plan enrollees in California; and from Indian Health Service beneficiaries from American Indian populations in Arizona and New Mexico.

Study Population

Youth with diabetes diagnosed at <20 years of age were identified from a

population-derived incident registry network at five U.S. sites by the SEARCH for Diabetes in Youth Registry Study (11). Case subjects with newly diagnosed T1D or T2D in 2002–2006 or 2008, who completed a SEARCH baseline examination for risk factors (on average 9.3 \pm 6.4 months from diagnosis) and had at least 5 years of diabetes duration between 2011 and 2015, were recruited into the SEARCH Cohort Study to measure standardized outcomes and complete repeat risk factor assessments at the cohort visit (on average at 7.9 \pm 1.9 years from diagnosis). The SEARCH Cohort Study enrolled 2,777 individuals. For this analysis, we excluded participants aged <10 years ($n = 134$), those with no antibody measures for etiological definition of diabetes ($n = 440$), and those with incomplete neuropathy assessment (Michigan Neuropathy Screening Instrument [MNSI] and cardiac autonomic neuropathy examination) ($n = 213$), which reduced the analysis sample size to 1,992 (Supplementary Fig. 1).

Prior to protocol implementation, local institutional review board approval was obtained for each center. Written informed consent was obtained from participants age ≥ 18 years, while assent with parental written informed consent was obtained for participants <18 years old.

Baseline and Cohort Visits

The SEARCH baseline and cohort visits included a participant survey; measurement of height, weight, waist circumference, blood pressure, and DPN (cohort visit only); and blood and urine collection. Race and ethnicity were self-reported and categorized as non-Hispanic white, non-Hispanic black, Hispanic, Asian or Pacific Islander, and other. Current cigarette smoking was defined as having smoked cigarettes on one or more of the 30 days preceding the study visit. Individuals who had tried smoking or smoked regularly (at least one cigarette every day for 30 days) but were not current smokers were considered former smokers. Youth who had never smoked a whole cigarette were considered nonsmokers.

BMI was defined as weight in kilograms divided by the square of height in meters. Waist circumference was measured using the natural waist location (11). For participants <20 years of age, the Centers for Disease Control and Prevention (CDC)-derived BMI z20 scores were used; for those ≥ 20 years of age the observed

mean and SD were used to standardize their BMI z20 values.

Resting systolic blood pressure (SBP) and diastolic blood pressure (DBP) were measured three times using an aneroid sphygmomanometer while the participants were seated for at least 5 min, and the average of the three measurements was taken. Hypertension was defined as an SBP z score or DBP z score >95th percentile after accounting for age, sex, and height for youth ≤ 18 years of age; an SBP ≥ 140 mmHg or a DBP ≥ 90 mmHg for youth age >18 years; or use of blood pressure-lowering medication, regardless of age.

A blood draw occurred after an 8-h overnight fast, and medications, including short-acting insulin, were withheld the morning of the visit. Blood samples were obtained under conditions of metabolic stability, as defined by no episodes of diabetic ketoacidosis in the prior month. Specimens were processed locally at the sites and shipped within 24 h to the central laboratory (Northwest Lipid Metabolism and Diabetes Research Laboratories, University of Washington, Seattle, Washington), where they were analyzed for measurement of HDL cholesterol (HDL-c), LDL cholesterol (LDL-c), triglycerides, and HbA_{1c} as previously described (11). Urinary albumin and creatinine levels were assessed on a random spot urine sample to evaluate renal function using the albumin-to-creatinine ratio (ACR).

In addition to SEARCH baseline and cohort visits, 57% ($n = 1,113$) of participants had one or more intermediate visits (1-, 2-, and 5-year follow-up visits after baseline visit) at which risk factors were measured. These measurements included HbA_{1c}, lipids, and BMI. The assay of the biological samples has remained consistent over time. The accuracy of HbA_{1c} data was monitored by participation in the National Glycohemoglobin Standardization Program (NGSP), and accuracy and the consistency of the laboratory assay for lipids were monitored regularly by comparing results obtained by the enzymatic methods with those obtained by the CDC reference methods (CDC Reference Laboratory) (12).

Diabetes Type

Diabetes type was defined using an etiological classification developed by SEARCH (10) based on one or more positive diabetes autoantibodies (GAD-65, insulinoma-associated-2 antibodies, and Zinc-T8

autoantibody) and estimated insulin sensitivity score (validated equation including waist circumference, HbA_{1c}, and triglyceride levels) at the baseline visit (11). T1D was defined as at least one positive antibody, regardless of insulin sensitivity, or no positive antibodies and insulin sensitivity (score ≥ 8.15). T2D was defined as negative antibodies and insulin resistance (score < 8.15) (13).

Assessment of DPN

DPN was assessed during the cohort visit using the MNSI, a validated screening tool for DPN (14,15). SEARCH staff from each center were centrally trained and certified to perform the MNSI. The MNSI is a 15-item self-administered questionnaire and structured examination of the feet (MNSIE) which is scored for abnormalities of appearance (deformities, infection, and dry skin/callus), presence of ulcers, vibration perception at the distal great toe, and ankle reflexes. Scores assigned during the MNSIE procedure include appearance of feet (normal = 0, abnormal = 1), ulceration (absent = 0, present = 1), ankle reflexes (absent = 1, present with reinforcement = 0.5, present = 0), and vibration perception (absent = 1, reduced = 0.5, present = 0) and are graded separately for each foot, for a range of score from 0 (normal examination) to 8. The threshold for DPN, established by prior validation studies performed among adults, is a score of > 2 on the MNSIE out of a total score of 8 (15).

Statistical Analyses

Cross-sectional Data

Anthropometric, demographic, and metabolic data collected at the cohort visit as described above were used to compare the characteristics distinguishing youth with and without DPN stratified by diabetes type. Wilcoxon two-sample tests were used to compare the distribution of continuous variables, and the χ^2 test was used for categorical variables separately for T1D and T2D participants. Fisher exact test was used whenever a cell count for a particular test was < 5 . The prevalence of DPN was estimated overall and based on the age at diagnosis (≥ 10 years and < 10 years) and duration of diabetes (5–10 years and > 10 years) separately for subjects with T1D and T2D.

Longitudinal Data

In addition to the data collected at baseline and at the cohort visit, the area under

the curve (AUC) was computed to summarize the longitudinal trajectory of HbA_{1c} and other continuous variables, such as lipids, height, blood pressure, and BMI, collected over time (at the baseline and 1-, 2-, and 5-year follow-up and cohort visits), with adjustment for the interval between the first and last measurement. For assessment of the effect of long-term glycemic control on DPN, logistic regression models treating presence of DPN as the outcome were fitted separately for participants with T1D and T2D. These models were adjusted for age and sex (model 2), height z score (model 2a), BMI (model 3), blood pressure (model 4), triglycerides (model 5), and ACR (model 6). A fully adjusted model that included all of these variables as covariates was also fitted (model 7). Models were stratified by diabetes type to limit confounding effects of age and adiposity. Diagnostic tests were performed to ensure that modeling assumptions were satisfied. The data were analyzed using SAS 9.4 (SAS Institute, Cary, NC).

RESULTS

There were 1,734 youth with T1D and 258 youth with T2D who participated in the SEARCH study and had complete data for the variables of interest. The clinical and metabolic characteristics of the participants with T1D and T2D at the cohort visit (2011–2015), stratified by DPN status, are presented in Table 1. Seven percent of the participants with T1D and 22% of those with T2D had evidence of DPN.

Among youth with T1D, those with DPN were older (21 vs. 18 years, $P < 0.0001$), had a longer duration of diabetes (8.7 vs. 7.8 years, $P < 0.0001$), and had higher DBP (71 vs. 69 mmHg, $P = 0.02$), BMI (26 vs. 24 kg/m², $P < 0.001$), and LDL-c levels (101 vs. 96 mg/dL, $P = 0.01$); higher triglycerides (85 vs. 74 mg/dL, $P = 0.005$); and lower HDL-c levels (51 vs. 55 mg/dL, $P = 0.01$) compared to those without DPN. The prevalence of DPN was 5% among nonsmokers vs. 10% among the current and former smokers ($P = 0.001$).

Among youth with T2D, those with DPN were older (23 vs. 22 years, $P = 0.01$), had longer duration of diabetes (8.6 vs. 7.6 years; $P = 0.002$), and had lower HDL-c (40 vs. 43 mg/dL, $P = 0.04$) compared with those without DPN. The prevalence of DPN was higher among males than among females: 30% of males had DPN compared with 18% of females

($P = 0.02$). The prevalence of DPN was twofold higher in current smokers (33%) compared with nonsmokers (15%) and former smokers (17%) ($P = 0.01$).

The AUC for various metabolic and clinical parameters collected over time is shown in Table 2. The AUCs for HbA_{1c}, LDL-c, triglycerides, SBP, and DBP were higher and the AUC for HDL-c was lower among youth with T1D with DPN compared with those without DPN (all $P < 0.002$). These variables, however, were not significantly different in youth with T2D with or without DPN, which could be due to the comparatively small sample size.

Logistic regression models for association between DPN and long-term glycemic control (HbA_{1c} AUC) adjusted for age and sex, BMI, blood pressure, triglycerides, ACR, and all variables combined were also examined (Table 3). HbA_{1c} AUC was significantly associated with risk of DPN in youth with T1D (independent of cardiovascular risk factors). These associations were not statistically significant in the T2D subgroup; however, association parameters were generally in the same direction as those observed in T1D. The association between DPN and HbA_{1c} AUC for individuals with only two measures (at baseline and cohort visit; $n = 879$) versus those with three to five measures remained the same for both youth with T1D and youth with T2D ($n = 1,113$) (data not shown).

Finally, the prevalence of DPN was further assessed by 5-year increment of diabetes duration in individuals with T1D or T2D (Fig. 1). There was an approximately twofold increase in the prevalence of DPN with an increase in duration of diabetes from 5–10 years to > 10 years for both the T1D group (5–13%) ($P < 0.0001$) and the T2D group (19–36%) ($P = 0.02$).

Individuals age ≥ 10 years who were excluded from the study owing to missing data ($n = 653$) were more likely to belong to a minority racial/ethnic group (non-Hispanic Black 20 vs. 14%, Hispanic 15 vs. 13%, and Asian Pacific Islander 2 vs. 1.5%, $P < 0.001$), had a longer duration of diabetes (8.3 vs. 7.8 years, $P < 0.001$), were younger at the time of diabetes diagnosis (9.2 vs. 10.5 years, $P < 0.001$), and were more likely to have hypertension (5.5 vs. 2.8%, $P = 0.003$) compared with those included in the analysis cohort ($n = 1,992$) (Supplementary Table 1).

Table 1—Characteristics of the youth with T1D and T2D from the SEARCH Cohort Study stratified by DPN status (cross-sectional data collected at cohort visit between 2011 and 2015)

	T1D (N = 1,734)			T2D (N = 258)		
	No DPN	DPN	P	No DPN	DPN	P
N (%)	1,620 (93)	114 (7)	N/A	202 (78)	56 (22)	N/A
Age, years	18 ± 4	21 ± 4	<0.0001	22 ± 3	23 ± 4	0.01
Age at dx ≥10 years	806 (90)	88 (10)	<0.0001	191 (78)	55 (22)	0.25
Age at dx <10 years	814 (97)	26 (3)		11 (92)	1 (8)	
Female sex	807 (94)	55 (6)	0.746	144 (82)	31 (18)	0.02
Male sex	813 (93)	59 (7)		58 (70)	25 (30)	
Diabetes duration, years	7.8 ± 1.8	8.7 ± 2	<0.0001	7.6 ± 1.9	8.6 ± 2	0.002
5–10	1,395 (95)	81 (5)	<0.0001	175 (81)	41 (19)	0.02
>10	225 (87)	33 (13)		27 (64)	15 (36)	
Race/ethnicity						
Non-Hispanic white	1,237 (94)	85 (6)	0.65	55 (80)	14 (20)	0.77
Non-Hispanic black	150 (92)	14 (8)		94 (80)	23 (20)	
Hispanic	197 (95)	11 (5)		37 (71)	15 (29)	
Asian or Pacific Islander	24 (89)	3 (11)		2 (67)	1 (33)	
Other	5 (100)	0 (0)		1 (100)	0 (0)	
Native American	7 (88)	1 (12)		13 (81)	3 (19)	
Smoking						
Nonsmoker	1,099 (95)	56 (5)	0.001	82 (85)	15 (15)	0.01
Former	281 (90)	33 (10)		62 (83)	13 (17)	
Current	207 (90)	23 (10)		53 (67)	26 (33)	
Hypertension, yes	26 (1.6)	5 (1.7)	0.25	23 (11.4)	2 (8.9)	0.25
SBP, mmHg	106.3 ± 10.9	107 ± 11	0.68	117.6 ± 13.3	119.8 ± 14.1	0.80
SBP z score	−0.3 ± 1.6	0.1 ± 1.9	0.08	1.9 ± 2.5	2.9 ± 4	0.27
DBP, mmHg	68.6 ± 8.7	70.8 ± 9.1	0.02	75.8 ± 10.3	76.3 ± 10.3	0.71
DBP z score	−0.1 ± 2.02	−0.2 ± 3.3	0.74	1.4 ± 3.8	1.1 ± 5.6	0.54
Waist circumference, cm	77.6 ± 12.1	83.4 ± 13.2	<0.001	104.9 ± 20.4	106.1 ± 17.3	0.36
Waist-to-height ratio	0.47 ± 0.06	0.49 ± 0.07	0.0022	0.62 ± 0.12	0.62 ± 0.1	0.63
BMI, kg/m ²	24 ± 5	26 ± 5.8	<0.001	35.6 ± 9.1	35.1 ± 8.8	0.81
BMI z20 score	0.6 ± 0.7	0.7 ± 0.9	0.16	1.8 ± 2	1.7 ± 1.9	0.73
HbA _{1c} , %	9.1 ± 1.8	9.5 ± 2.3	0.07	9.2 ± 3	9.1 ± 3	0.71
HbA _{1c} , mmol/mol	75 ± 7	80 ± 8	0.07	77 ± 9	76 ± 9	0.71
LDL-c, mg/dL	96 ± 28	101 ± 25	0.01	106 ± 37	100 ± 43	0.27
Triglycerides, mg/dL	74 (55, 105)	85 (61, 126)	0.0052	115 (79, 188)	121 (85, 195)	0.61
HDL-c, mg/dL	55.3 ± 13.8	51.2 ± 11.8	0.01	42.7 ± 12.4	39.6 ± 11.2	0.04
ACR, μg/mg	6.1 (4.2, 10.8)	6.1 (3.8, 12.4)	0.93	8.5 (4.3, 30.5)	12.4 (4.9, 69.2)	0.46

Data are mean ± SD, median (interquartile range), or n (%). dx, diagnosis; N/A, not applicable.

Finally, in an unadjusted logistic regression model, youth with T2D were four times more likely to develop DPN compared with those with T1D, and though this association was attenuated, it remained significant independent of age, sex, height, and glycemic control (OR 2.99 [1.91; 4.67], $P < 0.001$) (Supplementary Table 2).

CONCLUSIONS

The prevalence of DPN in this multiethnic cohort of youth with T1D and T2D was 7 and 22%, respectively. The risk factors for DPN among youth with T1D included older age, smoking, longer duration of diabetes, long-term poor glycemic control, and cardiovascular risk factors, while

those among youth with T2D included older age, smoking, and longer diabetes duration. Long-term poor glycemic control and dyslipidemia emerged as the modifiable risk factors for both groups.

The prevalence estimates for DPN found in our study for youth with T2D are similar to those in the Australian cohort (8) but lower for youth with T1D than those reported in the Danish (7) and Australian (8) cohorts. The nationwide Danish Study Group for Diabetes in Childhood reported a prevalence of 62% among 339 adolescents and youth with T1D (age 12–27 years, duration 9–25 years, and HbA_{1c} 9.7 ± 1.7%) using the vibration perception threshold to assess DPN (7).

The higher prevalence in this cohort compared with ours (62 vs. 7%) could be due to the longer duration of diabetes (9–25 vs. 5–13 years) and reliance on a single measure of neuropathy (vibration perception threshold) as opposed to our use of the MNSI, which includes vibration as well as other indicators of neuropathy. In the Australian study, Eppens et al. (8) reported abnormalities in peripheral nerve function in 27% of the 1,433 adolescents with T1D (median age 15.7 years, median diabetes duration 6.8 years, and mean HbA_{1c} 8.5%) and 21% of the 68 adolescents with T2D (median age 15.3 years, median diabetes duration 1.3 years, and mean HbA_{1c} 7.3%) based on thermal and

Table 2—AUC for cumulative risk for various continuous variables in youth with T1D and T2D enrolled in the SEARCH study, stratified by DPN status (longitudinal data)

AUC	T1D			T2D		
	No DPN	DPN	<i>P</i>	No DPN	DPN	<i>P</i>
BMI z score	20.6 ± 0.9	20.6 ± 1	0.37	21.9 ± 0.7	21.9 ± 0.7	0.93
Height, m	1.6 ± 0.2	1.7 ± 0.1	<0.001	1.7 ± 0.1	1.7 ± 0.1	0.02
Height z score	0.3 ± 1	0.5 ± 1.1	<0.001	0.4 ± 1	0.5 ± 1	0.32
SBP	103.5 ± 9.2	106.6 ± 8.4	<0.001	116.5 ± 9.7	117.8 ± 10.2	0.53
DBP	65.8 ± 7.3	68.2 ± 6.8	<0.001	72.9 ± 7.2	74.3 ± 8.3	0.25
HbA _{1c} %	8.5 ± 1.3	9.1 ± 1.8	<0.001	8.4 ± 2.3	8.7 ± 2.3	0.48
HbA _{1c} mmol/mol	69.1 ± 14.3	76 ± 19.8	<0.001	68 ± 25.5	71.1 ± 27.6	0.48
HDL-c	56.2 ± 11.8	51.9 ± 10.4	<0.001	42.8 ± 11.4	39.8 ± 7.8	0.09
LDL-c	93.1 ± 22.4	100.1 ± 22.8	0.002	102.8 ± 30.3	101.1 ± 30.7	0.67
Triglycerides	68.5 (53.5, 90.2)	78 (62.9, 106.2)	<0.001	117.7 (85.4, 181.6)	128.5 (90.5, 209)	0.45

Data are mean ± SD or median (interquartile range).

vibration perception threshold. These data are thus reminiscent of the persistent inconsistencies in the definition of DPN, which are reflected in the wide range of prevalence estimates being reported.

Our findings indicate that glycemic control over time (summarized as the AUC for HbA_{1c} adjusted for the time interval between the research visits) was significantly worse in those with DPN compared with those without DPN. The Diabetes Control and Complications Trial (DCCT) and UK Prospective Diabetes Study (UKPDS) have both reported that good glycemic control could potentially delay the development and progression of DPN and other microvascular complications in individuals with T1D and T2D (16,17). There are a number of pathological events that lead to functional and structural abnormalities of the peripheral nerves seen in DPN. These include the nonenzymatic glycosylation of the cytoskeletal proteins (tubulin, neurofilament, and actin), formation of advanced glycation end products, ischemic injury due to

inflammation, and dysfunction of endoneurial, perineurial, and epineurial blood vessels. These pathological events lead to axonal atrophy, degeneration, and impaired axonal transport and contribute to the functional and structural abnormalities (18). Thus, achieving and maintaining good glycemic control (HbA_{1c} ≤7.5%) could go a long way in preventing or delaying the development of DPN and other microvascular complication of diabetes in this young population.

The more than three times higher prevalence of DPN among youth with T2D (22%) compared with those with T1D (7%), despite the two groups having similar poor glycemic control and diabetes duration, is intriguing and could be attributed to their older age, although metabolic syndrome and a longer prediabetes phase among youth with T2D could also have played a significant role in the development of DPN (18). In recently published data from the SEARCH cohort, the rates of complications, including DPN, are nearly two to three times higher in youth

with T2D compared with T1D (10). The difference in the rate of DPN in youth with T1D and T2D suggests that some of the pathophysiological pathways driving the neural damage in these two groups could be divergent (19,20). Glycotoxicity and its downstream pathways could be a key player in T1D, while lipotoxicity and insulin resistance could be the major drivers among those with T2D (20). Moreover, youth with T1D and T2D are known to have different cardiovascular risk profiles, which could in part explain the difference in rates of DPN in this population at such a young age (21–23). Associations between diabetic dyslipidemia (higher triglycerides and lower HDL-c) and neuropathy have been reported in several landmark studies, such as the European Diabetes (EURODIAB) Prospective Complications Study (24) and the Pittsburgh Epidemiology of Diabetes Complications study (25). Interestingly, the lower HDL-c found in our cohort could be one of the key players in the pathogenesis of DPN. HDL-c has protective

Table 3—Logistic regression models for association between DPN and long-term glycemic control

Logistic regression model	T1D		T2D	
	OR (95% CI)	<i>P</i>	OR (95% CI)	<i>P</i>
Model 1: HbA _{1c} AUC	1.48 (1.24; 1.77)	<0.0001	1.13 (0.83; 1.52)	0.44
Model 2: model 1 + age and sex	1.44 (1.21; 1.71)	<0.0001	1.12 (0.83; 1.53)	0.45
Model 2a: model 1 + age, sex, and height z scores	1.50 (1.25; 1.79)	<0.0001	1.13 (0.83; 1.54)	0.44
Model 3: model 1 + BMI	1.49 (1.25; 1.78)	<0.0001	1.12 (0.82; 1.53)	0.48
Model 4: model 1 + SBP and DBP	1.45 (1.21; 1.73)	<0.0001	1.15 (0.84; 1.57)	0.37
Model 5: model 1 + TG	1.5 (1.24; 1.81)	<0.0001	1.1 (0.8; 1.5)	0.56
Model 6: model 1 + ACR	1.55 (1.28; 1.89)	<0.0001	1.07 (0.74; 1.54)	0.73
Model 7: model 1 + age, sex, BMI, SBP, DBP, TG, and ACR	1.53 (1.24; 1.88)	<0.0001	1.05 (0.7; 1.56)	0.82

Outcome variable: DPN. Independent variable: HbA_{1c} AUC. TG, triglycerides.




Figure 1—Prevalence of DPN among youth with T1D and T2D stratified by the duration of diabetes in years.

effects, including inhibition of inflammation, oxidation, and thrombosis, as well as vasodilatation via endothelial release of nitric oxide (26,27). HDL-c also removes lipids from peripheral cells via its effects on reverse cholesterol transport, which potentially ameliorates inflammation. Thus, targeting the lower HDL-c levels with interventions such as aerobic exercise, smoking cessation, weight loss, and dietary manipulation, which have been known to increase HDL-c levels by up to 20% and are generally associated with improved glycemic control (28), could be an additional therapeutic approach beyond optimal glycemic control in this young population.

Ours is the first population-derived study to assess the burden of DPN in an ethnically diverse cohort of youth with diabetes in the U.S. The alarming rise in rates of DPN for every 5-year increase in duration, coupled with poor glycemic control and dyslipidemia, in this cohort reinforces the need for clinicians rendering care to youth with diabetes to be vigilant in screening for DPN and identifying any risk factors that could potentially be modified to alter the course of the disease (28–30). The modifiable risk factors that could be targeted in this young population include better glycemic control, treatment of dyslipidemia, and smoking cessation (29,30)—approaches that are

also part of the Standards of Care recommendation by the American Diabetes Association (31). The sharp increase in rates of DPN over time is a reminder that DPN is one of the complications of diabetes that must be a part of the routine annual screening for youth with diabetes.

The population-based study design; large sample size; multiethnic composition of the cohort; use of a noninvasive, simple, and validated instrument to assess DPN; and evaluation of the longitudinal and cross-sectional risk factors are among the strengths of our study. The limited power to examine the association between long-term glycemic control and DPN among persons with T2D (despite similar levels of HbA_{1c}) was the result of a comparatively small sample size (although the association was in the same direction as that of the T1D group) and is one of the limitations of our study. The lack of longitudinal measures of DPN is also a limitation of our study, although a subset of this cohort (2002–2012 incidence case subjects ≥ 10 years of age with least 5 years of duration of diabetes) will be re-evaluated for DPN as a part of the next phase of SEARCH (2016–2020). Although the SEARCH Cohort Study is drawn from population-based registries of youth with diabetes, those excluded from the analytic sample were more likely to belong to a racial/ethnic minority, have

a longer duration of diabetes, and have hypertension. Each of these variables is associated with increased prevalence of DPN and may influence our estimates of prevalence of DPN among youth with diabetes.

Overall, the results of our study suggest that poor glycemic control over time and traditional cardiovascular risk factors are important risk factors associated with DPN and need to be targeted for the prevention of debilitating consequences of DPN in this young cohort.

Acknowledgments. The SEARCH study is indebted to the many youth and their families, as well as their health care providers, whose participation made this study possible.

Funding. The SEARCH study is funded by the National Institute for Health Research (1UC4DK108173-01) and National Institutes of Health (NIH) National Institute of Diabetes and Digestive and Kidney Diseases and supported by the CDC. The Population Based Registry of Diabetes in Youth Study (RFP DP15-002) is funded by the CDC and supported by the NIH National Institute of Diabetes and Digestive and Kidney Diseases. Specific funding for study sites is as follows: Kaiser Permanente Southern California (U18DP006133, U48/CCU919219, U01 DP000246, and U18DP002714), University of Colorado Denver (U18DP006139, U48/CCU819241-3, U01 DP000247, and U18DP000247-06A1), Cincinnati Children's Hospital Medical Center (U18DP006134, U48/CCU519239, U01 DP000248, and U18DP002709), University of North Carolina at Chapel Hill (U18DP006138, U48/CCU419249, U01 DP000254, and U18DP002708), Seattle Children's Hospital (U18DP006136, U58/CCU019235-4, U01 DP000244, and U18DP002710-01), and Wake Forest University School of Medicine (U18DP006131, U48/CCU919219, U01 DP000250, and 200-2010-35171). The authors acknowledge the involvement of the South Carolina Clinical and Translational Research Institute at the Medical University of South Carolina (NIH/National Center for Advancing Translational Sciences [NCATS] grant UL1 TR001450), Seattle Children's Hospital and the University of Washington (NIH/NCATS grant UL1 TR00423), University of Colorado Pediatric Clinical and Translational Research Center (NIH/NCATS grant UL1 TR000154), the Barbara Davis Center for Diabetes at the University of Colorado Denver (Diabetes and Endocrinology Research Center NIH grant P30-DK-57516), the University of Cincinnati (NIH/NCATS grant UL1 TR001425), and the Children with Medical Handicaps program managed by the Ohio Department of Health. This study includes data provided by the Ohio Department of Health.

The provision of data by the Ohio Department of Health should not be considered an endorsement of this study or its conclusions. The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the CDC or the National Institute of Diabetes and Digestive and Kidney Diseases.

Duality of Interest. No potential conflicts of interest relevant to this article were reported.

Author Contributions. M.J., J.D., and D.D. designed the study and analysis plan. M.J. wrote the manuscript. J.D. performed the analysis and provided critical input to the manuscript. D.D., S.I., R.A.B., C.L.M., D.J.P., S.S., C.P., D.A.S., L.M.D., S.M., B.L., A.D.L., R.P.-B., and E.L.F. critically reviewed the manuscript and provided input. J.D. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

References

- Dabelea D, Mayer-Davis EJ, Saydah S, et al.; SEARCH for Diabetes in Youth Study. Prevalence of type 1 and type 2 diabetes among children and adolescents from 2001 to 2009. *JAMA* 2014;311:1778–1786
- Pop-Busui R, Boulton AJ, Feldman EL, et al. Diabetic neuropathy: a position statement by the American Diabetes Association. *Diabetes Care* 2017;40:136–154
- Bao XH, Wong V, Wang Q, Low LC. Prevalence of peripheral neuropathy with insulin-dependent diabetes mellitus. *Pediatr Neurol* 1999;20:204–209
- Nordwall M, Hyllienmark L, Ludvigsson J. Early diabetic complications in a population of young patients with type 1 diabetes mellitus despite intensive treatment. *J Pediatr Endocrinol Metab* 2006;19:45–54
- Blankenburg M, Kraemer N, Hirschfeld G, et al. Childhood diabetic neuropathy: functional impairment and non-invasive screening assessment. *Diabet Med* 2012;29:1425–1432
- Weintrob N, Amitay I, Lilos P, Shalitin S, Lazar L, Josefsberg Z. Bedside neuropathy disability score compared to quantitative sensory testing for measurement of diabetic neuropathy in children, adolescents, and young adults with type 1 diabetes. *J Diabetes Complications* 2007;21:13–19
- Olsen BS, Johannesen J, Sjølie AK, et al.; Danish Study Group of Diabetes in Childhood. Metabolic control and prevalence of microvascular complications in young Danish patients with type 1 diabetes mellitus. *Diabet Med* 1999;16:79–85
- Eppens MC, Craig ME, Cusumano J, et al. Prevalence of diabetes complications in adolescents with type 2 compared with type 1 diabetes. *Diabetes Care* 2006;29:1300–1306
- Jaiswal M, Lauer A, Martin CL, et al.; SEARCH for Diabetes in Youth Study Group. Peripheral neuropathy in adolescents and young adults with type 1 and type 2 diabetes from the SEARCH for Diabetes in Youth follow-up cohort: a pilot study. *Diabetes Care* 2013;36:3903–3908
- Dabelea D, Stafford JM, Mayer-Davis EJ, et al.; SEARCH for Diabetes in Youth Research Group. Association of type 1 diabetes vs type 2 diabetes diagnosed during childhood and adolescence with complications during teenage years and young adulthood. *JAMA* 2017;317:825–835
- Myers GL, Kimberly MM, Waymack PP, Smith SJ, Cooper GR, Sampson EJ. A reference method laboratory network for cholesterol: a model for standardization and improvement of clinical laboratory measurements. *Clin Chem* 2000;46:1762–1772
- Hamman RF, Bell RA, Dabelea D, et al.; SEARCH for Diabetes in Youth Study Group. The SEARCH for Diabetes in Youth study: rationale, findings, and future directions. *Diabetes Care* 2014;37:3336–3344
- Dabelea D, Pihoker C, Talton JW, et al.; SEARCH for Diabetes in Youth Study. Etiological approach to characterization of diabetes type: the SEARCH for Diabetes in Youth study. *Diabetes Care* 2011;34:1628–1633
- Feldman EL, Stevens MJ, Thomas PK, Brown MB, Canal N, Greene DA. A practical two-step quantitative clinical and electrophysiological assessment for the diagnosis and staging of diabetic neuropathy. *Diabetes Care* 1994;17:1281–1289
- Herman WH, Pop-Busui R, Braffett BH, et al.; DCCT/EDIC Research Group. Use of the Michigan Neuropathy Screening Instrument as a measure of distal symmetrical peripheral neuropathy in type 1 diabetes: results from the Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications. *Diabet Med* 2012;29:937–944
- Diabetes Control and Complications Trial Research Group. Effect of intensive diabetes treatment on the development and progression of long-term complications in adolescents with insulin-dependent diabetes mellitus: Diabetes Control and Complications Trial. *J Pediatr* 1994;125:177–188
- UK Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet* 1998;352:837–853
- Cameron NE, Eaton SE, Cotter MA, Tesfaye S. Vascular factors and metabolic interactions in the pathogenesis of diabetic neuropathy. *Diabetologia* 2001;44:1973–1988
- Callaghan BC, Little AA, Feldman EL, Hughes RA. Enhanced glucose control for preventing and treating diabetic neuropathy. *Cochrane Database Syst Rev* 2012;6:CD007543
- Callaghan BC, Hur J, Feldman EL. Diabetic neuropathy: one disease or two? *Curr Opin Neurol* 2012;25:536–541
- Kershner AK, Daniels SR, Imperatore G, et al. Lipid abnormalities are prevalent in youth with type 1 and type 2 diabetes: the SEARCH for Diabetes in Youth study. *J Pediatr* 2006;149:314–319
- Rodriguez BL, Fujimoto WY, Mayer-Davis EJ, et al. Prevalence of cardiovascular disease risk factors in U.S. children and adolescents with diabetes: the SEARCH for Diabetes in Youth study. *Diabetes Care* 2006;29:1891–1896
- Petitti DB, Imperatore G, Palla SL, et al.; SEARCH for Diabetes in Youth Study Group. Serum lipids and glucose control: the SEARCH for Diabetes in Youth study. *Arch Pediatr Adolesc Med* 2007;161:159–165
- Tesfaye S, Chaturvedi N, Eaton SE, et al.; EURODIAB Prospective Complications Study Group. Vascular risk factors and diabetic neuropathy. *N Engl J Med* 2005;352:341–350
- Maser RE, Steenkiste AR, Dorman JS, et al. Epidemiological correlates of diabetic neuropathy. Report from Pittsburgh Epidemiology of Diabetes Complications Study. *Diabetes* 1989;38:1456–1461
- Barter PJ, Nicholls S, Rye KA, Anantharamaiah GM, Navab M, Fogelman AM. Antiinflammatory properties of HDL. *Circ Res* 2004;95:764–772
- Patel S, Drew BG, Nakhla S, et al. Reconstituted high-density lipoprotein increases plasma high-density lipoprotein anti-inflammatory properties and cholesterol efflux capacity in patients with type 2 diabetes. *J Am Coll Cardiol* 2009;53:962–971
- Drew BG, Rye KA, Duffy SJ, Barter P, Kingwell BA. The emerging role of HDL in glucose metabolism. *Nat Rev Endocrinol.* 2012;8:237–245
- Callaghan B, Feldman E. The metabolic syndrome and neuropathy: therapeutic challenges and opportunities. *Ann Neurol* 2013;74:397–403
- Clair C, Cohen MJ, Eichler F, Selby KJ, Rigotti NA. The effect of cigarette smoking on diabetic peripheral neuropathy: a systematic review and meta-analysis. *J Gen Intern Med* 2015;30:1193–1203
- American Diabetes Association. *Standards of Medical Care in Diabetes—2016*. *Diabetes Care* 2016;39(Suppl. 1):S1–S112