



Resolution of Hypoglycemia and Cardiovascular Dysfunction After Rituximab Treatment of Insulin Autoimmune Syndrome

Diabetes Care 2017;40:e80–e82 | <https://doi.org/10.2337/dc17-0264>

David Church,^{1,2,3} Robert W. Hunter,⁴ Marcus Lyall,^{4,5} Catriona Clarke,⁶ A.D. Bastiaan Vliegthart,⁴ James W. Dear,⁴ Robert Semple,^{1,2} Neeraj Dhaun,⁴ and Anna R. Dover^{4,5}

History and Examination

A 37-year-old female presented to the emergency department after having collapsed; venous glucose measured 50 mg/dL. She reported recurrent dizziness since the birth of her fifth child 10 months previously and had recorded low capillary blood glucose (36–52 mg/dL) when symptomatic. Her history included gestational diabetes during the most recent pregnancy, treated with insulin aspart, insulin glargine, and metformin. These therapies had been discontinued 1 month postpartum.

Investigation

Blood count and renal, liver, thyroid, and adrenal function were all normal. HbA_{1c} was 41 mmol/mol (5.9%). Continuous glucose monitoring (CGM) demonstrated early-morning (asymptomatic) hypoglycemia and postprandial hyperglycemia. During hypoglycemia the patient had high circulating insulin (39,181 pmol/L; C-peptide 1,046 pmol/L). Insulin assay was performed as previously described (1). Insulin eluted from a gel filtration column in high-molecular weight fractions (Fig. 1C). Serum anti-insulin IgG

concentration was 171 mg/L (reference range 0–5).

We diagnosed insulin autoimmune syndrome (IAS), a term commonly used in patients without previous exposure to exogenous insulin. Here, we cannot exclude that the insulin antibodies generated were in response to exogenous insulin, but severe hypoglycemia did not develop until many months after cessation of insulin therapy. Moreover, antibodies developing after exposure to exogenous insulin rarely bind insulin with sufficient capacity or affinity to perturb glycemia (2).

The patient was fitted with CGM with a hypoglycemia alarm and prescribed a diet of frequent low-glycemic index carbohydrate meals. As significant hypoglycemia continued, prednisone was commenced (1 mg/kg/day) and titrated according to capillary blood glucose readings to 10 mg/day over 3 months. To reduce anti-insulin antibodies, we gave the anti-CD20 monoclonal antibody rituximab.

Repeat CGM after 6 weeks revealed intermittent hypoglycemia and sustained daytime hyperglycemia. Over months, there were reductions in total insulin,

anti-insulin antibody concentration, and antibody-bound insulin (Fig. 1A and B). These reductions were associated with reduced hypoglycemia and improved hypoglycemic awareness. By 6 months, hypoglycemia was rare and postprandial hyperglycemia had improved (CGM peak glucose 162.0 mg/dL). There were no adverse events and prednisone was discontinued after 10 months, after which no further hypoglycemia was recorded.

Recurrent hypoglycemia is associated with endothelial dysfunction, inflammation, and increased cardiovascular risk (3). Thus, we explored the cardiovascular phenotype associated with IAS before and after treatment. Carotid-femoral pulse wave velocity was 7.6 m/s at presentation and 5.2 m/s at 6 months (5.2–8.0 m/s); there was no change in blood pressure over this period. This suggests elevated arterial stiffness at presentation, associated with cardiovascular risk. Other surrogate markers of endothelial dysfunction were all higher at disease presentation than at 6 months (Fig. 1D). Circulating miR-126 rose following treatment (Fig. 1D). This microRNA is endothelial enriched and is thought

¹University of Cambridge Metabolic Research Laboratories, Wellcome Trust-MRC Institute of Metabolic Science, Cambridge, UK

²National Institute for Health Research Cambridge Biomedical Research Centre, Cambridge, UK

³The Pathology Partnership, Department of Clinical Biochemistry and Immunology, Addenbrooke's Hospital, Cambridge, UK

⁴University/British Heart Foundation Centre for Cardiovascular Science, The Queen's Medical Research Institute, University of Edinburgh, Edinburgh, UK

⁵Edinburgh Centre for Endocrinology and Diabetes, Royal Infirmary of Edinburgh, Edinburgh, UK

⁶Department of Clinical Biochemistry, Western General Hospital, Edinburgh, UK

Corresponding author: Robert W. Hunter, robert.hunter@ed.ac.uk.

Received 3 February 2017 and accepted 28 March 2017.

D.C., R.W.H., and M.L. contributed equally to this study.

© 2017 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <http://www.diabetesjournals.org/content/license>.




Figure 1—A: Reportable insulin concentration, total insulin estimation, and free insulin estimation against time. Day 0 is the day of first presentation. Prednisone was commenced on day 44, and rituximab was administered on days 44 and 58. Insulin concentrations shown here were measured using the DiaSorin LIAISON assay; an alternative assay (Abbott Architect) gave consistent results. Follow-up insulin determination was undertaken on neat plasma (“measured insulin”) and after 1:49 dilution in 0.9% saline to promote insulin-antibody dissociation and reduce negative assay interference by antibodies (“total insulin”), as well as in supernatant following polyethylene glycol precipitation (“free insulin”). The open circle represents measured insulin concentration >3,000 pmol/L in neat plasma. B: Serum anti-insulin IgG concentration (in-house ImmunoCAP assay; reference range 0–5). C: Changes in plasma macroinsulin in response to immunosuppressive therapy. At presentation, only 4% of total immunoreactive insulin was recovered from plasma supernatant following polyethylene glycol precipitation, consistent with the presence of high-molecular weight insulin immunoreactivity. Predominantly high-molecular weight insulin consistent with macroinsulin was demonstrable using gel filtration chromatography at presentation. The elution volumes of immunoglobulin and monomeric insulin are shown by the black and white arrows, respectively; the majority of insulin coeluted with immunoglobulins. Follow-up investigations on day 271 confirmed a decrease in macroinsulin. D: Changes in circulating markers of endothelial/vascular function at presentation (“pre”) and at 6 months (“post”).

to maintain endothelial homeostasis and promote vasculogenesis (4). There was minimal change in control miR-122-5p.

Conclusions

Treatment of IAS is poorly defined. Historically, glucocorticoids and plasmapheresis were used for refractory cases. We show that B-cell depletion with rituximab induces a sustained reduction in anti-insulin antibodies, circulating insulin, and the frequency of hypoglycemia. Rituximab has been used successfully in two other cases of IAS (5,6). However, in one, concomitant use of plasmapheresis, methotrexate, and intravenous immunoglobulin makes it difficult to ascribe the beneficial therapeutic effect to rituximab.

Our report is novel in providing data to suggest an adverse vascular phenotype in IAS that is reversible when dysglycemia resolves. We speculate that recurrent dysglycemia may contribute to vascular dysfunction in IAS, but further study is required to determine the underlying mechanism.

Funding. D.C. is funded by a Diabetes Research & Wellness Foundation Sutherland-Earl Clinical Fellowship (RG68554). R.W.H. is supported by the Wellcome Trust–University of Edinburgh Institutional Strategic Support Fund. N.D. is supported by a British Heart Foundation Intermediate Clinical Research Fellowship (FS/13/30/29994). R.S. is funded by the Wellcome Trust (grant WT098498). **Duality of Interest.** No potential conflicts of interest relevant to this article were reported.

Author Contributions. D.C., R.W.H., and M.L. performed the study, analyzed the data, and drafted the manuscript. C.C., A.D.B.V., J.W.D., R.S., N.D., and A.R.D. analyzed the results. All authors revised the manuscript and approved the final version. A.R.D. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

References

- Church D, Cardoso L, Bradbury S, et al. Diagnosis of insulin autoimmune syndrome using polyethylene glycol precipitation and gel filtration chromatography with ex vivo insulin exchange. *Clin Endocrinol (Oxf)* 2017;86:347–353
- Trabucchi A, Iacono RF, Guerra LL, et al. Characterization of insulin antibodies by surface plasmon resonance in two clinical cases: brittle diabetes and insulin autoimmune syndrome. *PLoS One* 2013;8:e84099

3. Joy NG, Tate DB, Younk LM, Davis SN. Effects of acute and antecedent hypoglycemia on endothelial function and markers of atherothrombotic balance in healthy humans. *Diabetes* 2015;64:2571–2580
4. Wang S, Aurora AB, Johnson BA, et al. The endothelial-specific microRNA miR-126 governs vascular integrity and angiogenesis. *Dev Cell* 2008;15:261–271
5. Jassam N, Amin N, Holland P, et al. Analytical and clinical challenges in a patient with concurrent type 1 diabetes, subcutaneous insulin resistance and insulin autoimmune syndrome. *Endocrinol Diabetes Metab Case Rep* 2014;2014:130086
6. Saxon DR, McDermott MT, Michels AW. Novel management of insulin autoimmune syndrome with rituximab and continuous glucose monitoring. *J Clin Endocrinol Metab* 2016;101:1931–1934