



Incidence Rates of Type 2 Diabetes in People With Impaired Fasting Glucose (ADA vs. WHO Criteria) and Impaired Glucose Tolerance: Results From an Older Population (KORA S4/F4/FF4 Study)

Diabetes Care 2019;42:e18–e20 | <https://doi.org/10.2337/dc18-1473>

Bernd Kowall,¹
 Wolfgang Rathmann,^{2,3}
 Brenda Bongaerts,^{2,3} Oliver Kuss,^{2,3,4}
 Andreas Stang,¹ Michael Roden,^{3,5,6}
 Christian Herder,^{3,6}
 Wolfgang Koenig,^{7,8} Cornelia Huth,^{3,9}
 Margit Heier,⁹ Barbara Thorand,^{3,9}
 Karl-Heinz Ladwig,^{3,9,10} Rolf Holle,¹¹
 Christa Meisinger,¹² and
 Annette Peters^{3,9}

The American Diabetes Association (ADA) Expert Committee introduced impaired fasting glucose (IFG: 100–125 mg/dL [5.6–6.9 mmol/L]) as a category with an increased risk for diabetes (1). The World Health Organization (WHO) defined a higher IFG cut point of 110 mg/dL (6.1 mmol/L) because using the lower cut point would substantially increase IFG prevalence and there was no evidence it would lead to reduced progression to diabetes (2).

Our aim was to estimate the long-term incidence of type 2 diabetes in people with isolated IFG (i-IFG), isolated impaired glucose tolerance (i-IGT), and combined IFG/IGT. Contrary to previous studies, we split IFG into two subcategories (i-IFG_{low}: 100 to <110 mg/dL; i-IFG_{high}: 110 to <126 mg/dL) according to the ADA and WHO cut points. We used oral glucose tolerance test (OGTT) data from the Cooperative Health Research in the Region of Augsburg (KORA) S4/F4/FF4 cohort study collected on 1,231

participants without known diabetes aged 55–74 years at baseline (1999–2001) and in two follow-up examinations (2006–2008 and 2013–2014) (3). Incident type 2 diabetes was defined as fasting glucose ≥ 126 mg/dL, 2-h glucose ≥ 200 mg/dL, or a type 2 diabetes diagnosis confirmed by a physician. We included 913 people (74%) with an OGTT or physician-confirmed type 2 diabetes at one or both follow-up investigations. Participants who died after baseline were either excluded if no data on their glycemic status during follow-up were available or censored if these data were available at the first follow-up examination. The mean follow-up time was 10.2 years. Self-reported age at diabetes diagnosis was used for calculation of person-years with physician-diagnosed diabetes during follow-up. Person-years were from baseline to the earliest of diagnosis of diabetes or last study date. For the binary response “incident diabetes” a log-linear model with a Poisson

working likelihood and robust standard errors was fitted to estimate relative risks (RRs) with 95% CIs adjusted for age, sex, BMI, physical activity, smoking, and hypertension. To additionally account for the time until incident diabetes, we fitted an adjusted accelerated failure time (AFT) model for interval-censored data with a log-logistic distribution assumed for the event times. The expected time until incident diabetes for the prediabetes categories is e^{β} (β regression coefficient) times the expected time until diabetes for a person with normal glucose tolerance (NGT).

Table 1 shows the baseline characteristics and incidence rates of type 2 diabetes by OGTT categories. The mean age \pm SD of participants at baseline was 63 ± 5 years (51% men). Compared with the NGT group, people with prediabetes were older, included larger proportions of men (except in the i-IGT category), had higher BMI, and had a higher prevalence of hypertension. There were no

¹Center of Clinical Epidemiology, Institute for Medical Informatics, Biometry and Epidemiology, Medical Faculty, University Duisburg-Essen, Essen, Germany

²Institute for Biometrics and Epidemiology, German Diabetes Center, Leibniz Center for Diabetes Research at Heinrich Heine University, Düsseldorf, Germany

³German Center for Diabetes Research (DZD), München-Neuherberg, Germany

⁴Institute of Medical Statistics, Medical Faculty, Heinrich Heine University, Düsseldorf, Germany

⁵Division of Endocrinology and Diabetology, Medical Faculty, Heinrich Heine University, Düsseldorf, Germany

⁶Institute for Clinical Diabetology, German Diabetes Center, Leibniz Center for Diabetes Research at Heinrich Heine University, Düsseldorf, Germany

⁷Department of Internal Medicine II—Cardiology, University of Ulm Medical Center, Ulm, Germany

⁸Deutsches Herzzentrum München, Technische Universität München, and German Centre for Cardiovascular Research (DZHK), partner site Munich Heart Alliance, Munich Germany

⁹Institute of Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany

¹⁰Department of Psychosomatic Medicine and Psychotherapy, Klinikum rechts der Isar, Technische Universität München, Munich, Germany

¹¹Institute of Health Economics and Health Care Management, Helmholtz Zentrum München, Neuherberg, Germany

¹²Ludwig-Maximilians-Universität München, University Center for Health Sciences at the Klinikum Augsburg (UNIKA-T), Augsburg, Germany

Corresponding author: Wolfgang Rathmann, wolfgang.rathmann@ddz.de

Received 10 July 2018 and accepted 7 November 2018

B.K. and W.R. share first authorship and C.M. and A.P. share senior authorship.

© 2018 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <http://www.diabetesjournals.org/content/license>.

Table 1—Incidence rates and adjusted RRs for type 2 diabetes by glycemic status at baseline (IFG and IGT), with IFG split into two categories, i-IFG_{low} (100 to <110 mg/dL) and i-IFG_{high} (110 to <126 mg/dL): the KORA S4/F4/FF4 study^a

OGTT category	N (%)	Age, years (% men)	BMI (kg/m ²)	Hypertension ^b	Current smokers	Physically inactive ^c	Incident diabetes, n (%)	Incidence rate per 1,000 py (95% CI)	RR (95% CI) ^d	Exp(β) from AFT model (95% CI) ^{d,e}
NGT	463 (51)	62.7 ± 5.4 (43)	27.3 ± 4.0	39.9	14.9	50.5	29 (6.3)	5.8 (3.9–8.3)	1.0	1.0
i-IFG _{low} (100 to <110 mg/dL)	202 (22)	63.0 ± 5.4 (58)	28.4 ± 4.0	51.2	11.4	48.8	30 (14.9)	14.4 (9.7–20.5)	2.0 (1.2–3.3)	0.79 (0.66–0.93)
i-IFG _{high} (110 to <126 mg/dL)	75 (8)	63.1 ± 4.9 (69)	29.5 ± 3.7	62.7	9.3	60.0	34 (45.3)	47.4 (32.8–66.2)	5.6 (3.6–8.8)	0.52 (0.43–0.63)
i-IGT	63 (7)	65.6 ± 5.4 (41)	29.0 ± 4.1	69.8	6.4	58.7	21 (33.3)	35.2 (21.8–53.8)	4.3 (2.6–7.2)	0.56 (0.46–0.69)
Combined IFG _{low} (100 to <110 mg/dL) and IGT	59 (6)	64.5 ± 5.3 (58)	30.2 ± 4.3	59.3	3.4	54.2	27 (45.8)	52.2 (34.4–75.9)	5.5 (3.5–8.8)	0.46 (0.38–0.56)
Combined IFG _{high} (110 to <126 mg/dL) and IGT	51 (6)	63.8 ± 4.8 (63)	29.4 ± 3.6	70.6	15.7	64.7	30 (58.8)	76.0 (51.3–108.5)	7.2 (4.6–11.3)	0.37 (0.30–0.46)
Total	913 (100)	63.2 ± 5.4 (51)	28.1 ± 4.1	49.3	12.4	52.5	171 (18.7)	18.3 (15.6–21.2)	—	—

Data are means ± SD or percentages, unless otherwise indicated. The i-IFG group had normal 2-h glucose; the i-IGT group had normal fasting glucose. py, person-years. ^aOverall mean follow-up was 10.2 years. Follow-up was 10.9 years for NGT, 10.4 years for i-IFG_{low}, 9.6 years for i-IFG_{high}, 9.5 years for i-IGT, 8.8 years for IFG_{low}/IGT, and 7.7 years for IFG_{high}/IGT. ^bSystolic/diastolic blood pressure ≥140/90 mmHg and/or use of antihypertensive medication if the subjects were aware of being hypertensive. ^cLess than 1 h of leisure-time physical activity per week in summer or winter. ^dAdjusted for age, sex, BMI, hypertension, physical activity, and smoking. ^eFor example, for i-IGT exp(β) 0.56 means the expected time until incident diabetes for people in the i-IGT group is the expected time until diabetes for people with NGT × 0.56.

consistent differences in current smoking between the prediabetes and NGT groups, whereas low physical activity was more often reported in prediabetes (except in i-IFG_{low}).

When compared with the NGT group, the risk of type 2 diabetes was increased in all prediabetes categories. RR ranged from 2.0 (95% CI 1.2–3.3) in i-IFG_{low} to 5.6 (3.6–8.8) in i-IFG_{high} and 7.2 (4.6–11.3) in combined IFG_{high}/IGT. In the latter group, more than half of the people developed diabetes, whereas in those with i-IFG_{low} only 15% progressed to diabetes. All prediabetes groups were associated with a reduced time to incident diabetes compared with NGT. The expected time until diabetes for i-IGT_{low} was 79% of the expected time until diabetes for NGT, and for combined IFG_{high}/IGT it was 37%. i-IFG_{high} was related to a faster time to diabetes than i-IFG_{low}. Finally, the time to incident diabetes of i-IFG_{high} and i-IGT was similar.

In an additional analysis, we split i-IFG_{low} (100–109 mg/dL) into two categories: i-IFG (100–104 mg/dL) and i-IFG (105–109 mg/dL). The RR in these groups was 1.57 (95% CI 0.86–2.87) and 2.80 (1.57–4.99), respectively; in the AFT model, e^β was 0.84 (95% CI 0.69–1.04) and 0.72 (0.58–0.90), respectively.

Our study gives several insights. First, 10-year diabetes incidences >40% were only found in i-IFG_{high} and in combined IFG/IGT (for both IFG_{low} and IFG_{high}). Second, the OGTT is helpful for identifying high-risk groups for type 2 diabetes. For example, diabetes risk was much stronger in IFG_{low} when IGT co-occurred. Third, among participants with i-IFG_{low} only 15% progressed to diabetes, but as this group was large (prevalence 22%), the absolute number of persons with incident diabetes from this subgroup equaled the number from other prediabetes categories with higher diabetes risk. Generally, the numbers of incident diabetes cases were similar for all categories of glycemic status including NGT, indicating that population-based approaches should complement high-risk approaches in diabetes prevention. For all prediabetes categories, population-attributable risks lay between 0.18 and 0.27; the small range indicates again that all prediabetes categories contribute similarly to the diabetes burden. Fourth, i-IFG (100–109 mg/dL) is not a homogenous category, and there is a strong increase in

diabetes risk at the 105 mg/dL cut point. This is in line with results by Zhuo et al. (4), who did an economic analysis of fasting plasma glucose (FPG) thresholds and suggested a threshold of 105 mg/dL to identify persons at high risk of type 2 diabetes (4). Our data support model-based estimates in the U.S., which suggested that a great reduction in the absolute number of diabetes cases would be achieved through implementation of a structured lifestyle intervention in people with IFG_{high/low} (5). However, it must be noted that evidence for lifestyle intervention mainly comes from studies of people with IGT rather than IFG. Finally, our study shows that all prediabetes categories have an unfavorable risk profile (e.g., obesity, hypertension). Thus, those affected are at a higher cardiovascular risk.

In conclusion, in our older cohort—for younger ones results may be different—we found that persons with fasting glucose values of 100 to <110 mg/dL have a twofold higher long-term risk to progress to diabetes than those with NGT and that i-IFG_{low} is almost three times more prevalent than i-IFG_{high} with implications for the expected number of incident diabetes cases in this population. Lowering FPG thresholds and including more people into prevention programs leads to a gain in quality-

adjusted life years but also to an increase in costs and a reduction in cost-effectiveness. Therefore, selection of an FPG cut point depends on how many health care resources are accepted to be put toward prevention.

Funding. This work was supported by the Ministry for Culture and Science of the State of North Rhine-Westphalia and the German Federal Ministry of Health. The diabetes part of the KORA study was partly funded by a grant from Deutsche Forschungsgemeinschaft (RA 459/3-1). The KORA study was initiated and financed by Helmholtz Zentrum München—German Research Center for Environmental Health, which is funded by the German Federal Ministry of Education and Research (BMBF) and by the State of Bavaria. A.S. received a grant from the German Federal Ministry of Education and Research (grant 01ER1704).

Duality of Interest. W.R. reported receiving consulting fees for attending educational sessions or advisory boards from AstraZeneca, Boehringer Ingelheim, Eli Lilly, and Novo Nordisk and institutional research grants from Novo Nordisk outside the submitted work. W.K. reports personal fees from AstraZeneca, Novartis, Pfizer, The Medicines Company, GlaxoSmithKline, DalCor, Sanofi, Berlin-Chemie, Kowa, and Amgen and grants and nonfinancial support from Roche Diagnostics, Beckman (Krefeld, Germany), Singulex, and Abbott outside the submitted work. No other potential conflicts of interest relevant to this article were reported.

Author Contributions. B.K. and W.R. conceived and designed the study. W.R. drafted the manuscript. B.K. analyzed the data. All authors acquired and interpreted the data, critically reviewed the manuscript, provided technical or material support, and approved the final version to be published. W.R. is the guarantor of this work and, as such, had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

References

1. American Diabetes Association. 2. Classification and diagnosis of diabetes: *Standards of Medical Care in Diabetes—2018*. *Diabetes Care* 2018;41(Suppl. 1):S13–S27
2. World Health Organization, International Diabetes Federation. *Definition and Diagnosis of Diabetes Mellitus and Intermediate Hyperglycemia: Report of a WHO/IDF Consultation*. Geneva, World Health Organization, 2006
3. Rathmann W, Haastert B, Icks A, et al. High prevalence of undiagnosed diabetes mellitus in Southern Germany: target populations for efficient screening. *The KORA survey 2000*. *Diabetologia* 2003;46:182–189
4. Zhuo X, Zhang P, Kahn HS, Gregg EW. Cost-effectiveness of alternative thresholds of the fasting plasma glucose test to identify the target population for type 2 diabetes prevention in adults aged ≥ 45 years. *Diabetes Care* 2013;36:3992–3998
5. Gregg EW, Boyle JP, Thompson TJ, Barker LE, Albright AL, Williamson DF. Modeling the impact of prevention policies on future diabetes prevalence in the United States: 2010–2030. *Popul Health Metr* 2013;11:18