



# BMI, Weight Change, and Dementia Risk in Patients With New-Onset Type 2 Diabetes: A Nationwide Cohort Study

*Diabetes Care* 2019;42:1217–1224 | <https://doi.org/10.2337/dc18-1667>

Ga Eun Nam,<sup>1</sup> Yong Gyu Park,<sup>2</sup> Kyungdo Han,<sup>2</sup> Mee Kyoung Kim,<sup>3</sup> Eun Sil Koh,<sup>4</sup> Eun Sook Kim,<sup>5</sup> Min-Kyung Lee,<sup>6</sup> Bongsung Kim,<sup>7</sup> Oak-kee Hong,<sup>8</sup> and Hyuk-Sang Kwon<sup>3</sup>

## OBJECTIVE

This study examined the association between baseline BMI, percentage weight change, and the risk of dementia in patients newly diagnosed with type 2 diabetes.

## RESEARCH DESIGN AND METHODS

Using the South Korean National Health Insurance Service-National Health Screening Cohort database, we identified 167,876 subjects aged  $\geq 40$  years diagnosed with new-onset type 2 diabetes between 2007 and 2012. Their weight changes were monitored for  $\sim 2$  years after diagnosis, with follow-up assessments occurring for an average of 3.5 years. The hazard ratios (HRs) and Bonferroni-adjusted 95% CIs of all-cause dementia, Alzheimer disease (AD), and vascular dementia were estimated using multivariable Cox proportional hazards regression models.

## RESULTS

We identified 2,563 incident dementia cases during follow-up. Baseline BMI among patients with new-onset type 2 diabetes was inversely associated with the risk of all-cause dementia and AD, independent of confounding variables ( $P$  for trend  $< 0.001$ ). The percentage weight change during the 2 years after a diagnosis of type 2 diabetes showed significant U-shaped associations with the risk of all-cause dementia development ( $P < 0.001$ ); the HRs of the disease increased significantly when weight loss or gain was  $> 10\%$  (1.34 [95% CI 1.11–1.63] and 1.38 [1.08–1.76], respectively). Additionally, weight loss  $> 10\%$  was associated with an increased risk of AD (HR 1.26 [95% CI 1.01–1.59]).

## CONCLUSIONS

A lower baseline BMI was associated with increased risks of all-cause dementia and AD in patients with new-onset type 2 diabetes. Weight loss or weight gain after the diagnosis of diabetes was associated with an increased risk of all-cause dementia. Weight loss was associated with an increased risk of AD.

Type 2 diabetes and dementia are major chronic diseases with increasing prevalence and are expected to have substantial impacts on the aging population (1,2). Recent evidence implies that type 2 diabetes is associated with an increased risk of cognitive decline and incident dementia (3–8). Although the underlying mechanisms remain unclear, it has been hypothesized that cerebrovascular pathologies are influenced by hyperglycemia and insulin resistance (9,10). Moreover, any specific treatment for dementia in patients with type 2 diabetes has been unknown. It is therefore important

<sup>1</sup>Department of Family Medicine, Korea University Anam Hospital, Korea University College of Medicine, Seoul, Republic of Korea

<sup>2</sup>Department of Biostatistics, College of Medicine, The Catholic University of Korea, Seoul, Republic of Korea

<sup>3</sup>Division of Endocrinology and Metabolism, Department of Internal Medicine, Yeouido St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Seoul, Republic of Korea

<sup>4</sup>Division of Nephrology, Department of Internal Medicine, Yeouido St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Incheon, Republic of Korea

<sup>5</sup>Division of Endocrinology and Metabolism, Department of Internal Medicine, Incheon St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Incheon, Republic of Korea

<sup>6</sup>Division of Endocrinology and Metabolism, Department of Internal Medicine, Myongji Hospital, Hanyang University Medical Center, Gyeonggi-do, Republic of Korea

<sup>7</sup>Department of Statistics and Actuarial Science, Soongsil University, Seoul, Republic of Korea

<sup>8</sup>Department of Internal Medicine, College of Medicine, The Catholic University of Korea, Seoul, Republic of Korea

Corresponding authors: Hyuk-Sang Kwon, [drkwon@catholic.ac.kr](mailto:drkwon@catholic.ac.kr), and Yong Gyu Park, [ygpark@catholic.ac.kr](mailto:ygpark@catholic.ac.kr)

Received 4 August 2018 and accepted 31 March 2019

This article contains Supplementary Data online at <http://care.diabetesjournals.org/lookup/suppl/doi:10.2337/dc18-1667/-/DC1>.

© 2019 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <http://www.diabetesjournals.org/content/license>.

to identify risk factors for cognitive decline and dementia in patients with type 2 diabetes.

In the general population, there is a growing interest in the role of simple and feasible measures of adiposity in relation to predicting dementia development (11–14). High adiposity can lead to cognitive decline in late life through vascular and dysmetabolic pathways, and accumulation of brain lesions through these pathways may further increase the risk of dementia with their longer duration throughout the life course (15). Thus, the predictive roles of BMI and weight change for the onset of Alzheimer disease (AD) have been extensively investigated. However, methodological differences have resulted in discrepancies among studies regarding the effects of BMI and weight change on cognitive prognoses. Moreover, these studies have predominantly been conducted in Western nations, whereas obesity parameters and the frequency of dementia differ across ethnic groups (16). Although these parameters have clinical implications in the prevention and treatment of type 2 diabetes (17), there are limited data on the association between baseline BMI, weight change, and the risk of incident dementia in patients with type 2 diabetes. We conducted a nationwide population-based cohort study to explore the association of baseline BMI and weight change with the risk of dementia in patients newly diagnosed with type 2 diabetes.

## RESEARCH DESIGN AND METHODS

### Study Subjects

We used a database provided by the South Korean National Health Insurance Service (NHIS), which is a single insurer managed by the South Korean government. The NHIS manages a mandatory universal insurance system that covers ~97% of the South Korean population and provides biannual health examinations for all insured Koreans. Hence, the South Korean NHIS retains an extensive data set of 50,000,000 Koreans that includes patient demographics, medical treatments and procedures, and disease diagnoses according to the ICD-10-CM codes. Since 2015, the South Korean NHIS has released a nationally representative data set that includes nearly the entire Korean population and is open to all researchers whose study protocols are approved by an official review committee.

To identify patients with new-onset type 2 diabetes among all individuals ( $n = 25,315,068$ ) who had undergone health examinations between 1 January 2007 and 31 December 2012, we excluded individuals aged  $<40$  years ( $n = 6,928,382$ ), those with a fasting plasma glucose (FPG) level  $<126$  mg/dL at the first health examination ( $n = 17,415,192$ ), and those with a prior history of diabetes ( $n = 659,776$ ). We then excluded individuals without a follow-up examination within 2 years ( $n = 136,215$ ), those diagnosed with cancer ( $n = 5,528$ ), those with a prior diagnosis of dementia between 2002 and enrollment ( $n = 979$ ), and those with any missing variables ( $n = 1,120$ ). Finally, 167,876 individuals with new-onset type 2 diabetes (108,025 men and 59,851 women) with 2 years of body weight change measurements were enrolled in this study (Fig. 1). Participants were tracked until 31 December 2015. The mean follow-up duration was  $3.5 \pm 1.7$  years. This study was approved by the The Catholic University of Korea Institutional Review Board (No. SC18ZESI0022).

### Definition of Dementia

Dementia was diagnosed based on the relevant ICD-10-CM codes (F00 or G30 for AD; F01 for vascular dementia [VD]; and F02, F03, or G31 for other dementia) and the prescription of one or more medications for dementia (donepezil, rivastigmine, galantamine, or memantine). When there were codes for both AD and VD, the main diagnosis was extracted as the final diagnosis. If both codes for AD and VD were as an additional diagnosis, the main diagnosis at the next visit was extracted as the final diagnosis. If there was neither AD nor VD as a main diagnosis up to the second claim database, the diagnosis was defined as “other dementia.”

### BMI and Weight Change

Height, weight, and waist circumference (WC) were measured while participants wore lightweight clothing. BMI ( $\text{kg}/\text{m}^2$ ) was calculated as weight in kilograms divided by the square of height in meters. The World Health Organization recommendations for Asian populations were used to categorize individuals into five BMI groups:  $<18.5$   $\text{kg}/\text{m}^2$




Figure 1—Flowchart of the study population.

(underweight), 18.5–22.9 kg/m<sup>2</sup> (normal), 23.0–24.9 kg/m<sup>2</sup> (overweight), 25.0–29.9 kg/m<sup>2</sup> (class I obese), or ≥30 kg/m<sup>2</sup> (class II obese) (18). Weight change was calculated as the difference in weight from the time of diagnosis of type 2 diabetes until 2 years later. We defined ≤5% weight change as “stable weight” and categorized “weight change” into groups of 5% increments: 1) ≥10% weight loss, 2) 5–10% weight loss, 3) 5–10% weight gain, and 4) ≥10% weight gain.

### Covariates

We assessed subject demographics and lifestyle through standardized self-report questionnaires. Income level was divided into quartile groups. Smoking status was classified as nonsmokers, former smokers, or current smokers. Individuals who consumed ≥30 g/day of alcohol were defined as heavy alcohol drinkers (19). Physical activity was categorized by the frequency of ≥20 min of strenuous exercise (none, ≤4 times per week, or ≥5 times per week).

The health examination provided by the South Korean NHIS included blood pressure (BP) and laboratory measurements. Systolic and diastolic BP was measured with the individual seated after at least 5 min of rest. Blood samples were obtained after overnight fasting to measure serum glucose, total cholesterol, and creatinine levels. Baseline comorbidities, such as hypertension and dyslipidemia, were identified based on combinations of the medical history, health examination results, or ICD-10-CM and prescription codes. Hypertension was defined as BP ≥140/90 mmHg, or at least one claim per year for antihypertensive medication (ICD-10-CM codes I10–I15). Dyslipidemia was defined as a total cholesterol level ≥240 mg/dL or at least one claim per year for lipid-lowering medication (ICD-10-CM code E78). Estimated glomerular filtration rate (eGFR) was calculated using the equation from the Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI), and CKD was defined as an eGFR <60 mL/min/1.73 m<sup>2</sup> (20,21). Ischemic heart disease and stroke were identified based on the answers to the self-report questionnaire for a physician's diagnosis of angina or myocardial infarction, or stroke, respectively. Depressive disorder was defined as at least one claim per year for the ICD-10-CM codes F32 or F33.

### Statistical Analyses

Statistical analyses were conducted using SAS 9.4 software (SAS Institute, Cary, NC). Clinical and demographic characteristics of the study participants were compared according to weight change (%) status using an ANOVA for continuous variables and a  $\chi^2$  test for categorical variables. Data are presented as mean  $\pm$  SD for continuous variables or as number (percentage) for categorical variables. For each participant, the primary outcome between 1 January 2009 and 31 December 2015 was dementia, and the number of person-years of follow-up was counted. All-cause dementia, AD, and VD were assessed as incident dementia during  $3.5 \pm 1.7$  years after the last recorded weighing. Incidence rates of dementia were calculated by dividing the number of events by 1,000 person-years.

Cox proportional hazards regression models were performed to evaluate the association of baseline BMI and weight change with incident dementia, and hazard ratios (HRs) and Bonferroni-adjusted 95% CIs were calculated. Model 1 was unadjusted and model 2 was adjusted for age, sex, smoking status, alcohol consumption, physical activity, and income. Model 3 was additionally adjusted for hypertension, dyslipidemia, CKD, history of stroke, depressive disorder, FPG level, insulin use, and number of oral antidiabetic agents used. Model 3 was further adjusted for baseline BMI in the analysis regarding the association between weight change and dementia risk. Stratified analyses were performed according to age (<65 vs. ≥65 years), sex (men vs. women), and obesity status (BMI <25 kg/m<sup>2</sup> vs. ≥25 kg/m<sup>2</sup>), and interactions between subgroups were tested. In addition, a sensitivity analysis was performed to exclude participants with end points occurring within ≤1 year of the follow-up to account for the possibility of reverse causation. Reported *P* values were two tailed, and *P* < 0.05 was considered statistically significant.

## RESULTS

### Baseline Characteristics of Study Subjects

Our study enrolled 167,876 individuals aged ≥40 years with incident type 2 diabetes. We observed 2,563 events of all-cause dementia (1,846 AD and 381 VD) during an average 3.5 years of follow-up. Table 1 presents the general

characteristics of the study participants on the index date after being monitored for ~2 years. Weight, BMI, WC, systolic and diastolic BP, and FPG levels were higher in subjects with ≥10% weight gain than in subjects with ≥10% weight loss; individuals in the former group were more likely to be current smokers and heavy alcohol drinkers. The proportion of comorbidities, such as hypertension and dyslipidemia, was higher in subjects with ≥10% weight gain than in those with ≥10% weight loss. Conversely, the proportion of CKD and ischemic heart disease was lower in subjects with ≥10% weight gain than in those with ≥10% weight loss.

### Risk of Incident Dementia According to Baseline BMI

Table 2 provides the HRs (95% CIs) of incident dementia according to baseline BMI when subjects were diagnosed with type 2 diabetes. During  $5.5 \pm 1.7$  years of follow-up, after adjusting for age, sex, smoking status, alcohol consumption, physical activity, income, hypertension, dyslipidemia, CKD, stroke, depressive disorder, FPG level, insulin use, and number of oral antidiabetic agents used (model 3), the risk of all-cause dementia decreased significantly in those with BMI of 23–25 kg/m<sup>2</sup> (HR 0.80 [95% CI 0.69–0.91]), 25–30 kg/m<sup>2</sup> (HR 0.77 [95% CI 0.68–0.88]), and ≥30 kg/m<sup>2</sup> (HR 0.79 [95% CI 0.63–0.99]), compared with patients with a normal BMI of 18.5–23 kg/m<sup>2</sup>. The HRs for all-cause dementia tended to increase as BMI decreased (*P* for trend <0.001). The risk of AD was significantly higher for underweight patients (HR 1.59 [95% CI 1.04–2.43]) and also decreased in those with BMI of 23–25 kg/m<sup>2</sup> (HR 0.79 [95% CI 0.67–0.93]) and 25–30 kg/m<sup>2</sup> (HR 0.76 [95% CI 0.65–0.88]) compared with patients with a normal BMI (*P* for trend <0.001). Baseline BMI was not significantly associated with the risk of VD. The sensitivity analysis showed similar results (Supplementary Table 1).

### Risk of Incident Dementia According to Weight Change in the 2 Years After Being Diagnosed With Diabetes

Table 3 reports the HRs (95% CIs) of incident dementia according to the percentage weight change after the diagnosis of type 2 diabetes. After adjusting for all confounding variables, including baseline BMI, the risk of all-cause dementia

**Table 1—General characteristics of study subjects according to weight change status**

|                                  | Weight change (%)    |                               |                             |                           |                    | P      |
|----------------------------------|----------------------|-------------------------------|-----------------------------|---------------------------|--------------------|--------|
|                                  | ≥ -10<br>(n = 6,603) | < -10 to ≥ -5<br>(n = 24,550) | < -5 to <5<br>(n = 115,473) | ≥5 to <10<br>(n = 15,539) | ≥10<br>(n = 5,711) |        |
| Male sex                         | 3,061 (46.4)         | 13,684 (55.7)                 | 76,334 (66.1)               | 10,864 (69.9)             | 4,082 (71.5)       | <0.001 |
| Age (years)                      | 58.9 ± 10.9          | 58.0 ± 9.9                    | 57.1 ± 9.4                  | 56.2 ± 9.5                | 55.9 ± 9.8         | <0.001 |
| Age (years)                      |                      |                               |                             |                           |                    | <0.001 |
| 40–49                            | 1,405 (3.7)          | 5,204 (13.5)                  | 26,078 (67.9)               | 4,119 (10.7)              | 1,622 (4.2)        |        |
| 50–59                            | 2,160 (3.4)          | 8,871 (13.8)                  | 44,986 (70.1)               | 6,001 (9.3)               | 2,202 (3.4)        |        |
| 60–69                            | 1,661 (3.9)          | 6,498 (15.3)                  | 29,511 (69.4)               | 3,659 (8.6)               | 1,188 (2.8)        |        |
| ≥70                              | 1,377 (6.1)          | 3,977 (17.5)                  | 14,898 (65.6)               | 1,760 (7.8)               | 699 (18.5)         |        |
| Height (cm)                      | 160.4 ± 9.4          | 162.0 ± 9.1                   | 163.6 ± 8.7                 | 164.1 ± 8.7               | 164.1 ± 8.7        | <0.001 |
| Weight (kg)                      | 59.2 ± 10.9          | 63.7 ± 10.7                   | 68.6 ± 11.0                 | 70.6 ± 11.6               | 71.3 ± 11.9        | <0.001 |
| BMI (kg/m <sup>2</sup> )         | 22.9 ± 3.1           | 24.2 ± 3.0                    | 25.6 ± 3.1                  | 26.1 ± 3.3                | 26.4 ± 3.5         | <0.001 |
| BMI (kg/m <sup>2</sup> )         |                      |                               |                             |                           |                    | <0.001 |
| <18.5                            | 363 (5.5)            | 332 (1.4)                     | 577 (0.5)                   | 52 (0.3)                  | 20 (0.4)           |        |
| 18.5–23                          | 3,286 (49.8)         | 8,458 (34.5)                  | 21,843 (18.9)               | 2,378 (15.3)              | 838 (14.7)         |        |
| 23–25                            | 1,516 (23.0)         | 7,016 (28.6)                  | 30,247 (26.2)               | 3,468 (22.3)              | 1,188 (20.8)       |        |
| 25–30                            | 1,280 (19.4)         | 7,740 (31.5)                  | 53,349 (46.2)               | 7,858 (50.6)              | 2,853 (50.0)       |        |
| ≥30                              | 159 (2.4)            | 1,004 (4.1)                   | 9,457 (8.2)                 | 1,783 (11.5)              | 812 (14.2)         |        |
| WC (cm)                          | 80.3 ± 8.6           | 82.9 ± 8.0                    | 86.4 ± 8.0                  | 87.8 ± 8.3                | 88.3 ± 8.6         | <0.001 |
| BP (mmHg)                        |                      |                               |                             |                           |                    |        |
| Systolic                         | 123.8 ± 15.1         | 125.4 ± 14.4                  | 127.7 ± 14.4                | 129.1 ± 14.5              | 129.7 ± 14.9       | <0.001 |
| Diastolic                        | 76.3 ± 9.8           | 77.3 ± 9.5                    | 78.9 ± 9.5                  | 79.7 ± 9.5                | 80.1 ± 9.7         | <0.001 |
| Fasting glucose (mg/dL)          | 125.1 ± 44.5         | 129.4 ± 39.7                  | 135.9 ± 37.0                | 138.7 ± 37.7              | 139.3 ± 41.0       | <0.001 |
| Total cholesterol (mg/dL)        | 182.5 ± 40.7         | 187.4 ± 40.5                  | 190.2 ± 40.1                | 188.0 ± 39.4              | 187.4 ± 39.7       | <0.001 |
| Current smoker                   | 1,280 (19.4)         | 5,560 (22.7)                  | 29,986 (26.0)               | 4,334 (27.9)              | 1,637 (28.7)       | <0.001 |
| Heavy alcohol drinker            | 308 (4.7)            | 1,599 (6.5)                   | 10,392 (9.0)                | 1,509 (9.7)               | 582 (10.2)         | <0.001 |
| Regular exerciser                | 3,384 (51.3)         | 13,366 (54.5)                 | 64,547 (56.0)               | 8,430 (54.3)              | 3,020 (52.9)       | <0.001 |
| Income quartiles                 |                      |                               |                             |                           |                    | <0.001 |
| 1                                | 1,775 (26.9)         | 6,076 (24.8)                  | 28,214 (24.4)               | 4,105 (26.4)              | 1,592 (27.9)       |        |
| 2                                | 1,419 (21.5)         | 5,329 (21.7)                  | 25,193 (21.8)               | 3,680 (23.7)              | 1,388 (24.3)       |        |
| 3                                | 1,599 (24.2)         | 6,106 (24.9)                  | 28,649 (24.8)               | 3,927 (25.3)              | 1,464 (25.6)       |        |
| 4                                | 1,810 (27.4)         | 7,039 (28.7)                  | 33,417 (28.9)               | 3,827 (24.6)              | 1,267 (22.2)       |        |
| Comorbidities                    |                      |                               |                             |                           |                    |        |
| Hypertension                     | 3,323 (50.4)         | 12,544 (51.1)                 | 64,927 (56.3)               | 9,142 (58.9)              | 3,401 (59.6)       | <0.001 |
| Dyslipidemia                     | 3,016 (45.8)         | 11,760 (48.0)                 | 59,120 (51.3)               | 8,095 (52.2)              | 2,992 (52.4)       | <0.001 |
| CKD                              | 576 (8.7)            | 1,680 (6.9)                   | 7,535 (6.5)                 | 1,015 (6.5)               | 398 (7.0)          | <0.001 |
| Ischemic heart disease           | 321 (6.9)            | 1,079 (6.1)                   | 4,384 (5.1)                 | 520 (4.4)                 | 189 (4.5)          | <0.001 |
| Stroke                           | 155 (3.4)            | 396 (2.3)                     | 1,526 (1.8)                 | 208 (1.8)                 | 99 (2.4)           | <0.001 |
| Depressive disorder              | 519 (7.9)            | 1,458 (5.9)                   | 5,012 (4.3)                 | 709 (4.6)                 | 293 (5.1)          | <0.001 |
| Insulin use                      | 419 (6.4)            | 1,164 (4.7)                   | 5,011 (4.3)                 | 891 (5.7)                 | 417 (7.3)          | <0.001 |
| ≥2 oral antidiabetic agents used | 2,587 (39.2)         | 9,940 (40.5)                  | 61,872 (53.6)               | 11,399 (73.4)             | 4,367 (76.5)       | <0.001 |

Values are presented as mean ± SD or n (%).

was significantly higher among individuals with ≥10% weight loss (HR 1.34 [95% CI 1.11–1.63]) and ≥10% weight gain (HR 1.38 [95% CI 1.08–1.76]) than among those with <5% weight loss or gain. The correlation between weight change and all-cause dementia among patients with new-onset type 2 diabetes showed a significant U-shaped association ( $P < 0.001$ ). The risk of AD was significantly higher among individuals with ≥10% weight loss (HR 1.26 [95% CI 1.01–1.59]) than among those with <5% weight loss or gain, after correcting for confounding variables. No

significant association was found between weight change and risk of VD. The trends of results were not changed largely in the sensitivity analysis (Supplementary Table 2).

#### Risk of Dementia by Weight Change Status in Subgroups According to Age, Sex, and Baseline BMI

The relationship between weight change and sex in the development of VD was significant, with men with fluctuating weight at greater risk for developing VD than women ( $P = 0.020$ ). Any significant

interactions of weight change with age and obesity status were not shown in the development of all types of dementia (Table 4).

#### CONCLUSIONS

In this population-based longitudinal study, baseline BMI levels among patients with new-onset type 2 diabetes were inversely associated with the risk of all-cause dementia and AD, independent of confounding variables. Weight change during the 2 years after diagnosis showed a significant U-shaped association with the risk of developing all-cause dementia.

**Table 2—HR (95% CI) of incident dementia according to baseline BMI among patients with new-onset type 2 diabetes**

| BMI (kg/m <sup>2</sup> )  | N      | Events | Person-years | Incidence rate* | HR (95% CI) <sup>†</sup> |                  |                  |
|---------------------------|--------|--------|--------------|-----------------|--------------------------|------------------|------------------|
|                           |        |        |              |                 | Model 1‡                 | Model 2§         | Model 3¶         |
| <b>All-cause dementia</b> |        |        |              |                 |                          |                  |                  |
| <18.5                     | 1,214  | 49     | 4,235        | 11.57           | 1.97 (1.36–2.85)         | 1.49 (1.03–2.15) | 1.44 (0.99–2.10) |
| ≥18.5–23                  | 33,116 | 685    | 116,472      | 5.88            | 1 (ref.)                 | 1 (ref.)         | 1 (ref.)         |
| >23–25                    | 43,071 | 655    | 152,611      | 4.29            | 0.73 (0.64–0.84)         | 0.81 (0.70–0.92) | 0.80 (0.69–0.91) |
| >25–30                    | 76,677 | 1,017  | 267,897      | 3.80            | 0.65 (0.57–0.73)         | 0.79 (0.69–0.89) | 0.77 (0.68–0.88) |
| >30                       | 13,798 | 157    | 45,888       | 3.42            | 0.59 (0.47–0.74)         | 0.79 (0.63–0.99) | 0.79 (0.63–0.99) |
| <i>P</i> for trend        |        |        |              |                 | <0.001                   | <0.001           | <0.001           |
| <b>AD</b>                 |        |        |              |                 |                          |                  |                  |
| <18.5                     | 1,214  | 39     | 4,235        | 9.21            | 2.15 (1.42–3.25)         | 1.62 (1.07–2.45) | 1.59 (1.04–2.43) |
| ≥18.5–23                  | 33,116 | 500    | 116,472      | 4.29            | 1 (ref.)                 | 1 (ref.)         | 1 (ref.)         |
| >23–25                    | 43,071 | 471    | 152,611      | 3.09            | 0.72 (0.61–0.84)         | 0.79 (0.68–0.93) | 0.79 (0.67–0.93) |
| >25–30                    | 76,677 | 722    | 267,897      | 2.70            | 0.63 (0.54–0.73)         | 0.76 (0.66–0.89) | 0.76 (0.65–0.88) |
| >30                       | 13,798 | 114    | 45,888       | 2.48            | 0.59 (0.46–0.77)         | 0.77 (0.59–1.01) | 0.78 (0.60–1.01) |
| <i>P</i> for trend        |        |        |              |                 | <0.001                   | <0.001           | <0.001           |
| <b>VD</b>                 |        |        |              |                 |                          |                  |                  |
| <18.5                     | 1,214  | 8      | 4,235        | 1.89            | 2.29 (0.91–5.75)         | 1.73 (0.69–4.35) | 1.71 (0.68–4.31) |
| ≥18.5–23                  | 33,116 | 96     | 116,472      | 0.82            | 1 (ref.)                 | 1 (ref.)         | 1 (ref.)         |
| >23–25                    | 43,071 | 94     | 152,611      | 0.62            | 0.75 (0.52–1.07)         | 0.82 (0.57–1.18) | 0.80 (0.55–1.16) |
| >25–30                    | 76,677 | 160    | 267,897      | 0.60            | 0.73 (0.53–1.00)         | 0.89 (0.64–1.23) | 0.86 (0.62–1.20) |
| >30                       | 13,798 | 23     | 45,888       | 0.50            | 0.62 (0.34–1.10)         | 0.89 (0.49–1.60) | 0.86 (0.47–1.56) |
| <i>P</i> for trend        |        |        |              |                 | 0.001                    | 0.253            | 0.276            |

Ref., reference. \*Incidence per 1,000 person-years. †Bonferroni correction was used to control the overall significance level 0.05. Actual confidence level was 98.75%. ‡Unadjusted. §Adjusted for age, sex, smoking status, alcohol consumption, physical activity, and income. ¶Adjusted for age, sex, smoking status, alcohol consumption, physical activity, income, hypertension, dyslipidemia, CKD, stroke, depressive disorder, FPG level, insulin use, and number of oral antidiabetic agents used.

Losing or gaining >10% of body weight was significantly associated with an increased risk of all-cause dementia. In addition, weight loss of >10% was

significantly associated with an increased risk of AD. These trends persisted after adjusting for confounding variables, including baseline BMI.

Previous longitudinal studies have reported the impact of type 2 diabetes on cognitive decline, mild cognitive impairment (MCI), and dementia in various

**Table 3—HR (95% CI) of incident dementia according to weight change for 2 years after diagnosis for type 2 diabetes**

| Weight change (%)         | N       | Events | Person-years | Incidence rate* | HR (95% CI) <sup>†</sup> |                  |                  |
|---------------------------|---------|--------|--------------|-----------------|--------------------------|------------------|------------------|
|                           |         |        |              |                 | Model 1‡                 | Model 2§         | Model 3¶         |
| <b>All-cause dementia</b> |         |        |              |                 |                          |                  |                  |
| ≥ -10                     | 6,603   | 214    | 22,208       | 9.64            | 2.55 (2.12–3.06)         | 1.54 (1.28–1.85) | 1.34 (1.11–1.63) |
| < -10 to ≥ -5             | 24,550  | 451    | 83,307       | 5.41            | 1.43 (1.25–1.63)         | 1.13 (0.99–1.30) | 1.08 (0.94–1.24) |
| < -5 to <5                | 115,473 | 1,555  | 404,340      | 3.85            | 1 (ref.)                 | 1 (ref.)         | 1 (ref.)         |
| ≥5 to <10                 | 15,539  | 231    | 56,695       | 4.07            | 1.04 (0.88–1.25)         | 1.14 (0.95–1.36) | 1.11 (0.93–1.33) |
| ≥10                       | 5,711   | 112    | 20,552       | 5.45            | 1.40 (1.10–1.79)         | 1.38 (1.08–1.76) | 1.38 (1.08–1.76) |
| <i>P</i>                  |         |        |              |                 | <0.001                   | <0.001           | <0.001           |
| <b>AD</b>                 |         |        |              |                 |                          |                  |                  |
| ≥ -10                     | 6,603   | 152    | 22,208       | 6.84            | 2.53 (2.04–3.14)         | 1.47 (1.18–1.83) | 1.26 (1.01–1.59) |
| < -10 to ≥ -5             | 24,550  | 349    | 83,307       | 4.19            | 1.54 (1.32–1.80)         | 1.20 (1.03–1.40) | 1.13 (0.97–1.33) |
| < -5 to <5                | 115,473 | 1,116  | 404,340      | 2.76            | 1 (ref.)                 | 1 (ref.)         | 1 (ref.)         |
| ≥5 to <10                 | 15,539  | 154    | 56,695       | 2.72            | 0.97 (0.78–1.20)         | 1.06 (0.85–1.31) | 1.04 (0.84–1.30) |
| ≥10                       | 5,711   | 75     | 20,552       | 3.65            | 1.30 (0.97–1.76)         | 1.27 (0.95–1.72) | 1.29 (0.95–1.74) |
| <i>P</i>                  |         |        |              |                 | <0.001                   | <0.001           | 0.018            |
| <b>VD</b>                 |         |        |              |                 |                          |                  |                  |
| ≥ -10                     | 6,603   | 29     | 22,208       | 1.31            | 2.22 (1.36–3.63)         | 1.53 (0.93–2.52) | 1.38 (0.83–2.31) |
| < -10 to ≥ -5             | 24,550  | 51     | 83,307       | 0.61            | 1.04 (0.71–1.53)         | 0.88 (0.60–1.31) | 0.88 (0.59–1.30) |
| < -5 to <5                | 115,473 | 241    | 404,340      | 0.60            | 1 (ref.)                 | 1 (ref.)         | 1 (ref.)         |
| ≥5 to <10                 | 15,539  | 39     | 56,695       | 0.69            | 1.14 (0.74–1.76)         | 1.24 (0.80–1.91) | 1.19 (0.76–1.84) |
| ≥10                       | 5,711   | 21     | 20,552       | 1.02            | 1.70 (0.96–3.00)         | 1.72 (0.97–3.03) | 1.65 (0.93–2.93) |
| <i>P</i>                  |         |        |              |                 | <0.001                   | 0.017            | 0.056            |

Ref., reference. \*Incidence per 1,000 person-years. †Bonferroni correction was used to control the overall significance level 0.05. Actual confidence level was 98.75%. ‡Unadjusted. §Adjusted for age, sex, smoking status, alcohol consumption, physical activity, and income. ¶Adjusted for age, sex, smoking status, alcohol consumption, physical activity, income, hypertension, dyslipidemia, CKD, stroke, depressive disorder, FPG level, insulin use, number of oral antidiabetic agents used, and baseline BMI.

**Table 4—HR (95% CI) of dementia by weight change status in subgroups according to age, sex, and obesity status**

| Subgroup                     | Weight change (%) | All-cause dementia |       |                  | AD     |       |                  | VD     |      |                  |
|------------------------------|-------------------|--------------------|-------|------------------|--------|-------|------------------|--------|------|------------------|
|                              |                   | Events             | IR    | HR (95% CI)*     | Events | IR    | HR (95% CI)*     | Events | IR   | HR (95% CI)*     |
| <b>Age, years</b>            |                   |                    |       |                  |        |       |                  |        |      |                  |
| <65                          | ≥ -5              | 103                | 1.31  | 1.23 (0.94–1.61) | 62     | 0.79  | 1.08 (0.76–1.52) | 25     | 0.32 | 1.67 (0.96–2.92) |
|                              | -5 to 5           | 303                | 0.95  | 1 (ref.)         | 197    | 0.62  | 1 (ref.)         | 60     | 0.19 | 1 (ref.)         |
| ≥65                          | ≥5                | 70                 | 1.12  | 1.22 (0.90–1.65) | 37     | 0.59  | 1.00 (0.66–1.51) | 21     | 0.34 | 1.83 (1.02–3.27) |
|                              | ≥ -5              | 562                | 20.86 | 1.13 (1.00–1.28) | 439    | 16.30 | 1.18 (1.03–1.36) | 55     | 2.04 | 0.84 (0.59–1.21) |
|                              | -5 to 5           | 1,252              | 14.62 | 1 (ref.)         | 919    | 10.73 | 1 (ref.)         | 181    | 2.11 | 1 (ref.)         |
|                              | ≥5                | 273                | 18.64 | 1.19 (1.02–1.38) | 192    | 13.11 | 1.14 (0.96–1.37) | 39     | 2.66 | 1.15 (0.77–1.72) |
| <i>P</i> for interaction     |                   |                    |       | 0.677            |        |       | 0.740            |        |      | 0.030            |
| <b>Sex</b>                   |                   |                    |       |                  |        |       |                  |        |      |                  |
| Men                          | ≥ -5              | 225                | 3.98  | 1.03 (0.86–1.24) | 165    | 2.92  | 1.05 (0.85–1.30) | 31     | 0.55 | 0.90 (0.57–1.43) |
|                              | -5 to 5           | 722                | 2.72  | 1 (ref.)         | 491    | 1.85  | 1 (ref.)         | 131    | 0.49 | 1 (ref.)         |
|                              | ≥5                | 183                | 3.37  | 1.22 (1.01–1.47) | 104    | 1.91  | 1.02 (0.80–1.30) | 47     | 0.87 | 1.70 (1.16–2.51) |
| Women                        | ≥ -5              | 440                | 8.98  | 1.20 (1.05–1.38) | 336    | 6.85  | 1.22 (1.04–1.43) | 49     | 1.00 | 1.03 (0.69–1.55) |
|                              | -5 to 5           | 833                | 6.02  | 1 (ref.)         | 625    | 4.51  | 1 (ref.)         | 110    | 0.79 | 1 (ref.)         |
|                              | ≥5                | 160                | 6.98  | 1.13 (0.93–1.37) | 125    | 5.45  | 1.18 (0.94–1.47) | 13     | 0.57 | 0.73 (0.38–1.41) |
| <i>P</i> for interaction     |                   |                    |       | 0.369            |        |       | 0.574            |        |      | 0.020            |
| <b>BMI, kg/m<sup>2</sup></b> |                   |                    |       |                  |        |       |                  |        |      |                  |
| <25                          | ≥ -5              | 488                | 6.84  | 1.09 (0.95–1.25) | 367    | 5.15  | 1.10 (0.94–1.28) | 54     | 0.76 | 0.88 (0.60–1.29) |
|                              | -5 to 5           | 843                | 4.56  | 1 (ref.)         | 612    | 3.31  | 1 (ref.)         | 127    | 0.69 | 1 (ref.)         |
|                              | ≥5                | 162                | 5.59  | 1.17 (0.96–1.42) | 109    | 3.76  | 1.09 (0.86–1.38) | 31     | 1.07 | 1.46 (0.93–2.30) |
| ≥25                          | ≥ -5              | 177                | 5.18  | 1.28 (1.05–1.56) | 134    | 3.92  | 1.34 (1.08–1.66) | 26     | 0.76 | 1.31 (0.80–2.14) |
|                              | -5 to 5           | 712                | 3.24  | 1 (ref.)         | 504    | 2.30  | 1 (ref.)         | 114    | 0.52 | 1 (ref.)         |
|                              | ≥5                | 181                | 3.75  | 1.20 (0.99–1.45) | 120    | 2.49  | 1.13 (0.89–1.43) | 29     | 0.60 | 1.20 (0.75–1.93) |
| <i>P</i> for interaction     |                   |                    |       | 0.111            |        |       | 0.101            |        |      | 0.164            |

IR, incidence rate; ref., reference. \*HR (Bonferroni-adjusted 95% CI) was calculated using a multivariable Cox hazard regression analysis after adjusting for age, sex, smoking status, alcohol consumption, physical activity, income, hypertension, dyslipidemia, CKD, stroke, depressive disorder, FPG level, insulin use, number of oral antidiabetic agents used, and baseline BMI. The actual confidence level was 97.5%.

Downloaded from <http://diabetesjournals.org/care/article-pdf/42/7/1217/530364/dc181667.pdf> by guest on 03 December 2023

populations worldwide. However, the underlying mechanism seems to be complex and remains unclear. Diabetes has been found to increase the risk of cerebral infarction, but whether diabetes lowers the threshold to manifesting dementia through cerebrovascular disease or whether diabetes increases the risk for AD pathophysiology is unclear (9,10). Furthermore, a recent review suggested that the diabetes-related processes may contribute to the development of dementia but not directly to the hallmark pathologies of AD (22).

In this study, we found that the baseline BMI on diagnosis of type 2 diabetes was inversely associated with the risk of developing all-cause dementia or AD but not VD. The relationship between baseline BMI and dementia risk has been poorly studied in patients with diabetes. Several previous cohort studies in Western countries have explored the impact of BMI on the risk of dementia and produced conflicting results: high and low BMIs have both been reported to increase the risk of dementia (11–14). In general, however, low BMIs have been

associated with high rates of dementia, such that the optimal BMI for reducing dementia risk is within the overweight range (16). Moreover, a low baseline BMI has been linked with faster cognitive decline in MCI (23,24). Developing countries tend to have lower average BMIs compared with Western countries, and a Nigerian cohort study also reported low BMI was a risk factor for dementia (25).

Although the underlying mechanism is not fully understood, a low BMI may accompany changes in appetite regulation that precede overt dementia (26). Moreover, those with a low BMI have reported feelings of apathy that are a common prodromal sign of early dementia, especially in patients with MCI rapidly progressing to overt dementia (16,27). Our results support a previously demonstrated inverse association between baseline BMI and risk of dementia and include evidence of a similar trend in patients with incident diabetes. Of note, we found that baseline BMI was not associated with VD, suggesting that different pathophysiologic mechanisms are involved.

Our study found significant U-shaped relationships between weight change and all-cause dementia, and weight change >10% increased the risk of all-cause dementia. Furthermore, weight loss >10% was associated with increased risk of AD in patients with incident diabetes. While there is limited evidence regarding the impact of weight change on the risk of dementia in patients with diabetes, several other studies have similar findings. Epidemiologic studies have generally reported the effect of declining BMI and AD risk (28). A U.S. nationwide cohort study of elderly inpatients reported that weight loss increased dementia risk by 1.26 times and that the inpatient population was expected to have more comorbidities and to have diabetes (29). A randomized, controlled lifestyle intervention study of overweight Finnish patients with impaired glucose tolerance reported a decline in cognitive performance with decreasing BMI, although this may have reflected reverse causality (30).

Pathophysiologic pathways linking weight change and dementia may be mediated by cardiovascular risk factors,

insulin resistance, or lifestyle. Recent studies demonstrate a correlation between brain atrophy and rapid loss of lean body mass. Patients with incident diabetes are likely to maintain strict glycemic control and lose weight, which can lead to hypoglycemia and weakness that is associated with dementia (31). Weight loss in developing countries is often a consequence of low caloric intake that is insufficient for the demands of physical activity. Gradual weight loss might precede a diagnosis of dementia while being viewed as forgetting to eat, apathy and loss of initiative, impaired olfactory function, loss of taste, and difficulty in swallowing. Weight loss and the prodromal stages of dementia are likely to affect one another adversely (32,33).

Meanwhile, a 1.6-year follow-up study for individuals with MCI showed that regardless of baseline BMI, significant weight gain or loss increased the risk of progressing to probable AD (34). Also, intentional weight loss by obese and overweight individuals has a positive effect on cognitive function (35). Intentional weight loss has been correlated with cognitive improvement even in obese elderly patients with MCI (36). These studies support our findings that excessive weight gain significantly increases the risk of all-cause dementia in patients with type 2 diabetes. Although our study could not assess whether weight change was intentional, excessive weight gain is expected to have a detrimental effect on the link between type 2 diabetes and cognitive decline. Altered energy homeostasis can affect disease progression in dementia, and neurodegenerative diseases are also related to metabolic changes such as weight gain or loss (37). Of note, our study found that the correlation between weight change and VD risk was stronger in men than in women, implying that weight gain affects cognitive function in men with diabetes; however, further studies are needed to understand this relationship.

Our findings should be interpreted with several limitations in mind. First, the dementia diagnoses based on ICD-10-CM might be under- or overestimated, and we could not determine dementia severity from the medical records. Although we excluded patients with a dementia diagnosis made before the index date, there may have been reverse causality. Second, we could not consider whether weight change was intentional

or unintentional. Although the initial glycemic status can affect individual weight, the baseline FPG levels were not high enough to do so. We also corrected for the baseline FPG level in the analysis. Third, the observational duration of weight change after the diagnosis of type 2 diabetes and the follow-up duration for dementia development from the index date might not have been long enough to fully assess relationships. Fourth, the response rate of the self-report questionnaire in the South Korean NHIS was ~70%, and, therefore, there may have been possible selection bias. Fifth, our study included only a Korean population, and thus the findings cannot be generalized to other ethnicities.

Despite these limitations, the strengths of our study include a large sample size, longitudinal design, and abundant data regarding demographics, socioeconomics, biomedical information, comorbidity, and lifestyle variables. To our knowledge, this is the first study to examine the impact of baseline BMI and weight change on the risk of dementia in patients with new-onset type 2 diabetes. Additionally, new-onset diabetes is considered less affected by complications or other comorbidities that further effect weight change.

In conclusion, this large-scale cohort study of the South Korean population demonstrated that baseline BMI was inversely associated with the risk of all-cause dementia and AD among patients with new-onset type 2 diabetes. Furthermore, weight loss and weight gain after diagnosis were associated with increased risks of all-cause dementia. Weight loss was positively correlated with AD risk. In consideration of the risk of dementia, we suggest close monitoring for body weight in patients with diabetes.

**Funding.** This research was supported by Basic Science Research Program through the National Research Foundation of Korea funded by the Ministry of Education (2018R1D1A1B07049079).

**Duality of Interest.** No potential conflicts of interest relevant to this article were reported.

**Author Contributions.** G.E.N., Y.G.P., K.H., M.K.K., E.S.Ko., E.S.Ki., M.-K.L., B.K., O.-K.H., and H.-S.K. gave final approval of the version to be published. G.E.N., Y.G.P., and H.-S.K. contributed to the study concept and design, acquisition, analysis, and interpretation of data, drafting of the manuscript, and critical revision of the manuscript. K.H. contributed to the study

concept and design, acquisition, analysis and interpretation of data, and critical revision of the manuscript. M.K.K., E.S.Ko., E.S.Ki., M.-K.L., and O.-K.H. contributed to the interpretation of data and critical revision of the manuscript. B.K. contributed to the acquisition, analysis, and interpretation of data and critical revision of the manuscript. Y.G.P. and H.-S.K. are the guarantors of this work and, as such, had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

## References

- Cheng D, Noble J, Tang MX, Schupf N, Mayeux R, Luchsinger JA. Type 2 diabetes and late-onset Alzheimer's disease. *Dement Geriatr Cogn Disord* 2011;31:424–430
- Snowden MB, Steinman LE, Bryant LL, et al. Dementia and co-occurring chronic conditions: a systematic literature review to identify what is known and where are the gaps in the evidence? *Int J Geriatr Psychiatry* 2017;32:357–371
- Grodstein F, Chen J, Wilson RS, Manson JE; Nurses' Health Study. Type 2 diabetes and cognitive function in community-dwelling elderly women. *Diabetes Care* 2001;24:1060–1065
- Yaffe K, Blackwell T, Kanaya AM, Davidowitz N, Barrett-Connor E, Krueger K. Diabetes, impaired fasting glucose, and development of cognitive impairment in older women. *Neurology* 2004;63:658–663
- Roberts RO, Geda YE, Knopman DS, et al. Association of duration and severity of diabetes mellitus with mild cognitive impairment. *Arch Neurol* 2008;65:1066–1073
- Peila R, Rodriguez BL, Launer LJ; Honolulu-Asia Aging Study. Type 2 diabetes, APOE gene, and the risk for dementia and related pathologies: the Honolulu-Asia Aging Study. *Diabetes* 2002;51:1256–1262
- Luchsinger JA, Tang MX, Stern Y, Shea S, Mayeux R. Diabetes mellitus and risk of Alzheimer's disease and dementia with stroke in a multiethnic cohort. *Am J Epidemiol* 2001;154:635–641
- Xu WL, Qiu CX, Wahlin A, Winblad B, Fratiglioni L. Diabetes mellitus and risk of dementia in the Kungsholmen project: a 6-year follow-up study. *Neurology* 2004;63:1181–1186
- Luchsinger JA. Adiposity, hyperinsulinemia, diabetes and Alzheimer's disease: an epidemiological perspective. *Eur J Pharmacol* 2008;585:119–129
- Luchsinger JA. Type 2 diabetes and cognitive impairment: linking mechanisms. *J Alzheimers Dis* 2012;30(Suppl. 2):S185–S198
- Whitmer RA, Gunderson EP, Quesenberry CP Jr., Zhou J, Yaffe K. Body mass index in midlife and risk of Alzheimer disease and vascular dementia. *Curr Alzheimer Res* 2007;4:103–109
- Fitzpatrick AL, Kuller LH, Lopez OL, et al. Midlife and late-life obesity and the risk of dementia: cardiovascular health study. *Arch Neurol* 2009;66:336–342
- Anstey KJ, Cherbuin N, Budge M, Young J. Body mass index in midlife and late-life as a risk factor for dementia: a meta-analysis of prospective studies. *Obes Rev* 2011;12:e426–e437
- Sobów T, Fendler W, Magierski R. Body mass index and mild cognitive impairment-to-dementia progression in 24 months: a prospective study. *Eur J Clin Nutr* 2014;68:1216–1219

15. Gustafson D. Adiposity indices and dementia. *Lancet Neurol* 2006;5:713–720
16. García-Ptacek S, Faxén-Irving G, Cermáková P, Eriksdotter M, Religa D. Body mass index in dementia. *Eur J Clin Nutr* 2014;68:1204–1209
17. Van Gaal L, Scheen A. Weight management in type 2 diabetes: current and emerging approaches to treatment. *Diabetes Care* 2015;38:1161–1172
18. WHO Expert Consultation. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies [published correction appears in *Lancet* 2004;363:902]. *Lancet* 2004;363:157–163
19. Agarwal DP. Cardioprotective effects of light-moderate consumption of alcohol: a review of putative mechanisms. *Alcohol Alcohol* 2002;37:409–415
20. Levey AS, Stevens LA, Schmid CH, et al.; CKD-EPI (Chronic Kidney Disease Epidemiology Collaboration). A new equation to estimate glomerular filtration rate [published correction appears in *Ann Intern Med* 2011;155:408]. *Ann Intern Med* 2009;150:604–612
21. Levey AS, Coresh J, Balk E, et al.; National Kidney Foundation. National Kidney Foundation practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Ann Intern Med* 2003;139:137–147
22. Biessels GJ, Despa F. Cognitive decline and dementia in diabetes mellitus: mechanisms and clinical implications. *Nat Rev Endocrinol* 2018;14:591–604
23. Cronk BB, Johnson DK, Burns JM; Alzheimer's Disease Neuroimaging Initiative. Body mass index and cognitive decline in mild cognitive impairment. *Alzheimer Dis Assoc Disord* 2010;24:126–130
24. Besser LM, Gill DP, Monsell SE, et al. Body mass index, weight change, and clinical progression in mild cognitive impairment and Alzheimer disease. *Alzheimer Dis Assoc Disord* 2014;28:36–43
25. Ogunniyi A, Gao S, Unverzagt FW, et al. Weight loss and incident dementia in elderly Yoruba Nigerians: a 10-year follow-up study. *Int Psychogeriatr* 2011;23:387–394
26. Fadel JR, Jolivald CG, Reagan LP. Food for thought: the role of appetitive peptides in age-related cognitive decline. *Ageing Res Rev* 2013;12:764–776
27. Benoit M, Berrut G, Doussaint J, et al. Apathy and depression in mild Alzheimer's disease: a cross-sectional study using diagnostic criteria. *J Alzheimers Dis* 2012;31:325–334
28. Gu Y, Scarmeas N, Cosentino S, et al. Change in body mass index before and after Alzheimer's disease onset. *Curr Alzheimer Res* 2014;11:349–356
29. Beydoun MA, Beydoun HA, Gamaldo AA, et al. Nationwide inpatient prevalence, predictors, and outcomes of Alzheimer's disease among older adults in the United States, 2002–2012. *J Alzheimers Dis* 2015;48:361–375
30. Lehtisalo J, Lindström J, Ngandu T, et al.; Finnish Diabetes Prevention Study. Association of long-term dietary fat intake, exercise, and weight with later cognitive function in the Finnish Diabetes Prevention Study. *J Nutr Health Aging* 2016;20:146–154
31. Abdelhafiz AH, McNicholas E, Sinclair AJ. Hypoglycemia, frailty and dementia in older people with diabetes: reciprocal relations and clinical implications. *J Diabetes Complications* 2016;30:1548–1554
32. Suemoto CK, Gilsanz P, Mayeda ER, Glymour MM. Body mass index and cognitive function: the potential for reverse causation. *Int J Obes* 2015;39:1383–1389
33. Ready RE, Ott BR, Grace J, Cahn-Weiner DA. Apathy and executive dysfunction in mild cognitive impairment and Alzheimer disease. *Am J Geriatr Psychiatry* 2003;11:222–228
34. Ye BS, Jang EY, Kim SY, et al. Unstable body mass index and progression to probable Alzheimer's disease dementia in patients with amnesic mild cognitive impairment. *J Alzheimers Dis* 2016;49:483–491
35. Siervo M, Arnold R, Wells JC, et al. Intentional weight loss in overweight and obese individuals and cognitive function: a systematic review and meta-analysis. *Obes Rev* 2011;12:968–983
36. Horie NC, Serrao VT, Simon SS, et al. Cognitive effects of intentional weight loss in elderly obese individuals with mild cognitive impairment. *J Clin Endocrinol Metab* 2016;101:1104–1112
37. Vercauteren P, Vieau D, Blum D, Petersén Å, Dupuis L. Hypothalamic alterations in neurodegenerative diseases and their relation to abnormal energy metabolism. *Front Mol Neurosci* 2018;11:2