



# Chronic Kidney Disease, Diabetes, and Risk of Mortality After Acute Myocardial Infarction: Insight From the FAST-MI Program

*Diabetes Care* 2020;43:e43–e44 | <https://doi.org/10.2337/dc19-2209>

Louis Potier,<sup>1</sup> Ronan Roussel,<sup>1</sup>  
Marianne Zeller,<sup>2</sup> François Schiele,<sup>3</sup>  
Etienne Puymirat,<sup>4</sup>  
Tabassome Simon,<sup>5,6,7</sup> and  
Nicolas Danchin<sup>4</sup>

Diabetes is associated with a substantially increased risk of all-cause death, mainly driven by cardiovascular (CV) mortality. Furthermore, diabetes is associated with poorer outcomes after acute myocardial infarction (AMI) (1). Impaired glomerular filtration rate (GFR) is also associated with an increased risk of CV mortality (2). However, whether diabetes still confers a higher risk of mortality in patients with impaired GFR remains unknown.

The aim of this study was to assess the long-term prognostic significance of both diabetes and renal impairment in two prospective nationwide cohorts of AMI patients: FAST-MI (French Registry of Acute ST-Elevation or non-ST-elevation Myocardial Infarction) 2005 ( $n = 3,670$  [reg. no. NCT00673036, ClinicalTrials.gov]) and FAST-MI 2010 ( $n = 4,169$  [NCT01237418]) (3). Both registries consecutively included patients with AMI admitted to cardiac intensive care units within 48 h of symptom onset during a specified 1-month period. AMI was defined by increased levels of cardiac biomarkers together with either compatible symptoms or electrocardiogram changes. Vital

status at 5 years was available in >95%. We assessed all-cause mortality at 5 years according to estimated GFR (eGFR) (calculated with Chronic Kidney Disease Epidemiology Collaboration [CKD-EPI] formula and based on KDIGO [Kidney Disease: Improving Global Outcomes] GFR categories with eGFR <30 and <15 mL/min/1.73 m<sup>2</sup> pooled) and diabetes status at inclusion. Multivariable proportional hazards models (assumptions checked) were used with covariates chosen based on their potential prognostic relevance: year of inclusion; sex; age; BMI categories; hypertension; current smoking; prior AMI; peripheral artery disease; history of heart failure, stroke, cancer, and chronic obstructive pulmonary disease; type of myocardial infarction (MI) (STEMI or NSTEMI); GRACE risk score; percutaneous coronary intervention or coronary artery bypass during hospitalization; and left ventricular ejection fraction.

Among the 7,839 participants, 7,656 had creatinine values, assessed before percutaneous coronary intervention, at entry available. Mean  $\pm$  SD age was  $66.1 \pm 14.2$

years, 2,230 (29.1%) were female, and 2,090 (27.3%) had diabetes, with a duration of diabetes >10 years in 30.0% (missing data 35.6%). Mean  $\pm$  SD HbA<sub>1c</sub> at entry was  $7.5 \pm 1.5\%$ , and eGFR was <30 mL/min/1.73 m<sup>2</sup> in 398 (5.2%), 30–45 in 688 (9.0%), 45–60 in 1,151 (15.0%), 60–90 in 3,328 (43.5%), and >90 in 2,091 (27.3%). Five-year mortality was 38% ( $n = 796$ ) in patients with diabetes versus 19% ( $n = 1,058$ ) in patients without diabetes (adjusted hazard ratio [HR] 1.47, 95% CI 1.33–1.62,  $P < 0.001$ ). According to eGFR, among all participants, 5-year mortality rates were 8.1% ( $n = 170$ ) for eGFR >90 mL/min/1.73 m<sup>2</sup>, 17.7% (588) for eGFR 60–90, 36.4% ( $n = 419$ ) for eGFR 45–60, 57.6% ( $n = 396$ ) for eGFR 30–45, and 70.6% ( $n = 281$ ) for eGFR <30. Multivariable analysis suggested a gradual increase in mortality with decreasing renal function: adjusted HR versus eGFR >90 mL/min/1.73 m<sup>2</sup> as reference 0.91 (95% CI 0.75–1.11,  $P = 0.36$ ) for eGFR 60–90 mL/min/1.73 m<sup>2</sup>; 1.19 (0.96–1.46,  $P = 0.11$ ) for eGFR 45–60; 1.48 (1.19–1.85,  $P = 0.001$ ) for eGFR 30–45, and 1.86 (1.48–2.35,  $P < 0.001$ ) for eGFR <30.

<sup>1</sup>Department of Diabetology, Endocrinology and Nutrition, Hôpital Bichat, Assistance Publique–Hôpitaux de Paris, and Cordeliers Research Center, INSERM, U-1138, Université de Paris, Paris, France

<sup>2</sup>Equipe PEC2, EA 7460, UFR Sciences de Santé, Université de Bourgogne Franche Comté, Dijon, France

<sup>3</sup>Department of Cardiology, University Hospital Jean Minjot, Besançon, France

<sup>4</sup>Department of Cardiology, Hôpital Européen Georges Pompidou, Assistance Publique–Hôpitaux de Paris, and INSERM U-970, Université de Paris, Paris, France

<sup>5</sup>Department of Clinical Pharmacology and Clinical Research Platform (URCEST-CRC-CRB), Hôpital Saint-Antoine, Assistance Publique–Hôpitaux de Paris, Paris, France


<sup>6</sup>French Alliance for Cardiovascular Clinical Trials, Paris, France

<sup>7</sup>Faculté de Médecine Pierre et Marie Curie, Sorbonne Université, Paris, France

Corresponding author: Louis Potier, [louis.potier@gmail.com](mailto:louis.potier@gmail.com)

Received 3 November 2019 and accepted 21 December 2019

© 2020 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <https://www.diabetesjournals.org/content/license>.



**Figure 1**—HR for 5-year mortality according to eGFR and diabetes. *A*: HR according to eGFR in participants without and participants with diabetes (reference eGFR >90 mL/min/1.73 m<sup>2</sup>, with separate reference groups in the diabetes and no diabetes strata). *B*: HR for presence of diabetes compared with no diabetes within each renal function category.

Stratified analysis showed that increasing renal impairment was associated with an increased risk of death for participants with diabetes under a threshold of 60 mL/min/1.73 m<sup>2</sup> but only below a threshold of 45 mL/min/1.73 m<sup>2</sup> for participants without diabetes (Fig. 1A).

Compared with no diabetes, diabetes was associated with an increased risk of 5-year death throughout eGFR categories, except for eGFR <30 mL/min/1.73 m<sup>2</sup>: HR 1.45 (95% CI 1.01–2.09) for eGFR >90, HR 1.54 (1.28–1.84, *P* < 0.001) for eGFR 60–90, HR 1.64 (1.33–2.02, *P* < 0.001) for eGFR 45–60, HR 1.28 (1.04–1.59, *P* = 0.02) for eGFR 30–45, and HR 1.19 (0.92–1.55, *P* = 0.19) for eGFR <30 (Fig. 1B).

The trend of a lower prognostic impact of diabetes in patients with the most severe chronic kidney disease was observed both in men and in women, with no interaction between eGFR categories and diabetes (*P* = 0.52); likewise, the interaction between eGFR <30 vs. ≥30 mL/min/1.73 m<sup>2</sup> and diabetes was not significant (*P* = 0.17).

The study is observational, with some missing data such as proteinuria in all patients and HbA<sub>1c</sub> in some patients. Also, the group with the lowest eGFR was comparatively small. Finally, as we wished to describe long-term follow-up, we only included patients from the 2005 and 2010 cohorts. Since then, newer glucose-lowering medications have documented an impact on CV and renal outcomes, which might modify the relationship among diabetes, eGFR, and clinical outcomes.

Thus, the risk of 5-year all-cause mortality progressively increases with declining renal function in AMI patients; the eGFR threshold below which mortality increases, however, appears to be higher in patients with diabetes than in patients without diabetes (60 vs. 45 mL/min/1.73 m<sup>2</sup>). Likewise, diabetes is associated with higher mortality, although the increased risk of mortality associated with diabetes is attenuated for patients with eGFR <45 mL/min/1.73 m<sup>2</sup> and no longer significant for those with eGFR <30 mL/min/1.73 m<sup>2</sup>.

These results suggest that in post-MI patients, 1) chronically impaired renal failure and diabetes are both associated with an increased risk of mortality, 2) renal function requires specific attention in patients with diabetes as soon as it is mildly impaired (45–60 mL/min/1.73 m<sup>2</sup>), and 3) tight glycemic control, which is controversial in post-MI patients (4), may not be essential in patients with diabetes with eGFR <30 mL/min/1.73 m<sup>2</sup>.

**Acknowledgments.** The authors are deeply indebted to all patients who accepted to participate in the surveys and to the physicians who took care of the patients at the participating institutions.

**Funding.** This study was funded by the French Society of Cardiology.

**Duality of Interest.** The French Society of Cardiology received grants supporting the FAST-MI program from Amgen, AstraZeneca, Bayer, Bristol-Myers Squibb (BMS), Boehringer Ingelheim, Daiichi-Sankyo, Eli Lilly, Merck Sharp & Dohme (MSD), Pfizer, and Sanofi. None of the companies had a role in the design or conduct of the study, data collection, or management.

They were not involved in the analysis or interpretation of the data or in the preparation, review, or approval of the manuscript. L.P. reports grants, personal fees, and nonfinancial support from Novo Nordisk and Sanofi; personal fees and nonfinancial support from Eli Lilly; and nonfinancial support from Servier. R.R. reports grants, personal fees, and nonfinancial support from Sanofi; personal fees and nonfinancial support from MSD; grants from Amgen; personal fees from Physiogenex, AstraZeneca, Janssen, Eli Lilly, Abbott, Medtronic, Novo Nordisk, and Servier; and grants from Novo Nordisk. F.S. reports personal fees from Amgen, AstraZeneca, Bayer, BMS, MSD, Pfizer, and Sanofi. E.P. reports fees for lectures and/or consulting from Amgen, AstraZeneca, Bayer, BIOTRONIK, BMS, Boehringer Ingelheim, Daiichi-Sankyo, Lilly, MSD, The Medicines Company, Sanofi, Saint Jude Medical, Servier, and Siemens. T.S. reports grants from AstraZeneca, Daiichi-Sankyo, Eli Lilly, GlaxoSmithKline, MSD, Novartis, and Sanofi and personal fees for board membership and/or consultancy and/or lectures from AstraZeneca, BMS, Sanofi, and Novartis. N.D. reports having received grants, speaking fees, consulting fees, or nonfinancial support from Amgen, AstraZeneca, Bayer, BMS, Boehringer Ingelheim, Intercept, Novo Nordisk, Pfizer, Sanofi, and Servier. No other potential conflicts of interest relevant to this article were reported.

**Author Contributions.** M.Z., F.S., T.S., and N.D. researched data. L.P., R.R., and N.D. wrote the manuscript. M.Z., F.S., E.P., and T.S. reviewed and edited the manuscript and contributed to the discussion. L.P. and N.D. are the guarantors of this work and, as such, had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

## References

- Piccolo R, Franzone A, Koskinas KC, et al. Effect of diabetes mellitus on frequency of adverse events in patients with acute coronary syndromes undergoing percutaneous coronary intervention. *Am J Cardiol* 2016;118:345–352
- Matsushita K, Coresh J, Sang Y, et al.; CKD Prognosis Consortium. Estimated glomerular filtration rate and albuminuria for prediction of cardiovascular outcomes: a collaborative meta-analysis of individual participant data. *Lancet Diabetes Endocrinol* 2015;3:514–525
- Puymirat E, Simon T, Cayla G, et al.; USIK, USIC 2000, and FAST-MI Investigators. Acute myocardial infarction: changes in patient characteristics, management, and 6-month outcomes over a period of 20 years in the FAST-MI program (French Registry of Acute ST-Elevation or Non-ST-Elevation Myocardial Infarction) 1995 to 2015. *Circulation* 2017;136:1908–1919
- Heller SR, Bergenstal RM, White WB, et al.; EXAMINE Investigators. Relationship of glycated haemoglobin and reported hypoglycaemia to cardiovascular outcomes in patients with type 2 diabetes and recent acute coronary syndrome events: the EXAMINE trial. *Diabetes Obes Metab* 2017;19:664–671